

Lecture 5

Multivariate Linear Regression

Readings: Johnson & Wichern 2007, Chapter 7; DSA 8020 Lectures 1-4 [\[Link\]](#); Zelterman, 2015, Chapter 9

DSA 8070 Multivariate Analysis

Whitney Huang
Clemson University

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction
Motor Trend Car
Road Tests Data
Analysis

5.1

Notes

Agenda

- 1 Motivation
- 2 Model and Assumptions
- 3 Parameter Estimation
- 4 Inference and Prediction
- 5 Motor Trend Car Road Tests Data Analysis

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction
Motor Trend Car
Road Tests Data
Analysis

5.2

Notes

Example: Motor Trend Car Road Tests

```
> head(mtcars)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1
Hornet 4 Drive	21.4	6	258	110	3.08	3.215	19.44	1	0	3	1
Hornet Sportabout	18.7	8	360	175	3.15	3.440	17.02	0	0	3	2
Valiant	18.1	6	225	105	2.76	3.460	20.22	1	0	3	1

Suppose we would like to study the (linear) relationship between mpg, disp, hp, wt (responses) and cyl, am, carb (predictors)

Multiple regression predicts one outcome; multivariate regression predicts several simultaneously

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction
Motor Trend Car
Road Tests Data
Analysis

5.3

Notes

Why Multivariate Regression Instead of Separate Regressions?

- **Estimation:**
 - Coefficient estimates are the same as running separate regressions
- **Inference:** The real gain comes from joint modeling
 - Test hypotheses across multiple outcomes simultaneously
 - Accounts for correlations among responses \Rightarrow more powerful and accurate tests
- **Examples:**
 - Does a predictor affect all outcomes jointly?
 - Multivariate analog of ANOVA (MANOVA)

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.4

Notes

Review: Linear Regression Model

The multiple linear regression model has the form:

$$y_i = \beta_0 + \sum_{j=1}^p \beta_j x_{ij} + \varepsilon_i, \quad i = 1, \dots, n,$$

where

- y_i is the **response** for the i -th observation
- x_{ij} is the j -th **predictor** for the i -th observation
- β_0 and β_j 's are the **regression intercept** and **slopes** for the response, respectively
- ε_i is the **error** term for the response of the i -th observation

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.5

Notes

The Multivariate Linear Regression Model: Scalar Form

The multivariate (multiple) linear regression model has the form:

$$y_{ik} = \beta_{0k} + \sum_{j=1}^p \beta_{jk} x_{ij} + \varepsilon_{ik}, \quad i = 1, \dots, n, \quad k = 1, \dots, d,$$

where

- y_{ik} is the k -th **response** for the i -th observation
- x_{ij} is the j -th **predictor** for the i -th observation
- β_{0k} and β_{jk} 's are the **regression intercept** and **slopes** for k -th response, respectively
- ε_{ik} is the **error** term for the k -th response of the i -th observation

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.6

Notes

The Multivariate Linear Regression Model: Assumptions

The assumptions of the model are:

- Relationship between $\{x_j\}_{j=1}^p$ and Y_k is linear for each $k \in \{1, \dots, d\}$
- $(\varepsilon_{i1}, \dots, \varepsilon_{id})^T \stackrel{i.i.d.}{\sim} N(\mathbf{0}, \Sigma)$ is an unobserved random vector
- $[Y_{ik}|x_{i1}, \dots, x_{ip}] \sim N(\beta_{0k} + \sum_{j=1}^p \beta_{jk}x_{ij}, \sigma_{kk})$ for each $k \in \{1, \dots, d\}$

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.7

Notes

The Multivariate Linear Regression Model: Matrix Form

The multivariate multiple linear regression model has the form

$$Y = XB + E,$$

where

- $Y = [y_1, \dots, y_d]$ is the $n \times d$ response matrix, where $y_k = (y_{1k}, \dots, y_{nk})^T$ is the k -th response vector
- $X = [\mathbf{1}, x_1, \dots, x_p]$ is the $n \times (p + 1)$ design matrix
- $B = [\beta_1, \dots, \beta_d]$ is the $(p + 1) \times d$ matrix of regression coefficients
- $E = [\varepsilon_1, \dots, \varepsilon_d]$ is the $n \times d$ error matrix

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.8

Notes

Another Look of the Matrix Form

Matrix form writes the multivariate linear regression model for all $n \times d$ points simultaneously as

$$Y = XB + E$$

$$\begin{bmatrix} y_{11} & \cdots & y_{1d} \\ y_{21} & \cdots & y_{2d} \\ \vdots & \ddots & \vdots \\ y_{n1} & \cdots & y_{nd} \end{bmatrix} = \begin{bmatrix} 1 & \cdots & x_{1p} \\ 1 & \cdots & x_{2p} \\ \vdots & \ddots & \vdots \\ 1 & \cdots & x_{np} \end{bmatrix} \begin{bmatrix} \beta_{01} & \cdots & \beta_{0d} \\ \beta_{11} & \cdots & \beta_{1d} \\ \vdots & \ddots & \vdots \\ \beta_{p1} & \cdots & \beta_{pd} \end{bmatrix} + \begin{bmatrix} \varepsilon_{11} & \cdots & \varepsilon_{1d} \\ \varepsilon_{21} & \cdots & \varepsilon_{2d} \\ \vdots & \ddots & \vdots \\ \varepsilon_{n1} & \cdots & \varepsilon_{nd} \end{bmatrix}$$

Assuming that n subjects are independent, we have

- $\varepsilon_k \sim N(0, \sigma_{kk}), \quad k \in \{1, \dots, d\}$
- $\varepsilon_i \stackrel{i.i.d.}{\sim} N(\mathbf{0}, \Sigma), \quad i = 1, \dots, n$

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.9

Notes

Ordinary Least Squares

The **ordinary least squares** OLS estimate is

$$\underset{\mathbf{B} \in \mathbb{R}^{(p+1) \times d}}{\operatorname{argmin}} \|\mathbf{Y} - \mathbf{X}\mathbf{B}\|^2 = \underset{\mathbf{B} \in \mathbb{R}^{(p+1) \times d}}{\operatorname{argmin}} \sum_{i=1}^n \sum_{k=1}^d \left(y_{ik} - \beta_{0k} - \sum_{j=1}^p \beta_{jk} x_{ij} \right)^2,$$

where $\|\cdot\|$ denotes the Frobenius norm.

- $\text{OLS}(\mathbf{B}) = \|\mathbf{Y} - \mathbf{X}\mathbf{B}\|^2 = \text{tr}(\mathbf{Y}^T \mathbf{Y}) - 2\text{tr}(\mathbf{Y}^T \mathbf{X}\mathbf{B}) + \text{tr}(\mathbf{B}^T \mathbf{X}^T \mathbf{X}\mathbf{B})$
- $\frac{\partial \text{OLS}(\mathbf{B})}{\partial \mathbf{B}} = -2\mathbf{X}^T \mathbf{Y} + 2\mathbf{X}^T \mathbf{X}\mathbf{B}$

The OLS estimate has the form

$$\hat{\mathbf{B}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} \Rightarrow \hat{\beta}_k = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}_k, \quad k \in \{1, \dots, d\}$$




Notes

Expected Value of Least Squares Coefficients

The expected value of the estimated coefficients is given by

$$\begin{aligned} \mathbb{E}(\hat{\mathbf{B}}) &= \mathbb{E}[(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}] \\ &= (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbb{E}(\mathbf{Y}) \\ &= (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{X}\mathbf{B} \\ &= \mathbf{B} \end{aligned}$$

$\Rightarrow \hat{\mathbf{B}}$ is an **unbiased estimator** of \mathbf{B}



Notes

Fitted Values and Residuals

- Fitted values are given by

$$\hat{\mathbf{Y}} = \mathbf{X}\hat{\mathbf{B}},$$


i.e.,

$$\hat{y}_{ik} = \hat{\beta}_{0k} + \sum_{j=1}^p \hat{\beta}_{jk} x_{ij}, \quad i = 1, \dots, n, \quad k = 1, \dots, d$$

- Residuals are given by

$$\hat{\mathbf{E}} = \mathbf{Y} - \hat{\mathbf{Y}},$$

i.e., $\hat{e}_{ik} = y_{ik} - \hat{y}_{ik}, \quad i = 1, \dots, n, \quad k = 1, \dots, d$



Notes

Hat Matrix

Just like in univariate linear regression we can write the fitted values as

$$\begin{aligned}\hat{Y} &= X\hat{B} \\ &= X(X^T X)^{-1} X^T Y \\ &= HY,\end{aligned}$$

where $H = X(X^T X)^{-1} X^T$ is the **hat matrix**

$\Rightarrow H$ projects y_k onto the column space of X for $k \in \{1, \dots, d\}$

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation
Model and Assumptions
Parameter Estimation
Inference and Prediction
Motor Trend Car Road Tests Data Analysis

5.13

Notes

Partitioning the Total Variation

We can partition the total covariation in $\{y_i\}_{i=1}^n$ (SSCP_{Tot}) as

$$\begin{aligned}\text{SSCP}_{\text{tot}} &= \sum_{i=1}^n (y_i - \bar{y})^T (y_i - \bar{y}) \\ &= \sum_{i=1}^n (y_i - \hat{y}_i + \hat{y}_i - \bar{y}) (y_i - \hat{y}_i + \hat{y}_i - \bar{y})^T \\ &= \underbrace{\sum_{i=1}^n (\hat{y}_i - \bar{y})(\hat{y}_i - \bar{y})^T}_{\text{SSCP}_{\text{Reg}}} + \underbrace{\sum_{i=1}^n (y_i - \hat{y}_i)(y_i - \hat{y}_i)^T}_{\text{SSCP}_{\text{Err}}} \\ &\quad + 2 \underbrace{\sum_{i=1}^n (\hat{y}_i - \bar{y})(y_i - \hat{y}_i)}_{=0} \\ &= \text{SSCP}_{\text{Reg}} + \text{SSCP}_{\text{Err}}\end{aligned}$$

The corresponding **degrees of freedom** are $d(n-1)$ for SSCP_{Tot}; dp for SSCP_{Reg}; and $d(n-p-1)$ for SSCP_{Err}

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation
Model and Assumptions
Parameter Estimation
Inference and Prediction
Motor Trend Car Road Tests Data Analysis

5.14

Notes

Estimated Error Covariance

The estimated error covariance matrix is

$$\begin{aligned}\hat{\Sigma} &= \frac{\sum_{i=1}^n (y_i - \hat{y}_i)(y_i - \hat{y}_i)^T}{n-p-1} \\ &= \frac{\text{SSCP}_{\text{Err}}}{n-p-1}\end{aligned}$$

- $\hat{\Sigma}$ is an **unbiased estimate** of Σ
- The estimate $\hat{\Sigma}$ is the **mean SSCP_{Err}**

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation
Model and Assumptions
Parameter Estimation
Inference and Prediction
Motor Trend Car Road Tests Data Analysis

5.15

Notes

Sampling Distributions of \hat{B} , \hat{Y} , and \hat{E}

We would need to figure out the **sampling distributions** of estimator and predictor in order to drawn inference

Given the model assumptions, we have

$$\begin{aligned}\text{vec}(\hat{B}) &\sim N(\text{vec}(B), \Sigma \otimes (X^T X)^{-1}) \\ \text{vec}(\hat{Y}) &\sim N(\text{vec}(XB), \Sigma \otimes H) \\ \text{vec}(\hat{E}) &\sim N(0, \Sigma \otimes (I - H)),\end{aligned}$$

where $\text{vec}(\cdot)$ is the vectorization operator and \otimes is the Kronecker product

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction
Motor Trend Car
Road Tests Data
Analysis

5.16

Notes

Inference about Multiple $\hat{\beta}_{jk}$

Assume that $q < p$ and want to test if a reduced model is sufficient:

$$H_0 : B_2 = 0_{p-q \times d}, \quad \text{versus} \quad H_a : B_2 \neq 0_{p-q \times d},$$

where

$$B = \begin{bmatrix} B_1 \\ B_2 \end{bmatrix}$$

is the partitioned of the coefficient vector
We can compare the SSCP_{Err} for the **full model**:

$$y_{ik} = \beta_{0k} + \sum_{j=1}^p \beta_{jk} x_{ij} + \varepsilon_{ik}, \quad k = 1, \dots, d$$

and the **reduced model**:

$$y_{ik} = \beta_{0k} + \sum_{j=1}^q \beta_{jk} x_{ij} + \varepsilon_{ik}, \quad k = 1, \dots, d$$

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction
Motor Trend Car
Road Tests Data
Analysis

5.17

Notes

Some Test Statistics

Let $\tilde{E} = n\tilde{\Sigma}$ denote the SSCP_{Err} matrix from the **full model**, and let $\tilde{H} = n(\tilde{\Sigma}_1 - \tilde{\Sigma})$ denote the hypothesis SSCP_{Err} matrix
Some test statistics for

$$H_0 : B_2 = 0_{p-q \times d}, \quad \text{versus} \quad H_a : B_2 \neq 0_{p-q \times d} :$$

- Wilks Lambda

$$\Lambda^* = \frac{|\tilde{E}|}{|\tilde{H} + \tilde{E}|}$$

Reject H_0 if Λ^* is “small”

- Hotelling-Lawley Trace

$$T_0^2 = \text{tr}(\tilde{H}\tilde{E}^{-1})$$

Reject H_0 if T_0^2 is “large”

- Pillai Trace

$$V = \text{tr}(\tilde{H}(\tilde{H} + \tilde{E})^{-1})$$

Reject H_0 if V is “large”

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction
Motor Trend Car
Road Tests Data
Analysis

5.18

Notes

Interval Estimation

We would like to estimate the **expected value of the response** for a given predictor $\mathbf{x}_h = (1, x_{h1}, \dots, x_{hp})$.


Note that we have

$$\hat{\mathbf{y}}_h \sim N(\mathbf{B}^T \mathbf{x}_h, \mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h \Sigma)$$

We can exploit the duality between interval estimation and hypothesis testing. That is, we can test

$$H_0 : \mathbb{E}(\mathbf{y}_h) = \mathbf{y}_h^* \text{ versus } H_a : \mathbb{E}(\mathbf{y}_h) \neq \mathbf{y}_h^*$$

The $100(1 - \alpha)\%$ confidence region is the collection of \mathbf{y}_h^* values that fail to reject H_0 at α level



Notes

Interval Estimation (Cont'd)

Test statistics:

$$T^2 = \left(\frac{\hat{\mathbf{B}}^T \mathbf{x}_h - \mathbf{B}^T \mathbf{x}_h}{\sqrt{\mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h}} \right)^T \hat{\Sigma}^{-1} \left(\frac{\hat{\mathbf{B}}^T \mathbf{x}_h - \mathbf{B}^T \mathbf{x}_h}{\sqrt{\mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h}} \right)$$

$$\underset{H_0}{\sim} \frac{d(n-p-1)}{n-p-d} F_{d, n-p-d}$$

Therefore, the $100(1 - \alpha)\%$ simultaneous **confidence interval** for y_{hk} is

$$\hat{y}_{hk} \pm \sqrt{\frac{d(n-p-1)}{n-p-d} F_{d, n-p-d}} \sqrt{\mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h \hat{\sigma}_{kk}},$$

$$k \in \{1, \dots, d\}$$



Notes

Predicting New Observations


Here we want to predict the **observed value of response** for a given predictor

- **Note:** interested in actual $\hat{\mathbf{y}}_h$ instead of $\mathbb{E}(\hat{\mathbf{y}}_h)$
- Given $\mathbf{x}_h = (1, x_{h1}, \dots, x_{hp})$, the fitted value is still $\hat{\mathbf{y}}_h = \hat{\mathbf{B}}^T \mathbf{x}_h$

We can exploit the duality between interval estimation and hypothesis testing. That is, we can test

$$H_0 : \mathbf{y}_h = \mathbf{y}_h^* \text{ versus } H_a : \mathbf{y}_h \neq \mathbf{y}_h^*$$

The $100(1 - \alpha)\%$ prediction interval is the collection of \mathbf{y}_h^* values that fail to reject H_0 at α level



Notes

Predicting New Observations (Cont'd)

Test statistics:

$$T^2 = \left(\frac{\hat{B}^T \mathbf{x}_h - B^T \mathbf{x}_h}{\sqrt{1 + \mathbf{x}_h^T (X^T X)^{-1} \mathbf{x}_h}} \right)^T \hat{\Sigma}^{-1} \left(\frac{\hat{B}^T \mathbf{x}_h - B^T \mathbf{x}_h}{\sqrt{1 + \mathbf{x}_h^T (X^T X)^{-1} \mathbf{x}_h}} \right)$$
$$\stackrel{H_0}{\sim} \frac{d(n-p-1)}{n-p-d} F_{d,n-p-d}$$

Therefore, the 100(1 - α)% simultaneous **prediction interval** for y_{hk} is

$$\hat{y}_{hk} \pm \sqrt{\frac{d(n-p-1)}{n-p-d} F_{d,n-p-d}} \sqrt{(1 + \mathbf{x}_h^T (X^T X)^{-1} \mathbf{x}_h)} \hat{\sigma}_{kk},$$
$$k \in \{1, \dots, d\}$$

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction


Motor Trend Car Road Tests Data Analysis

5.22

Notes

Motor Trend Car Road Tests Data Analysis

Study the linear relationship between mpg, disp, hp, wt (**responses**) and cyl, am, carb (**predictors**) in the mtcars dataset



Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.23

Notes

Multivariate Regression Fit

- Model: $\text{lm}(Y \sim \text{cyl} + \text{am} + \text{carb}, \text{data} = \text{mtcars})$
- Key findings:
 - mpg: decreases with more cylinders, increases with manual transmission
 - disp: strongly influenced by cyl
 - hp: influenced by cyl and carb
 - wt: influenced by cyl, am, carb
- **Note:** Multivariate regression produces the *same point estimates* as running separate regressions for each response

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation

Model and Assumptions

Parameter Estimation

Inference and Prediction

Motor Trend Car Road Tests Data Analysis

5.24

Notes

SSCP & Error Covariance

SSCP decomposition:

SSCP_{Tot} = SSCP_{Reg} + SSCP_{Err}

Estimated error covariance matrix ($\hat{\Sigma}$)

	mpg	disp	hp	wt
mpg	7.8680094	-53.27166	-19.7015979	-0.6575443
disp	-53.2716607	2504.87095	425.1328988	18.1065416
hp	-19.7015979	425.13290	577.2703337	0.4662491
wt	-0.6575443	18.10654	0.4662491	0.2573503

	mpg	disp	hp	wt
mpg	1.0000000	-0.3794645	-0.29233405	-0.46209388
disp	-0.3794645	1.0000000	0.35354314	0.71314929
hp	-0.2923340	0.3535431	1.0000000	0.03825304
wt	-0.4620939	0.7131493	0.03825304	1.0000000

⇒ Captures residual dependencies among responses

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation
Model and Assumptions
Parameter Estimation
Inference and Prediction
Motor Trend Car Road Tests Data Analysis

5.25

Notes

Do We Need “cyl”?

```
> mvlm0 <- lm(Y ~ am + carb, data = mtcars)
> anova(mvlm, mvlm0, test = "Wilks")
Analysis of Variance Table
```

Model 1: Y ~ cyl + am + carb
Model 2: Y ~ am + carb

	Res.Df	Df	Gen.var.	Wilks	approx F	num Df	Df
1	27		29.862				
2	29	2	43.692	0.16395	8.8181		8

den Df

Pr(>F)

1							
2	48		2.525e-07	***			

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation
Model and Assumptions
Parameter Estimation
Inference and Prediction
Motor Trend Car Road Tests Data Analysis

5.26

Notes

Confidence and Prediction Intervals

Default predict() lacks multivariate CI/PI, so we use an R function from Prof. Helwig at the University of Minnesota

```
> newdata <- data.frame(cyl = factor(6, levels = c(4, 6, 8)),
+   am = 1, carb = 4)
> # confidence interval
> pred.mlm(mvlm, newdata)
      mpg      disp      hp      wt
fit 21.51824 159.2707 136.98500 2.631108
lwr 16.65593  72.5141  95.33649 1.751736
upr 26.38055 246.0273 178.63351 3.510479
> # prediction interval
> pred.mlm(mvlm, newdata, interval = "prediction")
      mpg      disp      hp      wt
fit 21.518240 159.27070 136.98500 2.6311076
lwr  9.680053 -51.95435  35.58397 0.4901152
upr 33.356426 370.49576 238.38603 4.7720999
```

Multivariate Linear Regression

CLEMSON UNIVERSITY

Motivation
Model and Assumptions
Parameter Estimation
Inference and Prediction
Motor Trend Car Road Tests Data Analysis

5.27

Notes

Summary

In this lecture, we learned about Multivariate Linear Regression

- Model and Assumptions
- Parameter Estimation
- Inference and Prediction

Multivariate
Linear
Regression

CLEMSON
UNIVERSITY

Motivation
Model and
Assumptions
Parameter
Estimation
Inference and
Prediction

Motor Trend Car
Road Tests Data
Analysis

5.28

Notes

Notes

Notes
