

Lecture 15

Analysis of Variance (ANOVA)

Text: Chapters 8

STAT 8010 Statistical Methods I

March 5, 2020

Whitney Huang
Clemson University



Notes

Analysis of Variance (ANOVA)

- We use a z-test or a t-test to compare **means** of 2 groups
- To compare means of 3+ groups we use ANOVA to perform a F-test
- Overall F-test:

$$H_0 : \mu_1 = \mu_2 = \dots = \mu_J$$

H_a : At least one mean is different

Assumptions:

- The distribution of each group is normal with equal variance (i.e. $\sigma_1^2 = \sigma_2^2 = \dots = \sigma_J^2$)
- Responses are independent to each other



Notes

Partition of Sums of Squares

"Sums of squares" refers to sums of squared deviations from some mean. ANOVA decomposes the total sum of squares into treatment sum of squares and error sum of squares:

- Total sum of square: $SSTo = \sum_{j=1}^J \sum_{i=1}^{n_j} (X_{ij} - \bar{X})^2$
- Treatment sum of square: $SSTr = \sum_{j=1}^J n_j (\bar{X}_j - \bar{X})^2$
- Error sum of square: $SSE = \sum_{j=1}^J (n_j - 1) s_j^2$

We can show that $SSTo = SSTr + SSE$



Notes

ANOVA Table and F Test

Source	df	SS	MS	F statistic
Treatment	$J - 1$	SSTr	$MSTr = \frac{SSTr}{J - 1}$	$F = \frac{MSTr}{MSE}$
Error	$N - J$	SSE	$MSE = \frac{SSE}{N - J}$	
Total	$N - 1$	SSTo		

F-Test

- $H_0 : \mu_1 = \mu_2 = \dots = \mu_J$
 $H_a : \text{At least one mean is different}$
- Test Statistic: $F^* = \frac{MSTr}{MSE}$
- Under H_0 , $F^* \sim F_{df_1=J-1, df_2=N-J}$
 - Rejection Region Method: Reject H_0 if $F_{obs} > F_{\alpha, df_1=J-1, df_2=N-J}$
 - P-value Method: Reject H_0 if P-value $= \mathbb{P}(F^* > F_{obs}) < \alpha$

Analysis of
Variance (ANOVA)

CLEMSON
UNIVERSITY

15.4

Notes

Example

An experiment was conducted to determine if experience has an effect on the time it takes for mice to run a maze. Four treatment groups, consisting of mice having been trained on the maze one, two, three and four times were run through the maze and their times recorded. Three mice were originally assigned to each group, but it was discovered that some lab assistants, in an attempt to win a bet, gave one mouse a stimulant and another mouse a sedative. These mice were removed from the analysis.

Training runs	1	2	3	4
Times	11, 9	7,8,9	6,5,7	5,3

Fill out the ANOVA table and test whether the time to run the maze is affected by training. Use a significant level of .05.

Analysis of
Variance (ANOVA)

CLEMSON
UNIVERSITY

15.5

Notes

Multiple Comparisons

If we reject $H_0 : \mu_1 = \mu_2 = \dots = \mu_J$, we'll want to know which group means are different.

Analysis of
Variance (ANOVA)

CLEMSON
UNIVERSITY

15.6

Notes

One-Way ANOVA & Overall F-Test

- We use **one-way ANOVA** to compare means of **J** (≥ 3) **groups/conditions**

$$H_0 : \mu_1 = \mu_2 = \dots = \mu_J$$

$$H_a : \text{at least a pair } \mu\text{'s differ}$$

- If H_0 is rejected, ANOVA just states that there is a significant difference between the groups **but not where those differences occur**
- We need to perform additional post hoc tests, **multiple comparisons**, to determine where the group differences are

Notes

Pairwise T-Tests

- Suppose we have 4 groups, i.e. $J = 4$, then we need to perform $\binom{4}{2} = 6$ two-sample tests to locate where the group differences are

$$H_0 : \mu_1 = \mu_2 \text{ vs. } H_a : \mu_1 \neq \mu_2$$

$$H_0 : \mu_1 = \mu_3 \text{ vs. } H_a : \mu_1 \neq \mu_3$$

$$H_0 : \mu_1 = \mu_4 \text{ vs. } H_a : \mu_1 \neq \mu_4$$

$$H_0 : \mu_2 = \mu_3 \text{ vs. } H_a : \mu_2 \neq \mu_3$$

$$H_0 : \mu_2 = \mu_4 \text{ vs. } H_a : \mu_2 \neq \mu_4$$

$$H_0 : \mu_3 = \mu_4 \text{ vs. } H_a : \mu_3 \neq \mu_4$$

- What if we simply perform these tests using, say, $\alpha = 0.05$ for each test?

$$\mathbb{P}(\text{making a least one type I error}) = 1 - (1 - 0.05)^6 = 0.265$$

if each test was independent

Notes

Family-Wise Error Rate (FWER)

Family-Wise Error Rate (FWER) $\bar{\alpha}$: the probability of making 1 or more type I errors in a set of hypothesis tests

For m independent tests, each with individual type I error rate α , then we have

$$\bar{\alpha} = 1 - (1 - \alpha)^m$$

m	α		
	0.1	0.05	0.01
1	0.100	0.050	0.010
3	0.271	0.143	0.030
6	0.469	0.265	0.059
10	0.651	0.401	0.096
15	0.794	0.537	0.140
21	0.891	0.659	0.190

Notes

The Bonferroni Correction

If we would like to control the FWER to be α , then we adjust the significant level for each of the m tests to be $\frac{\alpha}{m}$

$$FWER = \mathbb{P}(\cup_{i=1}^m p_i \leq \frac{\alpha}{m}) \leq \sum_{i=1}^m \mathbb{P}(p_i \leq \frac{\alpha}{m}) = m \frac{\alpha}{m} = \alpha$$


where p_i is the p-value for the i_{th} test

If we have 4 treatment groups, then we need to perform 6 tests ($m = 6$) \Rightarrow will need to set the significant level for each individual pairwise t-test to be $0.05/6 = 0.0083$ to ensure that FWER is less than 0.05

Remark: Bonferroni procedure can be very conservative but gives guaranteed control over FWER at the risk of reducing statistical power. Does not assume independence of the comparisons.

Notes

Me and the significant boys



Me and the significant boys after Bonferroni correction



Notes

Example


A researcher who studies sleep is interested in the effects of ethanol on sleep time. She gets a sample of 20 rats and gives each an injection having a particular concentration of ethanol per body weight. There are 4 treatment groups, with 5 rats per treatment. She records Rapid eye movement (REM) sleep time for each rat over a 24-period.

Treatment	Control	1g/kg	2g/kg	4g/kg
Mean	82.2	81.0	73.8	65.7
Std	9.6	5.3	9.4	7.9

Recall in last lecture we reject $H_0 : \mu_1 = \mu_2 = \mu_3 = \mu_4$ at 0.05 level. But where these differences are?

Notes

Example: Multiple Testing with Bonferroni Correction



Notes

Fisher's Protected Least Significant Difference (LSD) Procedure

- We conclude that μ_i and μ_j differ at α significance level if $|\bar{X}_i - \bar{X}_j| > LSD$, where
$$LSD = t_{\alpha/2, df=N-J} \sqrt{MSE \left(\frac{1}{n_i} + \frac{1}{n_j} \right)}$$
- This procedure builds on the equal variances t-test of the difference between two means
- The test statistic is improved by using MSE rather than s_p^2

Notes


Tukey's Honest Significance Difference (HSD) Test

- The test procedure:
 - Requires equal sample size n per populations
 - Find a critical value ω as follows:
$$\omega = q_{\alpha}(J, N - J) \sqrt{\frac{MSE}{n}}$$
where $q_{\alpha}(J, N - J)$ can be obtained from the **studentized range table**
- If $\bar{X}_{max} - \bar{X}_{min} > \omega \Rightarrow$ there is sufficient evidence to conclude that $\mu_{max} > \mu_{min}$
- Repeat this procedure for each pair of samples. Rank the means if possible

Notes

Facebook Friends Example

A researcher would like to investigate the relationship between Facebook social attractiveness and the number of Facebook friends. An experiment was conducted where five groups of participant judge the same Facebook profiles, except for the one aspect that was manipulated: the number of friends for that profile.



Analysis of
Variance (ANOVA)

CLEMSON
UNIVERSITY

15.16

Notes

Example: Descriptive Statistics

	Score				
	102	302	502	702	902
Valid	24	33	26	30	21
Missing	0	0	0	0	0
Mean	3.817	4.879	4.562	4.407	3.990
Std. Deviation	0.999	0.851	1.070	1.428	1.023
Minimum	2.200	2.600	2.000	1.000	1.600
Maximum	6.000	6.400	6.800	7.000	5.200


Analysis of
Variance (ANOVA)

CLEMSON
UNIVERSITY

15.17

Notes

Example: Checking Model Assumptions



Analysis of
Variance (ANOVA)

CLEMSON
UNIVERSITY

15.18

Notes

Example: ANOVA Table

ANOVA - Score

Cases	Homogeneity Correction	Sum of Squares	df	Mean Square	F	p
Friends	None	19.890	4.000	4.973	4.142	0.003
Friends	Brown-Forsythe	19.890	4.000	4.973	4.184	0.003
Friends	Welch	19.890	4.000	4.973	5.445	< .001
Residual	None	154.867	129.000	1.201		
Residual	Brown-Forsythe	154.867	114.185	1.356		
Residual	Welch	154.867	61.144	2.533		

Note. Type III Sum of Squares

Analysis of Variance (ANOVA)

CLEMSON UNIVERSITY

15.19

Notes

Example: Multiple Testing

Post Hoc Tests

Post Hoc Comparisons - Friends

		95% CI for Mean Difference			t	Cohen's d	P-value	Post
		Mean Difference	Lower	Upper				
102	302	-1.062	-1.875	-0.249	0.294	-3.613	-1.160	0.004
	502	-0.745	-1.603	0.113	0.310	-2.402	-0.718	0.121
702		-0.590	-1.420	0.240	0.300	-1.966	-0.470	0.288
902		-0.174	-1.080	0.732	0.327	-0.531	-0.172	0.884
302	502	0.317	-0.478	1.112	0.287	1.104	0.333	0.804
	702	0.472	-0.293	1.237	0.276	1.708	0.406	0.433
	902	0.888	0.042	1.735	0.306	2.904	0.964	0.035
502	702	0.155	-0.657	0.967	0.294	0.528	0.121	0.984
	902	0.571	-0.318	1.460	0.321	1.776	0.544	0.392
702	902	0.416	-0.446	1.279	0.312	1.335	0.326	0.670

Note. Cohen's d does not correct for multiple comparisons.
Note. Confidence interval adjustment: tukey method for comparing a family of 5 estimates

Analysis of Variance (ANOVA)

CLEMSON UNIVERSITY

15.20

Notes

Notes