

Lecture 5

Descriptive Statistics III

Text: Chapter III

STAT 8010 Statistical Methods I
August 30, 2019

Whitney Huang
Clemson University



Notes

Agenda

- 1 Review of Last Class
- 2 Numerical Summaries of Quantitative Variables



Notes


Last Lecture

- Graphical Summaries of **Quantitative Variables**
 - Dot plot
 - Stem-and-Leaf Plot
 - Histogram
 - Boxplot
- Numerical Summaries of **Quantitative Variables**
 - Mean \Rightarrow A measure of **central tendency**



Notes

Shapes of Distributions



Notes

Measures of Center

- A **measure of center** attempts to report a "typical" value for the variable
- When a measure of center is calculated with **sample data** it is a **statistic**
- When a measure of center is calculated with popular (e.g., census data) it is a **parameter**
- **Measures:** Mean, Median, Mode

Notes

Mean

- The **population mean**, denoted by μ_X , is the sum of all the population values ($\{X_1, \dots, X_N\}$) divided by the total number (N) of population values. That is,

$$\mu_X = \frac{\sum_{i=1}^N X_i}{N}$$

- The **sample mean**, denoted by \bar{X} is the sum of all the sample values ($\{X_1, \dots, X_n\}$) divided by the total number of sample values (n). That is

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$$

Notes

Median

The **median** is the value separating the higher half from the lower half of a data sample

How to compute the median: Order the n observations in a data set from smallest to largest, then

Median =
$$\begin{cases} \text{the single middle value,} & n \text{ odd} \\ \text{the average of the middle two values,} & n \text{ even} \end{cases}$$

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.7

Notes

Mode

The **mode** is the value of the observation that appears most frequently

How to compute the mode(s): Order the observations in a data set from smallest to largest, then find the number that is repeated more often than any other

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables


5.8

Notes

Example

Suppose we have the following list of values: 13, 18, 13, 14, 13, 16, 14, 21, 13

- Plot this "data set" and describe the shape of the distribution



Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.9

Notes

Example cont'd

Suppose we have the following list of values: 13, 18, 13, 14, 13, 16, 14, 21, 13

- Find the sample mean

$$\bar{X} = \sum_{i=1}^9 \frac{13 + 18 + 13 + 14 + 13 + 16 + 14 + 21 + 13}{9} = 15$$

- Find the sample median
 - Order the data first: 13, 13, 13, 13, 14, 14, 16, 18, 21
 - Compute the sample size n and identify (or compute) the median value
 - $n = 9 \Rightarrow$ the median is the 5th number, which is 14

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.10

Notes

Example cont'd

- Find the mode
 - Order the data first: 13, 13, 13, 13, 14, 14, 16, 18, 21
 - We have 3 13 and 2 14 \Rightarrow 13 is the mode

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.11

Notes

Example: Resistant (Robust) Statistics

Suppose we have the following list of values: 13, 18, 13, 14, 13, 16, 14, 210, 13

- Find the sample mean
$$\bar{X} = \sum_{i=1}^9 \frac{13 + 18 + 13 + 14 + 13 + 16 + 14 + 210 + 13}{9} = 36$$
- Find the sample median
 - Order the data first: 13, 13, 13, 13, 14, 14, 16, 18, 210
 - Compute the sample size n and identify (or compute) the median value
 - $n = 9 \Rightarrow$ the median is the 5th number, which is (still) 14

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.12

Notes

Example cont'd

- Find the mode
 - Order the data first: 13, 13, 13, 13, 14, 14, 16, 18, 210
 - We have 3 13 and 2 14 \Rightarrow 13 is (still) the mode

Descriptive Statistics III

CLEMSON UNIVERSITY


Review of Last Class

Numerical Summaries of Quantitative Variables

5.13

Notes

Measures of Spread



- Measures:** Range, Variance(Standard Deviation), Interquartile range (IQR)

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.14

Notes

Range

The **range** of a dataset is the difference between the largest and smallest values

Range = Largest Value – Smallest Value

- Compute the range of the following list of values: 13, 18, 13, 14, 13, 16, 14, 21, 13
- Compute the range of the following list of values: 13, 18, 13, 14, 13, 16, 14, **210**, 13

Question: Is Range a robust statistic?

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.15

Notes

Standard Deviation (Variance)

- The sample standard deviation (variance), denoted by s (s^2), can be used to estimate the population standard deviation (variance), denoted by σ (σ^2).
- s is calculated in the following way:
 - ➊ Calculate the sample mean \bar{x}
 - ➋ Calculate the deviation (from the sample mean) for each observation (i.e., $x_i - \bar{x}$, $i = 1, \dots, n$)
 - ➌ Square each deviation and add them (i.e., $\sum_{i=1}^n (x_i - \bar{x})^2$)
 - ➍ Divide by $n - 1$ and take the square root (i.e., $s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}}$)

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.16

Notes

Example

- Compute s of the following list of values: 13, 18, 13, 14, 13, 16, 14, 21, 13
- Compute s of the following list of values: 13, 18, 13, 14, 13, 16, 14, 210, 13

Question: Is standard deviation a robust statistic?

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.17

Notes

Interquartile range (IQR)

- $IQR = Q_3 - Q_1$, where Q_1 is the Lower Quartile (the median of the lower half of the data) and Q_3 is the Upper Quartile (the median of the upper half of the data)
- Compute the IQR of the following list of values: 13, 18, 13, 14, 13, 16, 14, 21, 13
- Compute the IQR of the following list of values: 13, 18, 13, 14, 13, 16, 14, 210, 13

Question: Is IQR a robust statistic?

Descriptive Statistics III

CLEMSON UNIVERSITY

Review of Last Class

Numerical Summaries of Quantitative Variables

5.18

Notes
