

Lecture 27

An Overview of Spatial Interpolation

STAT 8020 Statistical Methods II
December 3, 2020

Whitney Huang
Clemson University

An Overview of
Spatial
Interpolation

CLEMSON
UNIVERSITY

Gaussian Process
Spatial Model


Spatial
Interpolation

Parameter
estimation

27.1

Notes

Toy Examples of Spatial Interpolation



Question: What is your best guess of the value of the missing pixel, denoted as $Y(s_0)$, for each case?

An Overview of
Spatial
Interpolation

CLEMSON
UNIVERSITY

Gaussian Process
Spatial Model


Spatial
Interpolation

Parameter
estimation

27.2

Notes

Interpolating Paraná State Precipitation Data



Goal: To interpolate the values in the spatial domain

An Overview of
Spatial
Interpolation

CLEMSON
UNIVERSITY

Gaussian Process
Spatial Model

Spatial
Interpolation

Parameter
estimation

27.3

Notes

The Spatial Interpolation Problem

Given observations of a spatially varying quantity Y at n spatial locations

$y(s_1), y(s_2), \dots, y(s_n), \quad s_i \in \mathcal{S}, i = 1, \dots, n$

We want to estimate this quantity at any **unobserved location**

$Y(s_0), \quad s_0 \in \mathcal{S}$

Applications

- Mining: ore grade
- Climate: temperature, precipitation, ...
- Remote Sensing: CO₂ retrievals
- Environmental Science: air pollution levels, ...

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.4

Notes

Some History

- Mining (Krige 1951) Matheron (1960s), Forestry (Matérn 1960)
- More recent work: Cressie (1993) Stein (1999)



An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.5

Notes

Outline

- 1 Gaussian Process Spatial Model
- 2 Spatial Interpolation
- 3 Parameter estimation

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.6

Notes

Linear Interpolation


The best guess (in a statistical sense) should be based on the conditional distribution $[Y(s_0) | Y = y]$ where

$$y = (y(s_1), \dots, y(s_n))^T$$

- Calculating this conditional distribution can be difficult
- Instead we use a **linear predictor**:

$$\hat{Y}(s_0) = \lambda_0 + \sum_{i=1}^n \lambda_i y(s_i)$$

- The best linear predictor is completely determined by the **mean** and **covariance** of $\{Y(s), s \in S\}$, and the observations y



Notes

Gaussian Process (GP) Spatial Model

We assume that the observed data $\{y(s_i)\}_{i=1}^n$ is one partial realization of a (continuously indexed) spatial GP $\{Y(s)\}_{s \in S}$.

Model:

$$Y(s) = m(s) + \epsilon(s), \quad s \in S \subset \mathbb{R}^d$$


where

- **Mean function:**

$$m(s) = E[Y(s)] = X^T(s)\beta$$

- **Covariance function:**

$$\{\epsilon(s)\}_{s \in S} \sim \text{GP}(0, K(\cdot, \cdot)), \quad K(s_1, s_2) = \text{Cov}(\epsilon(s_1), \epsilon(s_2))$$



Notes

Assumptions on Covariance Function

In practice, the covariance must be estimated from the data $(y(s_1), \dots, y(s_n))^T$. We need to impose some structural assumptions

- **Stationarity:**

$$\begin{aligned} K(s_1, s_2) &= \text{Cov}(\epsilon(s_1), \epsilon(s_2)) = C(s_1 - s_2) \\ &= \text{Cov}(\epsilon(s_1 + h), \epsilon(s_2 + h)) \end{aligned}$$

- **Isotropy:**

$$K(s_1, s_2) = \text{Cov}(\epsilon(s_1), \epsilon(s_2)) = C(\|s_1 - s_2\|)$$



Notes

A Valid Covariance Function Must Be Positive Definite (p.d.)!

A covariance function is positive if

$$\sum_{i,j=1}^n a_i a_j C(\mathbf{s}_i - \mathbf{s}_j) \geq 0$$

for any finite locations $\mathbf{s}_1, \dots, \mathbf{s}_n$, and for any constants $a_i, i = 1, \dots, n$

Question: what is the consequence if a covariance function is NOT p.d.? \Rightarrow weird things can happen

Question: How to guarantee a $C(\cdot)$ is p.d.?

- Using a **parametric covariance function**
- Using **Bochner's Theorem** to construct a valid covariance function

Notes

Some Commonly Used Covariance Functions

- **Powered exponential:**

$$C(h) = \sigma^2 \exp\left(-\left(\frac{h}{\rho}\right)^\alpha\right), \quad \sigma^2 > 0, \rho > 0, 0 < \alpha \leq 2$$

- **Spherical:**

$$C(h) = \sigma^2 \left(1 - 1.5 \frac{h}{\rho} + 0.5 \left(\frac{h}{\rho}\right)^3\right) \mathbb{1}_{\{h \leq \rho\}}, \quad \sigma^2, \rho > 0$$

Note: it is only valid for 1, 2, and 3 dimensional spatial domain.

- **Matérn:**

$$C(h) = \sigma^2 \frac{(\sqrt{2\nu}h/\rho)^\nu K_\nu(\sqrt{2\nu}h/\rho)}{\Gamma(\nu)2^{\nu-1}}, \quad \sigma^2 > 0, \rho > 0, \nu > 0$$

"Use the Matérn model" – Stein (1999, pp. 14)

Notes

1-D Realizations from Matérn Model with Fixed σ^2, ρ





Figure: courtesy of Rasmussen & Williams 2006

Notes

2-D Realizations from Matérn Model with Fixed σ^2



An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.13

Notes

Outline

- 1 Gaussian Process Spatial Model
- 2 Spatial Interpolation
- 3 Parameter estimation

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.14

Notes

Conditional Distribution of Multivariate Normal

If
$$\begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix} \sim N \left(\begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix} \right)$$

Then
$$[Y_1|Y_2 = y_2] \sim N(\mu_{1|2}, \Sigma_{1|2})$$

where
$$\begin{aligned} \mu_{1|2} &= \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(y_2 - \mu_2) \\ \Sigma_{1|2} &= \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} \end{aligned}$$

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.15

Notes

GP-Based Spatial Interpolation: Kriging

If $\{Y(s)\}_{s \in \mathcal{S}}$ follows a GP, then

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim N \left(\begin{pmatrix} m_0 \\ \mathbf{m} \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & k^T \\ k & \Sigma \end{pmatrix} \right)$$

We have

$$[Y_0 | \mathbf{Y} = \mathbf{y}] \sim N \left(m_{Y_0 | \mathbf{Y} = \mathbf{y}}, \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 \right)$$

where

$$\begin{aligned} m_{Y_0 | \mathbf{Y} = \mathbf{y}} &= m_0 + k^T \Sigma^{-1} (\mathbf{y} - \mathbf{m}) \\ \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 &= \sigma_0^2 - k^T \Sigma^{-1} k \end{aligned}$$

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model


Spatial Interpolation

Parameter estimation

27.16

Notes

Spatial Prediction of Paraná State Rainfall



An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.17

Notes

GP-Based Spatial Interpolation: Kriging

If $\{Y(s)\}_{s \in \mathcal{S}}$ follows a GP, then

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim N \left(\begin{pmatrix} m_0 \\ \mathbf{m} \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & k^T \\ k & \Sigma \end{pmatrix} \right)$$

We have

$$[Y_0 | \mathbf{Y} = \mathbf{y}] \sim N \left(m_{Y_0 | \mathbf{Y} = \mathbf{y}}, \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 \right)$$

where

$$\begin{aligned} m_{Y_0 | \mathbf{Y} = \mathbf{y}} &= m_0 + k^T \Sigma^{-1} (\mathbf{y} - \mathbf{m}) \\ \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 &= \sigma_0^2 - k^T \Sigma^{-1} k \end{aligned}$$

Question: what if we don't know $m_0, \mathbf{m}, \sigma_0^2, \Sigma$?

⇒ We need to estimate the mean and covariance from the data \mathbf{y} .

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.18

Notes

Outline

- 1 Gaussian Process Spatial Model
- 2 Spatial Interpolation
- 3 Parameter estimation

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation


27.19

Notes

Gaussian Process

We assume that the observed data $\{y(s_i)\}_{i=1}^n$ is one partial realization of a (continuously indexed) spatial stochastic process $\{Y(s)\}_{s \in \mathcal{S}}$.

- Gaussian Processes GP $(m(\cdot), K(\cdot, \cdot))$ are widely used in modeling spatial stochastic processes
- Spatial statisticians often focus on the covariance function. e.g. $K(h) = \sigma^2 \frac{(\sqrt{2\nu}h/\gamma)^\nu \mathcal{K}_\nu(\sqrt{2\nu}h/\gamma)}{\Gamma(\nu)2^{\nu-1}}$



An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model


Spatial Interpolation

Parameter estimation

27.20

Notes

Semivariogram $\{\frac{1}{2} \text{Var}(Y(s_i, s_j))\}_{i,j}$



Source: fields vignette by Wiens and Krock, 2019

An Overview of Spatial Interpolation

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

27.21

Notes

Variogram, Semivariogram, and Covariance Function


Under the stationary and isotropic assumptions

Variogram:

$$\begin{aligned} 2\gamma(\mathbf{s}_i, \mathbf{s}_j) &= \text{Var}(Y(\mathbf{s}_i) - Y(\mathbf{s}_j)) \\ &= \text{E} \left\{ ((Y(\mathbf{s}_i) - \mu(\mathbf{s}_i)) - (Y(\mathbf{s}_j) - \mu(\mathbf{s}_j)))^2 \right\} \\ &= \text{E} \left\{ (Y(\mathbf{s}_i) - Y(\mathbf{s}_j))^2 \right\} \\ &= 2\gamma(\|\mathbf{s}_i - \mathbf{s}_j\|) = 2\gamma(h) \end{aligned}$$

Semivariogram and covariance function:



$$\gamma(h) = C(0) - C(h)$$



Notes

Estimation: Least Squares Method

$$\underset{\boldsymbol{\theta}}{\text{argmin}} \sum_u \frac{n_u}{\gamma(h_u; \boldsymbol{\theta})^2} [\hat{\gamma}(h_u) - \gamma(h_u; \boldsymbol{\theta})]^2$$



Notes

Maximum Likelihood Estimation (MLE)

Log-likelihood:

Given data $\mathbf{y} = (y(\mathbf{s}_1), \dots, y(\mathbf{s}_n))^T$

$$\ell_n(\boldsymbol{\beta}, \boldsymbol{\theta}; \mathbf{y}) \propto -\frac{1}{2} \log |\boldsymbol{\Sigma}_{\boldsymbol{\theta}}| - \frac{1}{2} (\mathbf{y} - \mathbf{X}^T \boldsymbol{\beta})^T [\boldsymbol{\Sigma}_{\boldsymbol{\theta}}]_{n \times n}^{-1} (\mathbf{y} - \mathbf{X}^T \boldsymbol{\beta})$$

where $\boldsymbol{\Sigma}_{\boldsymbol{\theta}}(i, j) = \sigma^2 \rho_{\rho, \nu}(\|\mathbf{s}_i - \mathbf{s}_j\|) + \tau^2 \mathbb{1}_{\{\mathbf{s}_i = \mathbf{s}_j\}}$, $i, j = 1, \dots, n$

for any fixed $\boldsymbol{\theta}_0 \in \Theta$ the unique value of $\boldsymbol{\beta}$ that maximizes ℓ_n is given by

$$\hat{\boldsymbol{\beta}} = \left(\mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}_0}^{-1} \mathbf{X} \right)^{-1} \mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}_0} \mathbf{y}$$


Then we obtain the profile log likelihood

$$\ell_n(\boldsymbol{\theta}; \mathbf{y}) \propto -\frac{1}{2} \log |\boldsymbol{\Sigma}_{\boldsymbol{\theta}}| - \frac{1}{2} \mathbf{y}^T P(\boldsymbol{\theta}) \mathbf{y}$$

where

$$P(\boldsymbol{\theta}) = \boldsymbol{\Sigma}_{\boldsymbol{\theta}}^{-1} - \boldsymbol{\Sigma}_{\boldsymbol{\theta}}^{-1} \mathbf{X} \left(\mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}}^{-1} \mathbf{X} \right)^{-1} \mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}}$$

Solve the maximization problem above to get the MLE



Notes
