

Lecture 8


Introduction to Probability II

Text: Chapter 4

STAT 8010 Statistical Methods I

September 6, 2019


Whitney Huang
Clemson University



Notes

Agenda

- 1 Union and Intersection
- 2 Logical Relationships among Events
- 3 Complement Rule and General Addition Rule
- 4 Venn Diagram



Notes


Intersection and Union

- **Intersection:** the intersection of two events A and B , denoted by $A \cap B$, is the event that contains all outcomes of A that also belong to $B \Rightarrow$ **AND**

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 4, 5\}$, then $A \cap B = \{1, 2\}$

- **Union:** the union of two events A and B , denoted by $A \cup B$, is the event of all outcomes that belong to either A or $B \Rightarrow$ **OR**

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 4, 5\}$, then $A \cup B = \{1, 2, 3, 4, 5\}$



Notes

Example

Suppose we flipped 3 fair coins. Let A be the event of **exactly 2 tails**. Let B be the event that the **first 2 tosses are tails**. Let C be the event that **all 3 tosses are tails**. What are $A \cap B$, $A \cup C$, and $(A \cap B) \cup C$?

Solution.

$A = \{(T, T, H), (T, H, T), (H, T, T)\}$
 $B = \{(T, T, T), (T, T, H)\}$
 $C = \{T, T, T\}$

- $A \cap B = \{(T, T, H)\}$
- $A \cup C = \{(T, T, H), (T, H, T), (H, T, T), (T, T, T)\}$
- $(A \cap B) \cup C = \{(T, T, H)\} \cup \{(T, T, T)\} = \{(T, T, H), (T, T, T)\}$

Notes

Logical Relationships among Events

- Mutually exclusive:** refers to two (or more) events that cannot both occur when the random experiment is formed.

$A \cap B = \emptyset$

- Exhaustive:** refers to event(s) that comprise the sample space.


$A \cup B = \Omega$

- Partition:** events that are both mutually exclusive and exhaustive.

$A \cap B = \emptyset \quad \text{and} \quad A \cup B = \Omega$

Notes

Complement



Notes

Complement Rule

- 1 By the definition of complement

$A \cup A^c = \Omega$

- 2 Apply the probability operator

$P(A \cup A^c) = P(\Omega) = 1$

- 3 Since A and A^c are mutually exclusive

$P(A \cup A^c) = P(A) + P(A^c)$

- 4 Hence we get $P(A) = 1 - P(A^c)$

Introduction to Probability II

CLEMSON UNIVERSITY

Union and Intersection

Logical Relationships among Events

Complement Rule and General Addition Rule

Venn Diagram

8.7

Notes

Example

Suppose we rolled a fair, six-sided die 10 times. Let T be the event that we roll at least 1 three. If one were to calculate T you would need to find the probability of 1 three, 2 threes, \dots , and 10 threes and add them all up. However, you can use the complement rule to calculate $P(T)$

Solution.

Let X be the times that we rolled a 3, then
 $P(T) = P(X \geq 1) =$
 $P(X = 1) + P(X = 2) + \dots + P(X = 10)$

need to compute 10 probabilities
If we apply the complement rule
 $P(T) = 1 - P(T^c) = 1 - P(X = 0)$

Introduction to Probability II

CLEMSON UNIVERSITY

Union and Intersection

Logical Relationships among Events

Complement Rule and General Addition Rule


Venn Diagram

8.8

Notes

Venn Diagram

A Venn diagram is a diagram that shows all possible logical relations between a finite collection of events.



Introduction to Probability II

CLEMSON UNIVERSITY

Union and Intersection

Logical Relationships among Events

Complement Rule and General Addition Rule


Venn Diagram

8.9

Notes

General Addition Rule

The general addition rule is a way of finding the probability of a union of 2 events. It is $P(A \cup B) = P(A) + P(B) - P(A \cap B)$



Introduction to Probability II

CLEMSON UNIVERSITY

Union and Intersection

Logical Relationships among Events

Complement Rule and General Addition Rule

Venn Diagram

8.10

Notes

Example

Three of the major commercial computer operating systems are Windows, Mac OS, and Red Hat Linux Enterprise. A Computer Science professor selects 50 of her students and asks which of these three operating systems they use. The results for the 50 students are summarized below.

- 30 students use Windows
- 16 students use at least two of the operating systems
- 9 students use all three operating systems
- 18 students use Mac OS
- 46 students use at least one of the operating systems
- 11 students use both Windows and Linux
- 11 students use both Windows and Mac OS

Introduction to Probability II

CLEMSON UNIVERSITY

Union and Intersection

Logical Relationships among Events


Complement Rule and General Addition Rule

Venn Diagram

8.11

Notes

Example cont'd



Introduction to Probability II

CLEMSON UNIVERSITY

Union and Intersection

Logical Relationships among Events

Complement Rule and General Addition Rule

Venn Diagram


8.12

Notes

Conditional Probability

Let A and B be events. The probability that event A occurs **given** (knowing) that event B occurs is called a **conditional probability** and is denoted by $P(A|B)$. The formula of conditional probability is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$



In a conditional probability problem, the sample space is "reduced" to the "space" of the given outcome (e.g. if given B , we now just care about the probability of A occurring "inside" of B)

Notes

Summary

In this lecture, we learned

- Union , Intersection
- Logical Relationships among Events Mutually Exclusive, Exhaustive, Partition
- Complement Rule and General Addition Rule
- Venn Diagram

Notes

Notes
