

Lecture 5

Law of Total Probability, Bayes' Rule, and Random Variables

Text: Chapter 4

STAT 8010 Statistical Methods I

January 23, 2020

Whitney Huang
Clemson University

Agenda

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Law of Total Probability

Bayes' Rule

Random Variables

1 Law of Total Probability

2 Bayes' Rule

3 Random Variables

Law of Partitions & Multiplication Rule

Law of partitions

Let A_1, A_2, \dots, A_k form a partition of Ω . Then, for all events B ,

$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(A_i \cap B)$$

Multiplication rule

- 2 events:

$$\mathbb{P}(B \cap A) = \mathbb{P}(A) \times \mathbb{P}(B|A) = \mathbb{P}(B) \times \mathbb{P}(A|B)$$

- More than 2 events:

$$\begin{aligned} \mathbb{P}(\cap_{i=1}^n A_i) &= \mathbb{P}(A_1) \times \mathbb{P}(A_2|A_1) \times \mathbb{P}(A_3|A_1 \cap A_2) \\ &\quad \times \dots \times \mathbb{P}(A_n|A_{n-1} \cap \dots \cap A_1) \end{aligned}$$

Let A_1, A_2, \dots, A_k form a partition of Ω . Then, for all events B ,

$$\begin{aligned}\mathbb{P}(B) &= \sum_{i=1}^k \mathbb{P}(A_i \cap B) \\ &\quad \underbrace{\hspace{10em}}_{\text{Law of partitions}} \\ &= \sum_{i=1}^k \underbrace{\mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}_{\text{Multiplication rule}}\end{aligned}$$

Example

Suppose that two factories supply light bulbs to the market. Factory X's bulbs work for over 5000 hours in 99% of cases, whereas factory Y's bulbs work for over 5000 hours in 95% of cases. It is known that factory X supplies 60% of the total bulbs available and Y supplies 40% of the total bulbs available. What is the chance that a purchased bulb will work for longer than 5000 hours?

The Monty Hall Problem

There was an old television show called Let's Make a Deal, whose original host was named Monty Hall. The set-up is as follows. You are on a game show and you are given the choice of three doors. Behind one door is a car, behind the others are goats. You pick a door, and the host, who knows what is behind the doors, opens another door (not your pick) which has a goat behind it. Then he asks you if you want to change your original pick. The question we ask you is, "Is it to your advantage to switch your choice?"

The Monty Hall Problem

Law of Total
Probability, Bayes'
Rule, and Random
Variables

CLEMSON
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables



The Monty Hall Problem Solution

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Law of Total Probability

Bayes' Rule

Random Variables

General form

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

Law of Total Probability

Bayes' Rule

Random Variables

Let A_1, A_2, \dots, A_k form a partition of the sample space. Then for every event B in the sample space,

$$\mathbb{P}(A_j|B) = \frac{\mathbb{P}(B|A_j) \times \mathbb{P}(A_j)}{\sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}, j = 1, 2, \dots, k$$

Example

Let us assume that a specific disease is only present in 5 out of every 1,000 people. Suppose that the test for the disease is accurate 99% of the time a person has the disease and 95% of the time that a person lacks the disease. What is the probability that the person has the disease given that they tested positive?

Solution.

Example

Let us assume that a specific disease is only present in 5 out of every 1,000 people. Suppose that the test for the disease is accurate 99% of the time a person has the disease and 95% of the time that a person lacks the disease. What is the probability that the person has the disease given that they tested positive?

Solution.

$$\mathbb{P}(D|+) = \frac{\mathbb{P}(D \cap +)}{\mathbb{P}(+)} = \frac{.005 \times .99}{.005 \times .99 + .995 \times .05} = \frac{.00495}{.0547} = .0905$$

Example

Let us assume that a specific disease is only present in 5 out of every 1,000 people. Suppose that the test for the disease is accurate 99% of the time a person has the disease and 95% of the time that a person lacks the disease. What is the probability that the person has the disease given that they tested positive?

Solution.

$$\mathbb{P}(D|+) = \frac{\mathbb{P}(D \cap +)}{\mathbb{P}(+)} = \frac{.005 \times .99}{.005 \times .99 + .995 \times .05} = \frac{.00495}{.0547} = .0905$$

The reason we get such a surprising result is because the disease is so rare that the number of false positives greatly outnumber the people who truly have the disease.

Review of Probability (we learned so far)

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Basic Concepts:

Law of Total Probability

Bayes' Rule

Random Variables

Review of Probability (we learned so far)

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event

Law of Total Probability

Bayes' Rule

Random Variables

Review of Probability (we learned so far)

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event

Law of Total Probability

Bayes' Rule

Random Variables

Review of Probability (we learned so far)

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework

Law of Total Probability

Bayes' Rule

Random Variables

Review of Probability (we learned so far)

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework

Law of Total Probability

Bayes' Rule

Random Variables

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection, Mutually Exclusive, Exhaustive, Partition

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection, Mutually Exclusive, Exhaustive, Partition

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection, Mutually Exclusive, Exhaustive, Partition
- Venn Diagram

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection, Mutually Exclusive, Exhaustive, Partition
- Venn Diagram

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection, Mutually Exclusive, Exhaustive, Partition
- Venn Diagram
- Independence and Conditional Probability

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$
- Conditional probability: $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$
- Conditional probability: $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$
- Conditional probability: $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- Law of total probability:
$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(B \cap A_i) = \sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)$$

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$
- Conditional probability: $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- Law of total probability:
$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(B \cap A_i) = \sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)$$

Review: Probability Rules


- $0 \leq \mathbb{P}(A) \leq 1$ for any event A , $\mathbb{P}(\emptyset) = 0$, $\mathbb{P}(\Omega) = 1$
- Complement rule: $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule: $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:
$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$$
- Conditional probability: $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- Law of total probability:
$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(B \cap A_i) = \sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)$$
- Independence: if A and B are independent, then
$$\mathbb{P}(A|B) = \mathbb{P}(A), \mathbb{P}(B|A) = \mathbb{P}(B), \text{ and } \mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$$

Random Variables

A **random variable** is a real-valued function whose domain is the sample space of a random experiment. In other words, a random variable is a function

$$X : \Omega \mapsto \mathbb{R}$$

where Ω is the sample space of the random experiment under consideration and \mathbb{R} represents the set of all real numbers.



Example

The following is a chart describing the number of siblings each student in a particular class has.

Siblings (X)	Frequency	Relative Frequency
0	8	.200
1	17	.425
2	11	.275
3	3	.075
4	1	.025
Total	40	1

Example cont'd

Let's define the event A as the event that a randomly chosen student has 2 or more siblings. What is $\mathbb{P}(X \in A)$?

Solution.

$$\begin{aligned}\mathbb{P}(X \in A) &= \mathbb{P}(X \geq 2) = \mathbb{P}(X = 2) + \mathbb{P}(X = 3) + \mathbb{P}(X = 4) \\ &= .275 + 0.75 + 0.25 = .375\end{aligned}$$

Types of Random Variables

There are two main types of quantitative random variables: **discrete** and **continuous**. A discrete random variable often involves a count of something. Examples may include number of cars per household, etc.

Discrete random variable

A random variable X is called a discrete random variable if the outcome of the random variable is limited to a countable set of real numbers (usually integers).

Let X be a discrete random variable. Then the probability mass function (pmf) of X is the real-valued function defined on \mathbb{R} by

$$p_X(x) = \mathbb{P}(X = x)$$

The capital letter, X , is used to denote random variable.
Lowercase letter, x , is used to denote possible values of the random variable.

Example

Flip a fair coin 3 times. Let X denote the number of heads tossed in the 3 flips. Create a pmf for X

Solution.

The random variable X maps any outcome to an integer (e.g. $X((T, T, T)) = 0$)

Example

Flip a fair coin 3 times. Let X denote the number of heads tossed in the 3 flips. Create a pmf for X

Solution.

The random variable X maps any outcome to an integer (e.g. $X((T, T, T)) = 0$)

X	0	1	2	3
$p_X(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Properties of a PMF

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Law of Total Probability

Bayes' Rule

Random Variables

- $0 \leq p_X(x) \leq 1, \forall x \in R$

Properties of a PMF

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Law of Total Probability

Bayes' Rule

Random Variables

- $0 \leq p_X(x) \leq 1, \forall x \in R$

Properties of a PMF

- $0 \leq p_X(x) \leq 1, \forall x \in R$
- $\{x \in \mathbb{R} : p_X(x) \neq 0\}$ is countable

Properties of a PMF

- $0 \leq p_X(x) \leq 1, \forall x \in R$
- $\{x \in \mathbb{R} : p_X(x) \neq 0\}$ is countable

Properties of a PMF

- $0 \leq p_X(x) \leq 1, \forall x \in \mathcal{R}$
- $\{x \in \mathbb{R} : p_X(x) \neq 0\}$ is countable
- $\sum_x p_X(x) = 1$

Example

Let X be a random variable with pmf defined as follows:

$$p_X(x) = \begin{cases} k(5-x) & \text{if } x = 0, 1, 2, 3, 4 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find the value of k that makes $p_X(x)$ a legitimate pmf.
- 2 What is the probability that X is between 1 and 3 inclusive?
- 3 If X is not 0, what is the probability that X is less than 3?

Mean of Discrete Random Variables

The mean of a discrete r.v. X , denoted by $\mathbb{E}[X]$, is defined by

$$\mathbb{E}[X] = \sum_x x \times p_X(x)$$

Remark:

The mean of a discrete r.v. is a weighted average of its possible values, and the weight used is its probability. Sometimes we refer to the expected value as the **expectation (expected value)**, or the **first moment**.

For any function, say $g(X)$, we can also find an expectation of that function. It is

$$\mathbb{E}[g(X)] = \sum_x g(x) \times p_X(x)$$

Example

$$\mathbb{E}[X^2] = \sum_x x^2 \times p_X(x)$$

Let X and Y be discrete r.v.s defined on the same sample space and having finite expectation (i.e. $\mathbb{E}[X], \mathbb{E}[Y] < \infty$). Let a and b be constants. Then the following hold:

Let X and Y be discrete r.v.s defined on the same sample space and having finite expectation (i.e. $\mathbb{E}[X], \mathbb{E}[Y] < \infty$). Let a and b be constants. Then the following hold:

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$

Let X and Y be discrete r.v.s defined on the same sample space and having finite expectation (i.e. $\mathbb{E}[X], \mathbb{E}[Y] < \infty$). Let a and b be constants. Then the following hold:

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$

Let X and Y be discrete r.v.s defined on the same sample space and having finite expectation (i.e. $\mathbb{E}[X], \mathbb{E}[Y] < \infty$). Let a and b be constants. Then the following hold:

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[aX + b] = a \times \mathbb{E}[X] + b$

Number of Siblings Example Revisited

Siblings (X)	Frequency	Relative Frequency
0	8	.200
1	17	.425
2	11	.275
3	3	.075
4	1	.025
Total	40	1

Find the expected value of the number of siblings

Solution.

Number of Siblings Example Revisited

Siblings (X)	Frequency	Relative Frequency
0	8	.200
1	17	.425
2	11	.275
3	3	.075
4	1	.025
Total	40	1

Find the expected value of the number of siblings

Solution.

$$\mathbb{E}[X] = \sum_x xp_X(x) = 0 \times .200 + 1 \times .425 + 2 \times .275 + 3 \times .075 + 4 \times .025 = 1.3$$

Variance/Standard Deviation of Discrete r.v.'s

The **variance** of a (discrete) r.v., denoted by $Var(X)$, is a measure of the spread, or variability, in the r.v. $Var(X)$ is defined by

$$Var(X) = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

or

$$Var(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$$

The **standard deviation**, denoted by $Sd(X)$, is the square root of its variance

Properties of Variance

Law of Total
Probability, Bayes'
Rule, and Random
Variables



Law of Total Probability

Bayes' Rule

Random Variables

Let c be a constant. Then the following hold:

Properties of Variance

Let c be a constant. Then the following hold:

- $Var(cX) = c^2 \times Var(X)$

Properties of Variance

Let c be a constant. Then the following hold:

- $Var(cX) = c^2 \times Var(X)$

Properties of Variance

Let c be a constant. Then the following hold:

- $Var(cX) = c^2 \times Var(X)$
- $Var(X + c) = Var(X)$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

4 $\mathbb{E}[X^2 - 4]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

4 $\mathbb{E}[X^2 - 4]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

4 $\mathbb{E}[X^2 - 4]$

5 $\mathbb{E}[(X - 4)^2]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

4 $\mathbb{E}[X^2 - 4]$

5 $\mathbb{E}[(X - 4)^2]$

Example

Suppose X and Y are random variables with $\mathbb{E}[X] = 3$, $\mathbb{E}[Y] = 4$ and $\text{Var}(X) = 4$. Find:

1 $\mathbb{E}[2X + 1]$

2 $\mathbb{E}[X - Y]$

3 $\mathbb{E}[X^2]$

4 $\mathbb{E}[X^2 - 4]$

5 $\mathbb{E}[(X - 4)^2]$

6 $\text{Var}(2X - 4)$