

Lecture 7

Introduction to Probability

Text: Chapter IV

STAT 8010 Statistical Methods I
September 3, 2019

Whitney Huang
Clemson University



Notes


Agenda

- 1 Probability and Statistics
- 2 Basic Concepts of Probability




Notes

Probability and Statistics



Notes

Probability and Statistics



Introduction to Probability

CLEMSON UNIVERSITY

Probability and Statistics

Basic Concepts of Probability

74

Notes

Definitions

The framework of Probability is based on the paradigm of a **random experiment**, i.e., an action whose outcome cannot be predicted beforehand.

- **Outcome:** A particular result of an (random) experiment. (e.g. rolling a 3 on a die roll)
- **Event:** A collection of one or more outcomes of an experiment. (e.g. rolling an odd number on a die roll)
- **Sample space:** the set of all possible outcomes for an experiment. We will use Ω to denote it
- **Probability:** A number between 0 and 1 that reflects the likelihood of occurrence of some events.

Introduction to Probability

CLEMSON UNIVERSITY

Probability and Statistics

Basic Concepts of Probability

75

Notes

Example

We are interested in whether the price of the *S&P 500* **decreases**, **stays the same**, or **increases**. If we were to examine the *S&P 500* over one day, then $\Omega = \{\text{decrease, stays the same, increases}\}$. What would Ω be if we looked at 2 days?

Solution.

Introduction to Probability

CLEMSON UNIVERSITY

Probability and Statistics

Basic Concepts of Probability

76

Notes

Example

Let us examine what happens in the flip of 3 fair coins. In this case $\Omega = \{(T, T, T), (T, T, H), (T, H, T), (H, T, T), (T, H, H), (H, T, H), (H, H, T), (H, H, H)\}$. Let A be the event of exactly 2 tails. Let B be the event that the first 2 tosses are tails. Let C be the event that all 3 tosses are tails. Write out the possible outcomes for each of these 3 events

Solution.

Introduction to Probability

CLEMSON UNIVERSITY

Probability and Statistics

Basic Concepts of Probability

7.7

Notes

Example

Start with a standard deck of 52 cards and remove all the hearts and all the spades, leaving 13 red and 13 black cards. List the cards in each of the following sets:

- N = not a face card
-
- R = neither red nor an ace
-
- E = either black, even, or a Jack
- {Ace, 2, 3, 4, 5, 6, 7, 8, 9, 10, Jack, Queen, King, 2, 4, 6, 8, 10, Jack}

Introduction to Probability

CLEMSON UNIVERSITY

Probability and Statistics

Basic Concepts of Probability

7.8

Notes

Example

Suppose a fair six-sided die is rolled twice. Determine the number of possible outcomes

- For this experiment
- The sum of the two rolls is 5
- The two rolls are the same
- The sum of the two rolls is an even number

Solution.

Introduction to Probability

CLEMSON UNIVERSITY

Probability and Statistics

Basic Concepts of Probability

7.9

Notes

Frequentist Interpretation of Probability

The probability of an event is the **long-run proportion** of times that the event occurs in independent repetitions of the random experiment. This is referred to as an **empirical probability** and can be written as

$$P(\text{event}) = \frac{\text{number of times that event occurs}}{\text{number of random experiment}}$$



Notes

Equally Likely Framework

$$P(\text{event}) = \frac{\text{number of outcomes for the event}}{\text{number of all possible outcomes}}$$

Remark:

- Any individual outcome of the sample space is equally likely as any other outcome in the sample space.
- In an equally likely framework, the probability of any event is the number of ways the event occurs divided by the number of total events possible.



Notes

Example

Find the probabilities associated with parts 2–4 of the previous example

Solution.

- The probability that the sum of the two rolls is 5:
 $\frac{4}{36} = \frac{1}{9}$
- The probability that the two rolls are the same:
 $\frac{6}{36} = \frac{1}{6}$
- The probability that the sum of the two rolls is an even number:
 $\frac{18}{36} = \frac{1}{2}$




Notes

Probability Rules

- ➊ Any probability must be between 0 and 1 inclusively
- ➋ The sum of the probabilities for all the experimental outcomes must equal 1

If a probability model satisfies the two rules above, it is said to be legitimate




Notes

Example

An experiment with three outcomes has been repeated 50 times, and it was learned that outcome 1 occurred 20 times, outcome 2 occurred 13 times, and outcome 3 occurred 17 times. Assign probabilities to the outcomes. What method did you use?

Solution.



Notes


Example

A decision maker subjectively assigned the following probabilities to the four possible outcomes of an experiment:

$$P(E_1) = 0.1 \ P(E_2) = 0.15 \ P(E_3) = 0.4 \ P(E_4) = 0.2$$

Are these probability assignments legitimate? Explain.


Solution.



Notes

Independence: A Motivating Example

Example
You toss a fair coin and it comes up “Heads” three times. What is the chance that the next toss will also be a “Head”?



Notes


Independence and Conditional Probability

Conditional Probability
Let A and B be events. The probability that event B occurs **given** (knowing) that event A occurs is called a **conditional probability** and is denoted by $P(B|A)$. The formula of conditional probability is

$$P(B|A) = \frac{P(B \cap A)}{P(A)}$$

Independent events
Suppose $P(A) > 0$, $P(B) > 0$. We say that event B is **independent** of event A if the occurrence of event A does not affect the probability that event B occurs.


$P(B|A) = P(B) \Rightarrow P(B \cap A) = P(B)P(A)$



Notes

Summary

- In this lecture, we learned
- **Some definitions:** Outcome, Event, Sample Space
 - The Frequentist Interpretation of Probability and the Equally Likely Framework
 - Probability Rules
 - Independence and Conditional Probability



Notes
