

Lecture 3

Stationary processes

Readings: Cryer & Chan Ch. 2 & Ch. 4.1-4.3; Brockwell & Davis Ch. 1.3-1.6; Shumway & Stoffer Ch. 1.2-1.6

MATH 8090 Time Series Analysis
August 31 & September 2, 2021

Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions

Whitney Huang
Clemson University

1 Mean and Autocovariance Functions

2 Stationarity

3 Some Examples of Stationary Processes

4 Estimation of Mean and Autocovariance Functions

Review: The Additive Decomposition

- The additive model for a time series $\{Y_t\}$ is

$$Y_t = \mu_t + s_t + \eta_t,$$

where

- μ_t is the **trend** component
- s_t is the **seasonal** component
- η_t is the **random (noise)** component with $\mathbb{E}(\eta_t) = 0$
- Standard procedure:
 - (1) Estimate/remove the trend and seasonal components
 - (2) Analyze the remainder, the residuals $\hat{\eta}_t = y_t - \hat{\mu}_t - \hat{s}_t$
- We will focus on (2) for the next few weeks

- A **time series model** is a specification of the probabilistic distribution of a sequence of random variables (RVs) η_t


(The observed time series is a **realization** of such a sequence of random variables)

- The simplest time series is **i.i.d. (*independent and identically distributed*) noise**
 - $\{\eta_t\}$ is a sequence of independent and identically distributed zero-mean (i.e., $\mathbb{E}(\eta_t) = 0, \forall t$) random variables
 \Rightarrow **no temporal dependence**
 - It is of little value of using i.i.d. noise model to conduct **forecast** as there is no information from the past observations
 - **But**, we will use i.i.d. model as a building block to develop time series models that can accommodate time dependence

Example Realizations of i.i.d. Noise


- Gaussian (normal) i.i.d. noise with mean 0 and variance $\sigma^2 > 0$

$$f(\eta_t) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{\eta_t^2}{2\sigma^2}\right)$$



- Bernoulli i.i.d. noise with “success” probability

$$\mathbb{P}(\eta_t = 1) = p = 1 - \mathbb{P}(\eta_t = -1)$$



A time series model could also be a specification of the **means** and **autocovariances** of the RVs

- The **mean function** of $\{\eta_t\}$ is

$$\mu_t = \mathbb{E}(\eta_t).$$

- μ_t is the population mean at time t , which can be computed as:

$$\mu_t = \begin{cases} \int_{-\infty}^{\infty} \eta_t f(\eta_t) d\eta_t & \text{when } \eta_t \text{ is a continuous RV;} \\ \sum_{-\infty}^{\infty} \eta_t p(\eta_t), & \text{when } \eta_t \text{ is a discrete RV,} \end{cases}$$

where $f(\cdot)$ and $p(\cdot)$ are the probability density function and probability mass function of η_t , respectively

- **Example 1:** What is the mean function for $\{\eta_t\}$, an i.i.d. $N(0, \sigma^2)$ process?
- **Example 2:** For each time point, let $Y_t = \beta_0 + \beta_1 t + \eta_t$ with β_0 and β_1 some constants and η_t is defined above. What is $\mu_Y(t)$?

Review: The Covariance Between Two RVs

- The **covariance** between the RVs X and Y is

$$\begin{aligned}\mathbb{Cov}(X, Y) &= \mathbb{E}\{(X - \mu_X)(Y - \mu_Y)\} \\ &= \mathbb{E}(XY) - \mu_X \mu_Y.\end{aligned}$$

It is a measure of **linear dependence** between the two RVs. When $X = Y$ we have

$$\mathbb{Cov}(X, X) = \mathbb{Var}(X).$$

- For constants a, b, c , and RVs X, Y, Z :

$$\begin{aligned}\mathbb{Cov}(aX + bY + c, Z) &= \mathbb{Cov}(aX, Z) + \mathbb{Cov}(bY, Z) \\ &= a\mathbb{Cov}(X, Z) + b\mathbb{Cov}(Y, Z)\end{aligned}$$

\Rightarrow

$$\begin{aligned}\mathbb{Var}(X + Y) &= \mathbb{Cov}(X, X) + \mathbb{Cov}(X, Y) + \mathbb{Cov}(Y, X) + \mathbb{Cov}(Y, Y) \\ &= \mathbb{Var}(X) + \mathbb{Var}(Y) + 2\mathbb{Cov}(X, Y)\end{aligned}$$

- The autocovariance function of $\{\eta_t\}$ is

$$\gamma(s, t) = \text{Cov}(\eta_s, \eta_t) = \mathbb{E}[(\eta_s - \mu_s)(\eta_t - \mu_t)]$$

It measures the strength of linear dependence between two RVs η_s and η_t

- **Properties:**

- $\gamma(s, t) = \gamma(t, s)$ for each s and t
- When $s = t$ we have

$$\gamma(t, t) = \text{Cov}(\eta_t, \eta_t) = \text{Cov}(\eta_t) = \sigma_t^2$$

the value of the variance function at time t

- $\gamma(s, t)$ is a non-negative definite function (will come back to this later)

- The autocorrelation function of $\{\eta_t\}$ is

$$\rho(s, t) = \text{Corr}(\eta_s, \eta_t) = \frac{\gamma(s, t)}{\sqrt{\gamma(s, s)\gamma(t, t)}}$$

It measures the “scale invariant” linear association between η_s and η_t

- **Properties:**

- $-1 \leq \rho(s, t) \leq 1$ for each s and t
- $\rho(s, t) = \rho(t, s)$ for each s and t
- $\rho(t, t) = 1$ for each t
- $\rho(\cdot, \cdot)$ is a non-negative definite function

Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions

- We typically need “replicates” to estimate population quantities. For example, we use

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

to be the estimate of μ_X , the population mean of the **single** RV, X

- However, in time series analysis, we have $n = 1$ (i.e., no replication) because we only have one realized value at each time point
- **Stationarity** means that some characteristic of $\{\eta_t\}$ does not depend on the time point, t , only on the “time lag” between time points **so that we can create “replicates”**

Next, we will talk about **strict stationarity** and **weak stationarity**

- A time series, $\{\eta_t\}$, is **strictly stationary** if

$$[\eta_1, \eta_2, \dots, \eta_T] \stackrel{d}{=} [\eta_{1+h}, \eta_{2+h}, \dots, \eta_{T+h}],$$

for all integers h and $T \geq 1 \Rightarrow$ the **joint distribution** are unaffected by time shifts

- Under such the strict stationarity
 - $\{\eta_t\}$ is **identically distributed** but not (necessarily) **independent**
 - When μ_t is finite, $\mu_t = \mu$ is independent of time t
 - When the variance function exists,

$$\gamma(s, t) = \gamma(s + h, t + h),$$

for any s, t , and h

- $\{\eta_t\}$ is **weakly stationary** if
 - $\mathbb{E}(\eta_t) = \mu_t = \mu$
 - $\text{Cov}(\eta_t, \eta_{t+h}) = \gamma(t, t+h) = \gamma(h)$, finite constant that can depend on h **but not on t**
- Other names for this type of stationarity include **second-order, covariance, wide sense**. The quantity h is called the **lag**
- Weak and strict stationarity
 - A strictly stationary process $\{\eta_t\}$ is also weakly stationary as long as μ is finite
 - **Weak stationarity does not imply strict stationarity!**

The autocovariance function (ACVF) of a stationary process $\{\eta_t\}$ is defined to be

$$\begin{aligned}\gamma(h) &= \text{Cov}(\eta_t, \eta_{t+h}) \\ &= \mathbb{E}[(\eta_t - \mu)(\eta_{t+h} - \mu)],\end{aligned}$$

which measures the lag- h time dependence

Properties of the ACVF:

- $\gamma(0) = \text{Var}(\eta_t)$
- $\gamma(-h) = \gamma(h)$ for each h
- $\gamma(s - t)$ as a function of $(s - t)$ is non-negative definite

The autocorrelation function (ACF) of a stationary process $\{\eta_t\}$ is defined to be

$$\rho(h) = \frac{\gamma(h)}{\gamma(0)}$$

which measures the “scale invariant” lag- h time dependence

Properties of the ACF:

- $-1 \leq \rho(h) \leq 1$ and $\rho(0) = 1$ for each h
- $\rho(-h) = \rho(h)$ for each h
- $\rho(s-t)$ as a function of $(s-t)$ is non-negative definite

Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions

The White Noise Process

Let's assume $\mathbb{E}(\eta_t) = \mu$ and $\text{Var}(\eta_t) = \sigma^2 < \infty$. $\{\eta_t\}$ is a **white noise** or **WN**(μ, σ^2) process if

$$\gamma(h) = 0,$$

for $h \neq 0$

- $\{\eta_t\}$ is stationary
- However, distributions of η_t and η_{t+1} **can be different!**
- All i.i.d. noise with finite variance ($\sigma^2 < \infty$) is **white noise** but **the converse need not be true**

Examples Realizations of White Noise Processes

Stationary processes





CLEMSON
UNIVERSITY

Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions



The Moving Average Process of First Order (MA(1))

Let $\{Z_t\}$ be a $WN(0, \sigma^2)$ process and θ be some constant $\in \mathbb{R}$.
For each integer t , let

$$\eta_t = Z_t + \theta Z_{t-1}.$$

- The sequences of RVs $\{\eta_t\}$ is called the **moving average process of order 1** or MA(1) process
- One can show that the MA(1) process $\{\eta_t\}$ is **stationary**

First-Order Moving Average Process: Mean Function

Need to show the mean function is NOT a function of time t

$$\begin{aligned}\mathbb{E}[\eta_t] &= \mathbb{E}[Z_t + \theta Z_{t-1}] \\ &= \mathbb{E}[Z_t] + \theta \mathbb{E}[Z_{t-1}] \\ &= 0 + \theta \times 0 \\ &= 0, \quad \forall t\end{aligned}$$



First-Order Moving Average Process: Covariance Function

Need to show the autocovariance function $\gamma(\cdot, \cdot)$ is a function of time lag only

$$\begin{aligned}\gamma(t, t+h) &= \text{Cov}(\eta_t, \eta_{t+h}) \\ &= \text{Cov}(Z_t + \theta Z_{t-1}, Z_{t+h} + \theta Z_{t+h-1}) \\ &= \text{Cov}(Z_t, Z_{t+h}) + \text{Cov}(Z_t, \theta Z_{t+h-1}) \\ &\quad + \text{Cov}(\theta Z_{t-1}, Z_{t+h}) + \text{Cov}(\theta Z_{t-1}, \theta Z_{t+h-1})\end{aligned}$$

$$\begin{aligned}\text{if } h = 0, \text{ we have} \quad & \gamma(t, t+h) = \sigma^2 + \theta^2 \sigma^2 = \sigma^2(1 + \theta^2) \\ \text{if } h = \pm 1, \text{ we have} \quad & \gamma(t, t+h) = \theta \sigma^2 \\ \text{if } |h| \geq 2, \text{ we have} \quad & \gamma(t, t+h) = 0\end{aligned}$$

$\Rightarrow \gamma(t, t+h)$ only depends on h but not on t 😊

First-Order Moving Average Process: ACVF & ACF

ACVF:

$$\gamma(h) = \begin{cases} \sigma^2(1 + \theta^2) & h = 0; \\ \theta\sigma^2 & |h| = 1; \\ 0 & |h| \geq 2 \end{cases}$$

We can get **ACF** by dividing everything by $\gamma(0) = \sigma^2(1 + \theta^2)$

$$\rho(h) = \begin{cases} 1 & h = 0; \\ \frac{\theta}{1+\theta^2} & |h| = 1; \\ 0 & |h| \geq 2. \end{cases}$$

Examples Realizations of MA(1) Processes

Stationary processes



Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions

First-order autoregressive process, AR(1)

Let $\{Z_t\}$ be a $WN(0, \sigma^2)$ process, and $-1 < \phi < 1$ be a constant.
Let's assume $\{\eta_t\}$ is a **stationary process** with

$$\eta_t = \phi\eta_{t-1} + Z_t,$$

for each integer t , where η_s and Z_t are **uncorrelated** for each $s < t \Rightarrow$ future noise is uncorrelated with the current time point)

We will see later there is only one unique solution to this equation. Such a sequence $\{\eta_t\}$ of RVs is called an **AR(1) process**

Properties of the AR(1) process

Want to find the mean value μ under the weakly stationarity assumption

$$\begin{aligned}\mathbb{E}[\eta_t] &= \mathbb{E}[\phi\eta_{t-1} + Z_t] \\ \mu &= \phi\mathbb{E}[\eta_{t-1}] + \mathbb{E}[Z_t] \\ \mu &= \phi\mu + 0 \\ \Rightarrow \mu &= 0, \quad \forall t\end{aligned}$$



Want to find $\gamma(h)$ under the weakly stationarity assumption

$$\begin{aligned}\text{Cov}(\eta_t, \eta_{t-h}) &= \text{Cov}(\phi\eta_{t-1} + Z_t, \eta_{t-h}) \\ \gamma(-h) &= \phi\text{Cov}(\eta_{t-1}, \eta_{t-h}) + \text{Cov}(Z_t, \eta_{t-h}) \\ \gamma(h) &= \phi\gamma(h-1) + 0 \\ \Rightarrow \gamma(h) &= \phi\gamma(h-1) = \dots = \phi^{|h|}\gamma(0)\end{aligned}$$

Next, need to figure out $\gamma(0)$

Properties of the AR(1) process Cont'd

$$\text{Var}(\eta_t) = \text{Var}(\phi\eta_{t-1} + Z_t)$$

$$\gamma(0) = \phi^2\gamma(0) + \sigma^2$$

$$\Rightarrow (1 - \phi^2)\gamma(0) = \sigma^2$$

$$\Rightarrow \gamma(0) = \frac{\sigma^2}{1 - \phi^2}$$



Therefore, we have

$$\gamma(h) = \begin{cases} \frac{\sigma^2}{1-\phi^2} & h = 0; \\ \phi^{|h|} \frac{\sigma^2}{1-\phi^2} & h \neq 0, \end{cases}$$

and

$$\rho(h) = \begin{cases} 1 & h = 0; \\ \phi^{|h|} & h \neq 0. \end{cases}$$

Examples Realizations of AR(1) Processes

Stationary processes



Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions

Let $\{Z_t\}$ be a $WN(0, \sigma^2)$ process and for $t \geq 1$ define

$$\eta_t = Z_1 + Z_2 + \cdots + Z_t = \sum_{s=1}^t Z_s.$$

- The sequence of RVs $\{\eta_t\}$ is called a **random walk process**
- **Special case:** If we have $\{Z_t\}$ such that for each t

$$\mathbb{P}(Z_t = z) = \begin{cases} \frac{1}{2}, & z = 1; \\ \frac{1}{2}, & z = -1, \end{cases}$$

then $\{\eta_t\}$ is a **simple symmetric random walk**

- **The random walk process is not stationary!**

Example Realizations of Random Walk Processes

Stationary processes



Mean and
Autocovariance
Functions

Stationarity

Some Examples of
Stationary Processes

Estimation of Mean
and Autocovariance
Functions

$\{\eta_t\}$ is a **Gaussian process (GP)** if the joint distribution of any collection of the RVs has a multivariate normal (aka Gaussian) distribution

- The distribution of a GP is fully characterized by $\mu(\cdot)$, the mean function, and $\gamma(\cdot, \cdot)$, the autocovariance function. The joint probability density function of $\boldsymbol{\eta} = (\eta_1, \eta_2, \dots, \eta_T)^T$ is

$$f(\boldsymbol{\eta}) = \frac{1}{(2\pi)^{\frac{T}{2}} |\Sigma|^{\frac{1}{2}}} \exp\left(-\frac{1}{2}(\boldsymbol{\eta} - \boldsymbol{\mu})^T \Sigma^{-1}(\boldsymbol{\eta} - \boldsymbol{\mu})\right),$$

where $\boldsymbol{\mu} = (\mu_1, \mu_2, \dots, \mu_T)^T$ and the (i, j) element of the covariance matrix Σ is $\gamma(i, j)$

- If a GP $\{\eta_t\}$ is **weakly stationary** then the process is also **strictly stationary**

Estimating the Mean of Stationary Processes

Let $\{\eta_t\}$ be stationary with mean μ and ACVF $\gamma(s, t)$

- A natural estimator of μ is the sample mean

$$\bar{\eta} = \frac{1}{T} \sum_{t=1}^T \eta_t.$$

$\bar{\eta}$ is an unbiased estimator of μ , i.e.

- Since $\{\eta_t\}$ is stationary, we have

$$\begin{aligned} \text{Var}(\bar{\eta}) &= \frac{1}{T^2} \text{Var}\left(\sum_{i=1}^T \eta_t\right) \\ &= \frac{1}{T^2} \sum_{s=1}^T \sum_{t=1}^T \text{Cov}(\eta_s, \eta_t) \\ &= \frac{1}{T^2} \sum_{s=1}^T \sum_{t=1}^T \gamma(s-t) \end{aligned}$$

- **Exercise:** Show

$$\text{Var}(\bar{\eta}) = \frac{1}{T} \sum_{h=-(T-1)}^{T-1} \left(1 - \frac{|h|}{T}\right) \gamma(h)$$

AR(1) Example

Suppose $\{\eta_1, \eta_2, \eta_3\}$ is an AR(1) process with $|\phi| < 1$ and innovation variance σ . Show that the variance of $\bar{\eta}$ is

$$\frac{\sigma^2}{9(1+\phi^2)}(3 + 4\phi + 2\phi^2)$$

Solution:

Let

$$v_T = \sum_{h=-(T-1)}^{(T-1)} \left(1 - \frac{|h|}{T}\right) \gamma(h)$$

- If $\{\eta_t\}$ is **Gaussian** we have

$$\sqrt{T}(\bar{\eta} - \mu) \sim N(0, v_T)$$

- The result above is **approximate** for many **non-Gaussian** time series
- In practice we also need to **estimate** $\gamma(h)$ from the data

- If $\gamma(h) \rightarrow 0$ as $h \rightarrow \infty$ then

$$v = \lim_{T \rightarrow \infty} v_T = \sum_{h=-\infty}^{\infty} \gamma(h) \text{ exists.}$$

- Further, if $\{\eta_t\}$ is **Gaussian** and

$$\sum_{h=-\infty}^{\infty} |\gamma(h)| < \infty,$$

then an **approximate large-sample** 95% CI for μ is given by

$$\left[\bar{\eta} - 1.96\sqrt{\frac{v}{T}}, \bar{\eta} + 1.96\sqrt{\frac{v}{T}} \right]$$

1. Parametric:

- Assume a parametric model $\gamma_{\theta}(\cdot)$, and calculate $v = \sum_{h=-(T-1)}^{T-1} \left(1 - \frac{|h|}{T}\right) \gamma_{\hat{\theta}}(h)$ based on the ACVF for that model
- The standard error, v , will depend on the parameters θ of the parametric model

Nonparametric:

- Estimate v by

$$\hat{v} = \sum \left(1 - \frac{|h|}{T}\right) \hat{\gamma}(h),$$

where $\hat{\gamma}(\cdot)$ is an nonparametric estimate of ACVF

Examples of Parametric Forms for v

- **i.i.d. Gaussian Noise:** $v = \gamma(0) = \sigma \Rightarrow$ CI reduces to the classical case:

$$\left[\bar{\eta} - 1.96\sqrt{\frac{\sigma}{T}}, \bar{\eta} + 1.96\sqrt{\frac{\sigma}{T}} \right]$$

- **MA(1) process:** We have

$$\begin{aligned} v &= \sum_{h=-\infty}^{\infty} \gamma(h) = \gamma(-1) + \gamma(0) + \gamma(1) \\ &= \gamma(0) + 2\gamma(1) \\ &= \sigma^2(1 + \theta^2 + 2\theta) = \sigma^2(1 + \theta)^2 \end{aligned}$$

- **Exercise:** Show for an **AR(1)** process we have

$$v = \frac{\sigma^2}{(1 - \phi)^2}$$

Goal: Want to estimate

$$\gamma(h) = \text{Cov}(\eta_t, \eta_{t+h}) = \mathbb{E}[(\eta_t - \mu)(\eta_{t+h} - \mu)]$$

using data $\{\eta_t\}_{t=1}^T$

- For $|h| < T$, consider $\hat{\gamma}(h) = \frac{1}{T} \sum_{t=1}^{T-|h|} (\eta_t - \bar{\eta})(\eta_{t+|h|} - \bar{\eta})$. We call $\hat{\gamma}(h)$ the **sample ACVF**
- The sample ACVF is a **biased** estimator of $\gamma(h)$, but, it is used as the **standard** estimate of $\gamma(h)$
- $\hat{\gamma}(h)$ are **even** and **non-negative definite**