

Lecture 4

Multivariate Normal Distribution, Copula, and Nonparametric Density Estimation

Readings: Zelterman, 2015 Chapters 5, 6, 7, Izeman, 2008 Chapter 4.1, 4.3, 4.5

DSA 8070 Multivariate Analysis

Whitney Huang
Clemson University



Notes

Agenda

1 Multivariate Normal Distribution

2 Geometry of the Multivariate Normal Density

3 Copula

4 Nonparametric Density Estimation



Notes

The Multivariate Normal Distribution

Just as the **univariate normal distribution** tends to be the most important distribution in **univariate statistics**, the **multivariate normal distribution** is the most important distribution in **multivariate statistics**

- **Mathematical Simplicity:** It is easy to obtain multivariate methods based on the multivariate normal distribution
- **Central Limit Theorem:** The *sample mean vector* is going to be approximately *multivariate normally distributed* when the sample size is sufficiently large
- Many natural phenomena may be modeled using this distribution (perhaps after transformation)




Notes

Review: Univariate Normal Distributions

The probability density function of the normal distribution is

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma^2} \exp\left\{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right\},$$

where μ and σ^2 are its mean and variance, respectively.




$\left(\frac{x-\mu}{\sigma}\right)^2 = (x-\mu)(\sigma^2)^{-1}(x-\mu)$ is the squared statistical distance between x and μ in standard deviation units

Notes

Multivariate Normal Distributions

If we have a p -dimensional random vector that is distributed according to a **multivariate normal distribution** with mean vector $\mu = (\mu_1, \mu_2, \dots, \mu_p)^T$ and covariance matrix $\Sigma = \{\sigma_{ij}\}$, the probability density function is

$$f(x) = \frac{1}{2\pi^{\frac{p}{2}}|\Sigma|^{\frac{1}{2}}} \exp\left\{-\frac{1}{2}(x-\mu)^T \Sigma^{-1}(x-\mu)\right\}.$$



Notes

Review: Central Limit Theorem (CLT)

The **sampling distribution** of the **mean** will become approximately **normally distributed** as the **sample size becomes larger, irrespective of the shape of the population distribution!**

Let X_1, X_2, \dots, X_n $i.i.d.$ F with $\mu = E[X_i]$ and $\sigma^2 = \text{Var}[X_i]$. Then $\bar{X}_n = \frac{\sum_{i=1}^n X_i}{n} \xrightarrow{d} N(\mu, \frac{\sigma^2}{n})$ as $n \rightarrow \infty$.

Notes

CLT In Action

- ① Generate 100 (n) random numbers from an Exponential distribution (population distribution)
- ② Compute the **sample mean** of these 100 random numbers
- ③ Repeat this process 120 times



Notes

Properties of the Multivariate Normal Distribution

- If $\mathbf{X} \sim N(\mu, \Sigma)$, then any subset of \mathbf{X} also has a multivariate normal distribution

Example: Each single variable

$$X_i \sim N(\mu_i, \sigma_i^2), \quad i = 1, \dots, p$$

- If $\mathbf{X} \sim N(\mu, \Sigma)$, then any linear combination of the variables has a univariate normal distribution

Example: If $\mathbf{Y} = \mathbf{a}^T \mathbf{X}$. Then $\mathbf{Y} \sim N(\mathbf{a}^T \boldsymbol{\mu}, \mathbf{a}^T \boldsymbol{\Sigma} \mathbf{a})$

- Any conditional distribution for a subset of the variables conditional on known values for another subset of variables is a multivariate distribution

Example: $X_1 | \mathbf{X}_2 = \mathbf{x}_2 \sim$

$$N(\boldsymbol{\mu}_1 + \Sigma_{12} \Sigma_{22}^{-1} (\mathbf{x}_2 - \boldsymbol{\mu}_2), \Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21})$$



Notes

Example: Linear Combination of the Cholesterol Measurements [source: Penn State Univ. STAT 505]

Cholesterol levels were taken 0, 2, and 4 days following the heart attack on n patients. The mean vector is:

Variable	Mean
X_1 (0-day)	259.5
X_2 (2-day)	230.8
X_3 (4-day)	221.5

and the covariance matrix

$$\mathbf{S} = \begin{bmatrix} 2276 & 1508 & 813 \\ 1508 & 2206 & 1349 \\ 813 & 1349 & 1865 \end{bmatrix}$$

Suppose we are interested in $\Delta = X_2 - X_1$, the difference between the 2-day and the 0-day measurements. We can write the linear combination of interest as

$$\Delta = \mathbf{a}^T \mathbf{X} = [-1 \ 1 \ 0] \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix}$$



Notes

Cholesterol Measurements Example Cont'd

- The mean value for the difference Δ is

$$[-1 \ 1 \ 0] \begin{bmatrix} 259.5 \\ 230.8 \\ 221.5 \end{bmatrix} = -28.7$$

- The variance for Δ is

$$\begin{aligned} & [-1 \ 1 \ 0] \begin{bmatrix} 2276 & 1508 & 813 \\ 1508 & 2206 & 1349 \\ 813 & 1349 & 1865 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \\ &= [-768 \ 698 \ 536] \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \\ &= 1466 \end{aligned}$$

- If we assume these three variables together follows a multivariate normal distribution, then Δ follows a univariate normal distribution





Notes

Bivariate Normal Distribution

Let's focus bivariate normal distributions first as we can visualize them to facilitate our understanding. Suppose we have X_1 and X_2 jointly follows a bivariate normal distribution:

$$\begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \sim N \left[\begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix} \right]$$

Let's fix $\mu_1 = \mu_2 = 0$ and $\sigma_1^2 = \sigma_2^2 = 1$



Notes

Exponent of Multivariate Normal Distribution

Recall the multivariate normal density:

$$f(x) = \frac{1}{2\pi^{\frac{p}{2}} |\Sigma|^{\frac{1}{2}}} \exp \left\{ -\frac{1}{2}(x - \mu)^T \Sigma^{-1} (x - \mu) \right\}.$$


This density function only depends on x through the squared Mahalanobis distance: $(x - \mu)^T \Sigma^{-1} (x - \mu)$

- For bivariate normal, we get an ellipse whose equation is $(x - \mu)^T \Sigma^{-1} (x - \mu) = c^2$ which gives all $x = (x_1, x_2)$ pairs with constant density

- These ellipses are call contours and all are centered around μ

- A constant probability contour equals


$$\begin{aligned} &= \text{all } x \text{ such that } (x - \mu)^T \Sigma^{-1} (x - \mu) = c^2 \\ &= \text{surface of ellipsoid centered at } \mu \end{aligned}$$



Notes

Multivariate Normality and Outliers

The variable $d^2 = (\mathbf{X} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{X} - \boldsymbol{\mu})$ has a chi-square distribution with p degrees of freedom, i.e., $d^2 \sim \chi^2_p$ if $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ \Rightarrow we can exploit this result to check multivariate normality and to detect outliers




- Sort $(\mathbf{x}_i - \bar{\mathbf{x}})^T S^{-1} (\mathbf{x}_i - \bar{\mathbf{x}})$ in an increasing order to get sample quantiles
- Calculate the theoretical quantiles using the chi-square quantiles with $p = \frac{i-0.5}{n}$, $i = 1, \dots, n$
- Plot sample quantile against theoretical quantiles



Notes

Eigenvalues and Eigenvectors of $\boldsymbol{\Sigma}$ and the Geometry of the Multivariate Normal Density

Let $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, where $\boldsymbol{\mu} = (10, 5)^T$ and $\boldsymbol{\Sigma} = \begin{bmatrix} 64 & 16 \\ 16 & 9 \end{bmatrix}$.
The 95% probability contour is shown below



Next, we talk about how to "draw" this contour



Notes

Probability Contours

- The solid ellipsoid of values x satisfy

$$(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \leq c^2 = \chi^2_{df=p,\alpha}$$


Here we have $p = 2$ and $\alpha = 0.05 \Rightarrow c = \sqrt{\chi^2_{2,0.05}} = 2.4478$

- Major axis: $\boldsymbol{\mu} \pm c\sqrt{\lambda_1} \mathbf{e}_1$, where $(\lambda_1, \mathbf{e}_1)$ is the first eigenvalue/eigenvector of $\boldsymbol{\Sigma}$.

$$\Rightarrow \lambda_1 = 68.316, \quad \mathbf{e}_1 = \begin{bmatrix} -0.9655 \\ -0.2604 \end{bmatrix}$$


- Minor axis: $\boldsymbol{\mu} \pm c\sqrt{\lambda_2} \mathbf{e}_2$, where $(\lambda_2, \mathbf{e}_2)$ is the second eigenvalue/eigenvector of $\boldsymbol{\Sigma}$.

$$\Rightarrow \lambda_2 = 4.684, \quad \mathbf{e}_2 = \begin{bmatrix} 0.2604 \\ -0.9655 \end{bmatrix}$$



Notes

Graph of 95% Probability Contour




Notes

Example: Wechsler Adult Intelligence Scale [source: Penn State Univ. STAT 505]

We have data (wechslet.txt) on 37 subjects ($n = 37$) taking the Wechsler Adult Intelligence Test, which consists four different components: 1) Information; 2) Similarities; 3) Arithmetic; 4) Picture Completion.

- ① Calculate the sample mean vector \bar{x} and covariance matrix S
- ② Compute the eigenvalues and eigenvectors of S and give a geometry interpretation
- ③ Diagnostic the multivariate normal assumption



Notes

Beyond Normality: Copula [Sklar, 1959; Joe, 1997]

A copula is a multivariate cumulative distribution function for which the marginal probability distribution of each variable is uniform on the interval $[0, 1]$

$$\begin{aligned} F(x_1, \dots, x_p) &= \Pr(X_1 \leq x_1, \dots, X_p \leq x_p) \\ &= \Pr(F_1^{-1}(U_1) \leq x_1, \dots, F_p^{-1}(U_p) \leq x_p) \\ &= \Pr(U_1 \leq F_1(x_1), \dots, U_p \leq F_p(x_p)) \\ &= C(F_1(x_1), \dots, F_p(x_p)) \end{aligned}$$

- Copulas are used to model the **dependence** between random variables
- Copula approach has becomes popular in many areas, e.g., quantitative finance as it allows for **separate modeling of marginal distributions and dependence structure**




Notes

An Illustration of Bivariate Gaussian Copula

Left: Normal marginals + Gaussian Copula ($\rho = 0.7$)

Right: Exponential marginals + Gaussian Copula ($\rho = 0.7$)



The copula approach allows us to “build” multivariate distributions with non-normal marginals

Notes


Notes

More Examples

Marginal: normal
Copula: Gaussian

Marginal: normal
and normal

Marginal: Beta
and Log-normal
Copula: Clayton
 $\theta = 0.95$




⇒ The copula approach allows for more options for dependence modeling

Notes

A Financial Application Using Copula


Here we illustrate how to use a copula to model the bivariate joint distribution of S&P 500 and Nasdaq (log) returns



- 1 Transform the data $(x_{1i}, x_{2i})_{i=1}^n$ to $(u_{1i}, u_{2i})_{i=1}^n$ and fit a copula model to it
 - 2 Fit a distribution to $\{x_{1i}\}_{i=1}^n$ and $\{x_{2i}\}_{i=1}^n$, respectively
 - 3 Combine the fitted copula and marginal distributions to form the fitted bivariate distribution

Notes

Marginals, Copula, and Joint Distribution




Notes

Notes

Old Faithful Geyser Data


Waiting time between eruptions and the duration of the eruption for the Old Faithful geyser in Yellowstone NP



Notes

Notes

Histograms of Old Faithful Data




Notes

Transition from Histogram to Kernel Density

Goal: to estimate the probability density function

$$f(x)$$




- Histogram:

$$\hat{f}(x) = \sum_{j=1}^m \frac{\#\text{ of } x_i \in B_j}{nh} \mathbb{1}(x \in B_j),$$

Multivariate
Normal
Distribution

where B_j is the j th bin and h is the binwidth





- Kernel Density:

$$\hat{f}(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - x_i}{h}\right),$$

where $K(\cdot)$ is the kernel function

Notes

Kernel Density Estimates of Old Faithful



Notes

Summary

In this lecture, we learned about:

- Multivariate Normal Distribution
 - Copula Modeling
 - Non-parametric Density Estimation

In the next lecture, we will learn about making inferences for a mean vector