

Lecture 6

Comparisons of Several Mean Vectors

DSA 8070 Multivariate Analysis
September 20 - September 24, 2021


Whitney Huang
Clemson University



Notes

Agenda

- 1 Comparisons of Two Mean Vectors
- 2 Multivariate Analysis of Variance



Notes

Motivating Example: Swiss Bank Notes (Source: PSU stat 505)

Suppose there are two distinct populations for 1000 franc Swiss Bank Notes:

- The first population is the population of Genuine Bank Notes
- The second population is the population of Counterfeit Bank Notes

For both populations the following measurements were taken:

- 1 Length of the note
- 2 Width of the Left-Hand side of the note
- 3 Width of the Right-Hand side of the note
- 4 Width of the Bottom Margin
- 5 Width of the Top Margin
- 6 Diagonal Length of Printed Area

We want to determine if counterfeit notes can be distinguished from the genuine Swiss bank notes



Notes

Review: Two Sample t-Test

Suppose we have data from a single variable from population 1: $X_{11}, X_{12}, \dots, X_{1n_1}$ and population 2: $X_{21}, X_{22}, \dots, X_{2n_2}$. Here we would like to draw inference about their population means μ_1 and μ_2 .

Assumptions:

- **Homoskedasticity**: The data from both populations have common variance σ^2
- **Independence**: The subjects from both populations are independently sampled. Also $\{X_{1i}\}_{i=1}^{n_1}$ and $\{X_{2j}\}_{j=1}^{n_2}$ are independent to each other
- **Normality**: The data from both populations are normally distributed (not that crucial for "large" sample)

Here we are going to consider testing $H_0 : \mu_1 = \mu_2$ against $H_a : \mu_1 \neq \mu_2$



Notes

Review: Two Sample t-Test

We define the sample means for each population using the following expression:

$$\bar{x}_1 = \frac{\sum_{j=1}^{n_1} x_{1j}}{n_1}, \quad \bar{x}_2 = \frac{\sum_{j=1}^{n_2} x_{2j}}{n_2}.$$

We denote the sample variance

$$s_1^2 = \frac{\sum_{j=1}^{n_1} (x_{1j} - \bar{x}_1)^2}{n_1 - 1}, \quad s_2^2 = \frac{\sum_{j=1}^{n_2} (x_{2j} - \bar{x}_2)^2}{n_2 - 1}.$$

Under the **homoskedasticity** assumption, we can "pool" two samples to get the pooled sample variance

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

Test statistic

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}} \stackrel{H_0}{\sim} t_{n_1 + n_2 - 2}$$

We can use this result to construct confidence intervals and to perform hypothesis tests



Notes

The Two Sample Problem: The Multivariate Case

Now we would like to use two independent samples $\{X_{11}, \dots, X_{12}, \dots, X_{1n_1}\}$ and $\{X_{21}, \dots, X_{22}, \dots, X_{2n_2}\}$, where

$$\mathbf{X}_{ij} = \begin{bmatrix} X_{ij1} \\ X_{ij2} \\ \vdots \\ X_{ijp} \end{bmatrix}$$

to infer the relationship between μ_1 and μ_2 , where

$$\mu_i = \begin{bmatrix} \mu_{i1} \\ \mu_{i2} \\ \vdots \\ \mu_{ip} \end{bmatrix}$$

Assumptions

- Both populations have common covariance matrix, i.e., $\Sigma_1 = \Sigma_2$
- **Independence**: The subjects from both populations are independently sampled
- **Normality**: Both populations are normally distributed



Notes

The Multivariate Two-Sample Problem

Here we are testing

$$H_0 : \begin{bmatrix} \mu_{11} \\ \mu_{12} \\ \vdots \\ \mu_{1p} \end{bmatrix} = \begin{bmatrix} \mu_{21} \\ \mu_{22} \\ \vdots \\ \mu_{2p} \end{bmatrix}, \quad H_a : \mu_{1k} \neq \mu_{2k} \text{ for at least one } k \in \{1, 2, \dots, p\}$$

Under the **common covariance** assumption we have

$$S_p = \frac{(n_1 - 1)S_1 + (n_2 - 1)S_2}{n_1 + n_2 - 2},$$

where

$$S_i = \frac{1}{n_i - 1} \sum_{j=1}^{n_i} (x_{ij} - \bar{x}_i)(x_{ij} - \bar{x}_i)^T, \quad i = 1, 2$$



Notes

The Two-Sample Hotelling's T-Square Test Statistic

The two-sample t test is equivalent to

$$t^2 = (\bar{x}_1 - \bar{x}_2)^T \left[s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \right]^{-1} (\bar{x}_1 - \bar{x}_2).$$

Under H_0 , $t^2 \sim F_{1, n_1 + n_2 - 2}$. We can use this result to perform a hypothesis test

We can extend this to the multivariate situation:

$$T^2 = (\bar{x}_1 - \bar{x}_2)^T \left[S_p \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \right]^{-1} (\bar{x}_1 - \bar{x}_2)$$

Under H_0 , we have

$$F = \frac{n_1 + n_2 - p - 1}{p(n_1 + n_2 - 2)} T^2 \sim F_{p, n_1 + n_2 - p - 1}$$

We can use this result to perform inferences for multivariate cases



Notes

Two-Sample Test for Swiss Bank Notes

```
> (xbar1 <- colMeans(dat[real, -1]))
  V2   V3   V4   V5   V6   V7
214.969 129.943 129.720  8.305 10.168 141.517
> (xbar2 <- colMeans(dat[fake, -1]))
  V2   V3   V4   V5   V6   V7
214.823 130.300 130.193 10.530 11.133 139.450
> Sigma1 <- cov(dat[real, -1])
> Sigma2 <- cov(dat[fake, -1])
> n1 <- length(real); n2 <- length(fake); p <- dim(dat[, -1])[2]
> Sp <- ((n1 - 1) * Sigma1 + (n2 - 1) * Sigma2) / (n1 + n2 - 2)
> # Test statistic
> T.squared <- as.numeric(t(xbar1 - xbar2) %*% solve(Sp * (1 / n1 + 1 / n2)) %*% (xbar1 - xbar2))
> Fobs <- T.squared * ((n1 + n2 - p - 1) / ((n1 + n2 - 2) * p))
> # p-value
> pf(Fobs, p, n1 + n2 - p - 1, lower.tail = F)
[1] 3.378887e-105
```

Conclusion

The counterfeit notes can be distinguished from the genuine notes on at least one of the measurements \Rightarrow which ones?



Notes

Simultaneous Confidence Intervals

$$\bar{x}_{1k} - \bar{x}_{2k} \pm \sqrt{\frac{p(n_1 + n_2 - 2)}{n_1 + n_2 - p - 1} F_{p, n_1 + n_2 - p - 1, \alpha}} \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2}\right) s_{k,p}^2}$$

where $s_{k,p}^2$ is the pooled variance for the variable k

Variable	95% CI
Length of the note	(−0.04, 0.34)
Width of the Left-Hand note	(−0.52, −0.20)
Width of the Right-Hand note	(−0.64, −0.30)
Width of the Bottom Margin	(−2.70, −1.75)
Width of the Top Margin	(−1.30, −0.63)
Diagonal Length of Printed Area	(1.81, 2.33)



Notes

Checking Model Assumptions

Assumptions:

- **Homoskedasticity:** The data from both populations have common covariance matrix Σ

Will return to this in next slide

- **Independence:**

This assumption may be violated if we have clustered, time-series, or spatial data

- **Normality:**

Multivariate QQplot, univariate histograms, bivariate scatter plots



Notes

Testing for Equality of Mean Vectors when $\Sigma_1 \neq \Sigma_2$

- **Bartlett's test** can be used to test if $\Sigma_1 = \Sigma_2$ but this test is **sensitive** to departures from normality
- As a crude rule of thumb: if $s_{1,k}^2 > 4s_{2,k}^2$ or $s_{2,k}^2 > 4s_{1,k}^2$ for some $k \in \{1, 2, \dots, p\}$, then it is likely that $\Sigma_1 \neq \Sigma_2$
- Life gets difficult if we cannot assume that $\Sigma_1 = \Sigma_2$. However, if both n_1 and n_2 are "large", we can use the following approximation to conduct inferences:

$$T^2 = (\bar{X}_1 - \bar{X}_2)^T \left[\frac{1}{n_1} S_1 + \frac{1}{n_2} S_2 \right]^{-1} (\bar{X}_1 - \bar{X}_2) \stackrel{H_0}{\sim} \chi_p^2$$



Notes

Comparing More Than Two Populations:
Romano-British Pottery Example (source: PSU stat 505)

- Pottery shards are collected from four sites in the British Isles:
 - Llanedynr (L)
 - Caldicot (C)
 - Isle Thorns (I)
 - Ashley Rails (A)
- The concentrations of five different chemicals were be used
 - Aluminum (Al)
 - Iron (Fe)
 - Magnesium (Mg)
 - Calcium (Ca)
 - Sodium (Na)
- **Objective:** to determine whether the chemical content of the pottery depends on the site where the pottery was obtained

Comparisons of Several Mean Vectors

CLEMSON UNIVERSITY

Comparisons of Two Mean Vectors

Multivariate Analysis of Variance

6.13

Notes

Review: Analysis of Variance (ANOVA)

- $H_0 : \mu_1 = \mu_2 = \dots = \mu_g$
 $H_a : \text{At least one mean is different}$

Source	df	SS	MS	F statistic
Treatment	$g - 1$	SSTr	$MSTr = \frac{SSTr}{g-1}$	$F = \frac{MSTr}{MSE}$
Error	$N - g$	SSE	$MSE = \frac{SSE}{N-g}$	
Total	$N - 1$	SSTo		
- Test Statistic: $F^* = \frac{MSTr}{MSE}$. Under H_0 ,
 $F^* \sim F_{df_1=g-1, df_2=N-g}$
- **Assumptions:**
 - The distribution of each group is normal with equal variance (i.e. $\sigma_1^2 = \sigma_2^2 = \dots = \sigma_g^2$)
 - Responses for a given group are independent to each other

Comparisons of Several Mean Vectors

CLEMSON UNIVERSITY

Comparisons of Two Mean Vectors

Multivariate Analysis of Variance

6.14

Notes

One-way Multivariate Analysis of Variance (One-way MANOVA)

Treatment

12...g

1

2

...

g

Subject

1

2

...

n_i

$$\mathbf{Y}_{11} = \begin{pmatrix} Y_{111} \\ Y_{112} \\ \vdots \\ Y_{11p} \end{pmatrix} \quad \mathbf{Y}_{21} = \begin{pmatrix} Y_{211} \\ Y_{212} \\ \vdots \\ Y_{21p} \end{pmatrix} \quad \dots \quad \mathbf{Y}_{g1} = \begin{pmatrix} Y_{g11} \\ Y_{g12} \\ \vdots \\ Y_{g1p} \end{pmatrix}$$
$$\mathbf{Y}_{12} = \begin{pmatrix} Y_{121} \\ Y_{122} \\ \vdots \\ Y_{12p} \end{pmatrix} \quad \mathbf{Y}_{22} = \begin{pmatrix} Y_{221} \\ Y_{222} \\ \vdots \\ Y_{22p} \end{pmatrix} \quad \dots \quad \mathbf{Y}_{g2} = \begin{pmatrix} Y_{g21} \\ Y_{g22} \\ \vdots \\ Y_{g2p} \end{pmatrix}$$
$$\vdots$$
$$\mathbf{Y}_{1n_i} = \begin{pmatrix} Y_{1n_i1} \\ Y_{1n_i2} \\ \vdots \\ Y_{1n_ip} \end{pmatrix} \quad \mathbf{Y}_{2n_i} = \begin{pmatrix} Y_{2n_i1} \\ Y_{2n_i2} \\ \vdots \\ Y_{2n_ip} \end{pmatrix} \quad \dots \quad \mathbf{Y}_{gn_i} = \begin{pmatrix} Y_{gn_i1} \\ Y_{gn_i2} \\ \vdots \\ Y_{gn_ip} \end{pmatrix}$$

Source: PSU stat 505

- **Notation:** \mathbf{Y}_{ij} is the vector of variables for subject j in group i ; n_i is the sample size in group i ;
 $N = n_1 + n_2 + \dots + n_g$ the total sample size
- **Assumptions:** 1) common covariance matrix Σ ; 2) Independence; 3) Normality

Comparisons of Several Mean Vectors

CLEMSON UNIVERSITY

Comparisons of Two Mean Vectors

Multivariate Analysis of Variance

6.15

Notes

Test Statistics for MANOVA

- We are interested in testing the null hypothesis that the group mean vectors are all equal

$$H_0 : \mu_1 = \mu_2 = \dots = \mu_g.$$

The alternative hypothesis:

$$H_a : \mu_{ik} \neq \mu_{jk} \text{ for at least one } i \neq j \text{ and at least one variable } k$$

- Mean vectors:**

- Sample Mean Vector:** $\bar{\mathbf{y}}_i = \frac{1}{n_i} \mathbf{Y}_{ij}, \quad i = 1, \dots, g$

- Grand Mean Vector:** $\bar{\mathbf{y}}_{..} = \frac{1}{N} \sum_{i=1}^g \sum_{j=1}^{n_i} \mathbf{Y}_{ij}$

- Total Sum of Squares:**

$$T = \sum_{i=1}^g \sum_{j=1}^{n_i} (\mathbf{Y}_{ij} - \bar{\mathbf{y}}_{..})(\mathbf{Y}_{ij} - \bar{\mathbf{y}}_{..})^T$$

Notes

MANOVA Decomposition and MANOVA Table

$$\begin{aligned} T &= \sum_{i=1}^g \sum_{j=1}^{n_i} (\mathbf{Y}_{ij} - \bar{\mathbf{y}}_{..})(\mathbf{Y}_{ij} - \bar{\mathbf{y}}_{..})^T \\ &= \sum_{i=1}^g \sum_{j=1}^{n_i} [(\mathbf{Y}_{ij} - \bar{\mathbf{y}}_i) + (\bar{\mathbf{y}}_i - \bar{\mathbf{y}}_{..})][(\mathbf{Y}_{ij} - \bar{\mathbf{y}}_i) + (\bar{\mathbf{y}}_i - \bar{\mathbf{y}}_{..})]^T \\ &= \underbrace{\sum_{i=1}^g \sum_{j=1}^{n_i} (\mathbf{Y}_{ij} - \bar{\mathbf{y}}_i)(\mathbf{Y}_{ij} - \bar{\mathbf{y}}_i)^T}_E + \underbrace{\sum_{i=1}^g n_i (\bar{\mathbf{y}}_i - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_i - \bar{\mathbf{y}}_{..})^T}_H \end{aligned}$$

MANOVA Table

Source	df	SS
Treatment	$g - 1$	H
Error	$N - g$	E
Total	$N - 1$	T

Reject $H_0 : \mu_1 = \mu_2 = \dots = \mu_g$ if the matrix H is "large" relative to the matrix E

Notes

Test Statistics for MANOVA

There are several different test statistics for conducting the hypothesis test:

- Wilks Lambda**

$$\Lambda^* = \frac{|E|}{|H + E|}$$

Reject H_0 if Λ^* is "small"

- Hotelling-Lawley Trace**

$$T_0^2 = \text{trace}(\mathbf{H}\mathbf{E}^{-1})$$

Reject H_0 if T_0^2 is "large"

- Pillai Trace**

$$V = \text{trace}(\mathbf{H}(\mathbf{H} + \mathbf{E})^{-1})$$

Reject H_0 if V is "large"

Notes

Romano–British Pottery Example

```
> dat <- read.table("pottery.txt", header = F)
> out <- manova(cbind(V2, V3, V4, V5, V6) ~ V1, data = dat)
> summary(out, test = "Wilks")
      Df  Wilks approx F num Df den Df  Pr(>F)
V1      3 0.012301   13.088    15 50.091 1.84e-12 ***
Residuals 22
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
> summary(out)
      Df Pillai approx F num Df den Df  Pr(>F)
V1      3 1.5539   4.2984    15   60 2.413e-05 ***
Residuals 22
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Comparisons of
Several Mean
Vectors

CLEMSON
UNIVERSITY

Comparisons of
Two Mean Vectors

Multivariate
Analysis of
Variance

6.19

Notes

Notes

Notes
