

Lecture 1

Overview

DSA 8070 Multivariate Analysis
August 22-26, 2022

Whitney Huang
Clemson University



Notes

Agenda

- 1 Introduction
- 2 Objectives of Multivariate Analysis
- 3 Useful Tools for Multivariate Analysis



Notes

Introduction


- In many observational or experimental studies, measurements are collected simultaneously on **more than one variable** on each unit

```
> head(Boston)
  crim zn indus chas nox rm age dis rad tax ptratio black lstat medv
1 0.08632 18 2.31 0 0.538 6.575 65.2 4.0900 1 296 15.3 396.90 4.98 24.0
2 0.02731 0 7.07 0 0.469 6.421 78.9 4.9671 2 242 17.8 396.90 9.14 21.6
3 0.02729 0 7.07 0 0.469 7.185 61.1 4.9671 2 242 17.8 392.83 4.03 34.7
4 0.03237 0 2.18 0 0.458 6.998 45.8 6.0622 3 222 18.7 394.63 2.94 33.4
5 0.00905 0 2.18 0 0.458 7.147 54.2 6.0622 3 222 18.7 396.90 5.33 36.2
6 0.02985 0 2.18 0 0.458 6.430 58.7 6.0622 3 222 18.7 394.12 5.21 28.7
```
- **Multivariate analysis** is the collection of statistical methods that can be used to (jointly) analyze these multiple measurements
⇒ *some are extensions of familiar methods (t-test, ANOVA, linear regression,...) while others are unique to multivariate analysis*
- Idea is to exploit potential **"correlations"** among the multiple measurements to improve inference (see an example in the next slide)



Notes

Using Multivariate Methods Could Lead to Sharper Inference



Source: Fig. 1.1 of Applied Multivariate Statistics with R by Zelterman

Introduction

Objectives of Multivariate Analysis

Useful Tools for Multivariate Analysis

1.4

Notes

Objectives of Multivariate Analysis

Introduction

Objectives of Multivariate Analysis

Useful Tools for Multivariate Analysis

1.5

Notes

Dimensionality Reduction or Structural Simplification

- **Goal:** to reduce the “dimensionality” by considering a small number of (linear) combinations of a large number of measurements without losing important information
- **Examples:**
 - A single index of patient reaction to radiotherapy can be constructed from measurements on several response variables
 - Wildlife ecologists can construct a few indices of habitat preference from measurements of dozens of features of nesting sites selected by a certain bird species
- **Techniques:**
 - **Principal Component Analysis** (Week 9)
 - **Factor Analysis** (Week 10)
 - **Multidimensional Scaling** (Week 14)

Introduction

Objectives of Multivariate Analysis


Useful Tools for Multivariate Analysis

1.6

Notes

Grouping or Classification


- **Goal:** to **identify** groups of “similar” units or to **classify** units into previously defined groups
- **Examples:**
 - Using the concentration of elements (copper, silver, tin, antimony) in the lead alloy used in bullets, the FBI **identifies** ‘similar’ bullets that may be used to infer whether bullets were produced from the same batch of lead
 - The US IRS uses data collected from tax returns (income, amount withheld, deductions, ...) to **classify** taxpayers into two groups: those who will be audited and those who will not
- **Techniques:**
 - **Classification Analysis** (Week 12)
 - **Cluster Analysis** (Week 13)



Notes

Dependence among Variables and Prediction


- **Goal:** to estimate the relationship among variables and to predict the value of some of them given information on the others
- **Examples:**
 - The associations between measures of risk-taking propensity and measures of socioeconomic characteristics for top-level business executives were used to assess the relation between risk-taking behavior and performance
 - The association between test scores, and several college performance variables were used to develop predictors of success in college
- **Techniques:**
 - **Multivariate Regression** (Week 7)
 - **Repeated Measures Analysis** (Week 8)
 - **Canonical Correlation Analysis** (Week 11)



Notes

Hypothesis Testing

- **Goal:** to test if differences in sets of response mean vectors for two or more groups large enough to be distinguished from sampling variation
- **Examples:**
 - A transportation company wants to know if means for gasoline mileage, repair costs, downtime due to repairs differ for different truck models
 - An insurance company wants to know if changing case management practices leads to changes in mean length of hospital stay, mean infection rates, and mean costs
- **Techniques:**
 - **Hotelling's T^2 and MAVONA** (Week 5 and Week 6)




Notes

Useful Concepts/Tools for Multivariate Analysis

1.10

Notes

Exploratory Data Analysis [EDA, Tukey 1977]




1.11

Notes

Statistical Distance

Multivariate methods rely on “distances” between data points: **clustering** (group units that are “close”); **classification** (allocate each unit to the “closest” group)



Question: which one ((x_0, y_0) or (x_1, y_1)) is closer the center of the observations? \Rightarrow We will learn **Mahalanobis distance** to formally answer this question

1.12

Notes

Matrix Algebra (Week 3)

The study of multivariate methods is greatly facilitated by the use of matrix algebra

- Many operations performed on multivariate data are presented using vector/matrix notation, e.g., $\mathbf{X}_{n \times p}$ (Data matrix); $\hat{\boldsymbol{\mu}}_{p \times 1}$ (estimated mean vector); $\hat{\boldsymbol{\Sigma}}_{p \times p}$ (estimated covariance matrix)
- The computation of **eigenvalues** and **eigenvectors** (i.e., the **spectral decomposition**) plays an important role in multivariate analysis
- We will use \mathbb{R} to perform the needed matrix operations

Notes

Multivariate Normal Distribution (Week 4)

- We will often assume the joint distribution of $\mathbf{X} = (X_1, X_2, \dots, X_p)^T$ follows a multivariate normal distribution with the probability density function:

$$f(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{\frac{d}{2}} \det(\boldsymbol{\Sigma})^{\frac{1}{2}}} \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right]$$

- The multivariate normal assumption is often appropriate:
 - Variables can sometimes be assumed to be multivariate normal (perhaps after transformation)
 - **Central limit theorem** tells us that distribution of many **multivariate sample statistics** is approximately normal, regardless of the form of the population distribution

Notes

Data Mining, Machine Learning, and Multivariate Analysis

- **Data Mining** is the process of extracting and discovering patterns (e.g., unexpected structures or relationships, trends, clusters, and outliers) in **massive data sets**
- **Supervised learning** and **unsupervised learning** are two most common problems in **machine learning**
- Data mining/machine learning applications usually involve **many variables**, often **related in complex ways**, hence techniques from **multivariate analysis** play an important role

Notes