

Reading: Zelterman Chapter 8.1-8.4; DSA 8020 Lecture 12 [\[Link\]](#)

DSA 8070 Multivariate Analysis
October 11-October 15, 2021

Whitney Huang
Clemson University

Principle
Components
Analysis

CLEMSON
UNIVERSITY

Background

Finding Principal
Components

Principal
Components
Analysis in
Practice

9.1

3 Principal Components Analysis in Practice

Principle
Components
Analysis

CLEMSON
UNIVERSITY

Background

Finding Principal
Components

Principal
Components
Analysis in
Practice

9.2

- First introduced by [Karl Pearson \(1901\)](#) as a procedure for finding lines and planes which best fit a set of points in p -dimensional space

Y. K. Pearson (1991) *Curving lines and planar dimensional spaces* (1991). In *On Linear and Planar Curves* (Five Essays of Patrick J. Y. K. Pearson), ed. by K. H. Parker, D. S. J. University College, London, pp. 1–10.

- **Harold Hotelling (1933)** published a paper on PCA to find a smaller “fundamental set of independent variables” that “determines the values of the original set of p variables”

Principle
Components
Analysis

CLEMSON
UNIVERSITY

Background

Finding Principal
Components

Principal
Components
Analysis in
Practice

9.3

Basic Idea

Reduce the **dimensionality** of a data set in which there is a large number (i.e., p is "large") of inter-related variables while retaining as much as possible the **variation** in the original set of variables

- The reduction is achieved by transforming the original variables to a new set of variables, "**principal components**", that are **uncorrelated**
- These principal components are **ordered** such that **the first few retains most of the variation present in the data**
- **Goals/Objectives**
 - Reduction and summary
 - Study the structure of **covariance/correlation matrix**

Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components

Principal Components Analysis in Practice

94

Notes

Some Applications

- Interpretation (by studying the structure of covariance/correlation matrix)
- Select a sub-set of the original variables, that are uncorrelated to each other, to be used in other multivariate procedures (e.g., multiple regression, classification)
- Detect outliers or clusters of multivariate observations

Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components

Principal Components Analysis in Practice

95

Notes

Multivariate Data

We display a multivariate data that contains n units on p variables using a matrix

$$X = \begin{pmatrix} X_{1,1} & X_{2,1} & \cdots & X_{p,1} \\ X_{1,2} & X_{2,2} & \cdots & X_{p,2} \\ \vdots & \vdots & \ddots & \vdots \\ X_{1,n} & X_{2,n} & \cdots & X_{p,n} \end{pmatrix}$$

Summary Statistics

- **Mean Vector:** $\bar{X} = (\bar{X}_1, \bar{X}_2, \dots, \bar{X}_p)^T$
- **Covariance Matrix:** $\Sigma = \{\sigma_{ij}\}_{i,j=1}^p$, where $\sigma_{ii} = \text{Var}(X_i)$, $i = 1, \dots, p$ and $\sigma_{ij} = \text{Cov}(X_i, X_j)$, $i \neq j$

Next, we are going to discuss how to find **principal components**

Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components

Principal Components Analysis in Practice

96

Notes

Finding Principal Components

Principal Components (PCs) are uncorrelated **linear combinations** $\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_p$ determined sequentially, as follows:

- 1 The first PC is the linear combination $\tilde{X}_1 = \mathbf{c}_1^T \mathbf{X} = \sum_{i=1}^p c_{1i} X_i$ that maximize $\text{Var}(\tilde{X}_1)$ subject to $\mathbf{c}_1^T \mathbf{c}_1 = 1$

- 2 The second PC is the linear combination $\tilde{X}_2 = \mathbf{c}_2^T \mathbf{X} = \sum_{i=1}^p c_{2i} X_i$ that maximize $\text{Var}(\tilde{X}_2)$ subject to $\mathbf{c}_2^T \mathbf{c}_2 = 1$ and $\mathbf{c}_2^T \mathbf{c}_1 = 0$

\vdots

- p_{th} The p_{th} PC is the linear combination $\tilde{X}_p = \mathbf{c}_p^T \mathbf{X} = \sum_{i=1}^p c_{pi} X_i$ that maximize $\text{Var}(\tilde{X}_p)$ subject to $\mathbf{c}_p^T \mathbf{c}_p = 1$ and $\mathbf{c}_p^T \mathbf{c}_k = 0, \forall k < p$



Notes

Finding Principal Components by Decomposing Covariance Matrix

- Let Σ , the covariance matrix of \mathbf{X} , have eigenvalue-eigenvector pairs $(\lambda_i, \mathbf{e}_i)_{i=1}^p$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$. Then, the k_{th} principal component is given by

$$\tilde{X}_k = \mathbf{e}_k^T \mathbf{X} = e_{k1} X_1 + e_{k2} X_2 + \dots + e_{kp} X_p$$

\Rightarrow we can perform a single matrix operation to get the coefficients to form all the PCs!

- Then,

$$\text{Var}(\tilde{X}_i) = \lambda_i, \quad i = 1, \dots, p$$

Moreover $\text{Var}(\tilde{X}_1) \geq \text{Var}(\tilde{X}_2) \geq \dots \geq \text{Var}(\tilde{X}_p) \geq 0$

$$\text{Cov}(\tilde{X}_j, \tilde{X}_k) = 0, \quad \forall j \neq k$$

\Rightarrow different PCs are **uncorrelated** with each other



Notes

PCA and Proportion of Variance Explained

- It can be shown that

$$\sum_{i=1}^p \text{Var}(\tilde{X}_i) = \lambda_1 + \lambda_2 + \dots + \lambda_p = \sum_{i=1}^p \text{Var}(X_i)$$

- The proportion of the total variance associated with the k_{th} principal component is given by

$$\frac{\lambda_k}{\lambda_1 + \lambda_2 + \dots + \lambda_p}$$

- If a large proportion of the total population variance (say 80% or 90%) is explained by the first k PCs, then we can restrict attention to the first k PCs without much loss of information \Rightarrow **we achieve dimension reduction by considering $k < p$ uncorrelated components rather than the original p correlated variables**




Notes

Toy Example 1

Suppose we have $\mathbf{X} = (X_1, X_2)^T$ where $X_1 \sim N(0, 4)$, $X_2 \sim N(0, 1)$ are independent

- Total variation = $\text{Var}(X_1) + \text{Var}(X_2) = 5$
- X_1 axis explains 80% of total variation
- X_2 axis explains the remaining 20% of total variation




Notes

Toy Example 2

Suppose we have $\mathbf{X} = (X_1, X_2)^T$ where $X_1 \sim N(0, 4)$, $X_2 \sim N(0, 1)$ and $\text{Cor}(X_1, X_2) = 0.8$

- Total variation
= $\text{Var}(X_1) + \text{Var}(X_2) = \text{Var}(\tilde{X}_1) + \text{Var}(\tilde{X}_2) = 5$
- $\tilde{X}_1 = .9175X_1 + .3975X_2$ explains 93.9% of total variation
- $\tilde{X}_2 = .3975X_1 - .9176X_2$ explains the remaining 6.1% of total variation



Notes

PCs of Standardized versus Original Variables

If we use standardized variables, i.e.,

$Z_j = \frac{X_j - \mu_j}{\sqrt{\sigma_{jj}}}$, $j = 1, \dots, p$ ("z-scores"). Then we are going to work with the **correlation matrix** instead of the **covariance matrix** of $(X_1, \dots, X_p)^T$


- We can obtain PCs of standardized variables by applying spectral decomposition of the correlation matrix
- However, the PCs (and the proportion of variance explained) are, in general, different than those from original variables
- If units of p variables are comparable, covariance PCA may be more informative, if units of p variables are incomparable, correlation PCA may be more appropriate



Notes

Example: Men's 100k Road Race

The data consists of the times (in minutes) to complete successive 10k segments ($p = 10$) of the race. There are 80 racers in total ($n = 80$)



Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components

Principal Components Analysis in Practice

9.13

Notes

Eigenvalues of $\hat{\Sigma}$

	Eigenvalue	Proportion	Cumulative
PC1	735.77	0.75	0.75
PC2	98.47	0.10	0.85
PC3	53.27	0.05	0.90
PC4	37.30	0.04	0.94
PC5	26.04	0.03	0.97
PC6	17.25	0.02	0.98
PC7	8.03	0.01	0.99
PC8	4.25	0.00	1.00
PC9	2.40	0.00	1.00
PC10	1.29	0.00	1.00

Much of the total variance can be explained by the first three PCs

Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components


Principal Components Analysis in Practice

9.14

Notes

How Many Components to Retain?

A **scree plot** displays the variance explained by each component



Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components

Principal Components Analysis in Practice

9.15

Notes

Men's 100k Road Race Component Weights

	Comp.1	Comp.2	Comp.3
0-10 time	0.13	0.21	0.36
10-20 time	0.15	0.25	0.42
20-30 time	0.20	0.31	0.34
30-40 time	0.24	0.33	0.20
40-50 time	0.31	0.30	-0.13
50-60 time	0.42	0.21	-0.22
60-70 time	0.34	-0.05	-0.19
70-80 time	0.41	-0.01	-0.54
80-90 time	0.40	-0.27	0.15
90-100 time	0.39	-0.69	0.35

What these numbers mean?

Principle Components Analysis

CLEMSON UNIVERSITY

Background


Finding Principal Components

Principal Components Analysis in Practice

9.16

Notes

Visualizing the Weights to Gain Insight



First component: overall speed
Second component: contrast long and short races

Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components


Principal Components Analysis in Practice

9.17

Notes

Looking for Patterns

Mature runners: Age < 40 (M); Senior runners: Age >= 40 (S)



Principle Components Analysis

CLEMSON UNIVERSITY

Background


Finding Principal Components

Principal Components Analysis in Practice

9.18

Notes

Relating to Original Data: Profile Plot



Principle Components Analysis

CLEMSON UNIVERSITY

Background


Finding Principal Components

Principal Components Analysis in Practice

9.19

Notes

Correlation PCA versus Covariance PCA



Principle Components Analysis

CLEMSON UNIVERSITY

Background

Finding Principal Components

Principal Components Analysis in Practice

9.20

Notes

Notes
