

Lecture 26

Linear Contrasts

STAT 8010 Statistical Methods I
October 23, 2019

Whitney Huang
Clemson University



Notes

Facebook Friends Example Revisited

Too Much of a Good Thing? The Relationship Between Number of Friends and Interpersonal Impressions on Facebook

Stephanie Tom Tong
Brandon Van Der Heide
Lindsey Langwell

Department of Communication

Joseph B. Walther

Departments of Communication and Telecommunication, Information Studies & Media
Michigan State University

A central feature of the online social networking system, Facebook, is the connection to and links among friends. The sum of the number of one's friends is a feature displayed on users' profiles as a vestige of the friend connections a user has accrued. In contrast to offline social networks, individuals in online network systems frequently accrue friends numbering several hundred. The uncertain meaning of friend status in these systems raises questions about whether and how sociometric popularity conveys attractiveness in non-traditional, non-linear ways. An experiment examined the relationship between the number of friends a Facebook profile featured and observers' ratings of attractiveness and extraversion. A curvilinear effect of sociometric popularity and social attractiveness emerged, as did a quartic relationship between friend count and perceived extraversion. These results suggest that an overabundance of friend connections raises doubts about Facebook users' popularity and desirability.




Notes

Facebook Friends Example

There are 5 "treatment groups" in this study:

Participants examined one of five stimuli, each containing a Facebook profile mockup. Elements of these stimuli (e.g. photographs, wall posts, etc.) remained constant over the five versions, with the exception of the number of friends which appeared on the profile as 102, 302, 502, 702, or 902.... – Tong et al., 2008



Notes

Facebook Friends Example: Descriptive Statistics

	Score				
	102	302	502	702	902
Valid	24	33	26	30	21
Missing	0	0	0	0	0
Mean	3.817	4.879	4.562	4.407	3.990
Std. Deviation	0.999	0.851	1.070	1.428	1.023
Minimum	2.200	2.600	2.000	1.000	1.600
Maximum	6.000	6.400	6.800	7.000	5.200

Linear Contrasts

CLEMSON
UNIVERSITY

26.4

Notes

Facebook Friends: Overall F-Test

Question: Are Facebook attractiveness affected by # of friends?

$H_0 : \mu_1 = \mu_2 = \dots = \mu_5$
 H_a : At least one group mean is different from others

Analysis of Variance Table

Response: Score				
	Df	Sum Sq	Mean Sq	F value
Friends	4	19.89	4.9726	4.142
Residuals	129	154.87	1.2005	
	Pr(>F)			
Friends	0.00344	**		
Residuals				

Linear Contrasts

CLEMSON
UNIVERSITY

26.5

Notes

Facebook Friends Example: Fisher's LSD

We conclude that μ_i and μ_j differ at α level if $|\bar{X}_i - \bar{X}_j| > LSD$, where

$$LSD = t_{\alpha/2, df=N-J} \sqrt{MSE \left(\frac{1}{n_i} + \frac{1}{n_j} \right)}$$

> LSD_none\$groups		> LSD_bon\$groups	
Score groups		Score groups	
302	4.878788	a	302 4.878788 a
502	4.561538	ab	502 4.561538 ab
702	4.406667	abc	702 4.406667 ab
902	3.990476	bc	902 3.990476 b
102	3.816667	c	102 3.816667 b


Linear Contrasts

CLEMSON
UNIVERSITY

26.6

Notes

Me and the significant boys



Me and the significant boys after Bonferroni correction



Notes

Facebook Friends Example: Tukey's HSD Test

We conclude that μ_i and μ_j differ at α level if $|\bar{X}_i - \bar{X}_j| > \omega$, where

$$\omega = q_{\alpha}(J, N - J) \sqrt{\frac{MSE}{n}},$$

$q_{\alpha}(J, N - J)$ can be obtained from the [studentized range table](#)


Denominator DF	Number of Groups (s.k.a. Treatments)						
	3	4	5	6	7	8	9
51	3.414	3.756	3.999	4.187	4.340	4.469	4.580
52	3.412	3.753	3.996	4.184	4.337	4.465	4.576
53	3.410	3.751	3.994	4.181	4.334	4.462	4.572
54	3.408	3.749	3.991	4.178	4.331	4.459	4.569
55	3.406	3.747	3.989	4.176	4.328	4.455	4.566
56	3.405	3.745	3.986	4.173	4.325	4.452	4.562
57	3.403	3.743	3.984	4.170	4.322	4.449	4.559
58	3.402	3.741	3.982	4.168	4.319	4.447	4.556
59	3.400	3.739	3.979	4.165	4.317	4.444	4.553
60	3.399	3.737	3.977	4.163	4.314	4.441	4.550

Note: Here α is the FWER

Notes

Facebook Example: Tukey's HSD Test

	diff	lwr	upr	p adj
302-102	1.0621212	0.2488644	1.87537798	0.003889635
502-102	0.7448718	-0.1132433	1.60298691	0.121456224
702-102	0.5900000	-0.2402014	1.42020143	0.288431585
902-102	0.1738095	-0.7320145	1.07963355	0.984016816
502-302	-0.3172494	-1.1121910	0.47769215	0.804080046
702-302	-0.4721212	-1.2368466	0.29260420	0.432633745
902-302	-0.8883117	-1.7345313	-0.04209203	0.034535577
702-502	-0.1548718	-0.9671402	0.65739661	0.984391504
902-502	-0.5710623	-1.4604793	0.31835479	0.391768065
902-702	-0.4161905	-1.2787075	0.44632652	0.669927748



Notes

Linear Contrasts

Suppose we have J populations (e.g. response for J different treatments) of interest. We have seen how to perform multiple comparisons. For example, the comparison between μ_1 and μ_2 can be conducted using the test: $H_0 : \mu_1 - \mu_2 = 0$ vs. $H_a : \mu_1 - \mu_2 \neq 0$. This comparison is actually a special case of **linear contrasts**

Linear Contrasts

Let c_1, c_2, \dots, c_J are constants where $\sum_{j=1}^J c_j = 0$, then $\sum_{j=1}^J c_j \mu_j$ is called a **linear contrast** of the population means.

Example: Suppose $J = 4$

- ➊ $\mu_1 - \mu_3 : c_1 = 1, c_2 = 0, c_3 = -1, c_4 = 0$
- ➋ $\mu_2 - \mu_4 : c_1 = 0, c_2 = 1, c_3 = 0, c_4 = -1$
- ➌ $\mu_1 - \frac{1}{3}\mu_2 - \frac{1}{3}\mu_3 - \frac{1}{3}\mu_4 : c_1 = 1, c_2 = c_3 = c_4 = -\frac{1}{3}$

Notes

Inferences for Linear Contrasts

If we want to make a inference about $L = \sum_{j=1}^J c_j \mu_j$. Then we use

$$\hat{L} = \sum_{j=1}^J c_j \bar{X}_j$$

as the point estimate. Furthermore, we can construct a $100(1 - \alpha)\%$ CI for L :

$$(\hat{L} - t_{(\alpha/2, df=N-J)} \hat{se}_{\hat{L}}, \hat{L} + t_{(\alpha/2, df=N-J)} \hat{se}_{\hat{L}}),$$

$$\text{where } \hat{se}_{\hat{L}} = \sqrt{\text{MSE} \left(\frac{c_1^2}{n_1} + \dots + \frac{c_J^2}{n_J} \right)}$$

To test whether L is significantly different from 0, we can conduct the following test:

$$H_0 : \sum_{j=1}^J c_j \mu_j = 0 \text{ vs. } H_a : \sum_{j=1}^J c_j \mu_j \neq 0$$

Notes

Hypothesis Testing for Linear Contrasts

- ➊ Null and Alternative Hypotheses:

$$H_0 : \sum_{j=1}^J c_j \mu_j = 0 \text{ vs. } H_a : \sum_{j=1}^J c_j \mu_j \neq 0$$

- ➋ Test Statistic:

$$t_{obs} = \frac{\hat{L} - 0}{\hat{se}_{\hat{L}}} = \frac{\sum_{j=1}^J c_j \bar{X}_j}{\sqrt{\text{MSE} \left(\frac{c_1^2}{n_1} + \dots + \frac{c_J^2}{n_J} \right)}}$$

- ➌ Decision:

Reject H_0 if $|t_{obs}| > t_{\alpha/2, df=N-J}$ (or p-value $< \alpha$)

Notes

Facebook Example: Linear Contrast

Suppose we'd like to compare μ_1 vs. $\frac{\mu_3 + \mu_4}{2}$. Let $L = 1\mu_1 - \frac{1}{2}\mu_3 - \frac{1}{2}\mu_4$. Then the above comparison is equivalent to test whether L is different from 0

1 $H_0 : L = 0$ vs. $H_a : L \neq 0$

2 $t_{obs} = \frac{\hat{l}}{se_L} = \frac{1 \times 3.817 - 0.5 \times 4.562 - 0.5 \times 4.407}{\sqrt{1.2005 \times (\frac{1^2}{24} + \frac{0.5^2}{26} + \frac{0.5^2}{30})}} = \frac{-0.6674}{0.2675} = -2.495$

3 Since $|t_{obs}| = |-2.495| = 2.495 > t_{0.025, df=129} = 1.9785$.
We reject H_0 at 0.05 level

Note: If we are performing several tests for different linear contrasts, we'll need to adjust α level accordingly to control the FWER

Notes

Notes

Notes
