

# Lecture 8

## Normal Random Variables

Text: Chapter 4

*STAT 8010 Statistical Methods I*  
February 3, 2020

Whitney Huang  
Clemson University

# Agenda

## 1 Normal Density Curves

## 2 Standard Normal

## 3 Sums of Normal Random Variables

# Probability Density Curve for Normal Random Variable


Normal Random  
Variables

CLEMSON  
UNIVERSITY

Normal Density  
Curves


Standard Normal

Sums of Normal  
Random Variables



# Normal Density Curves

Different  $\mu$  but same  $\sigma^2$




Normal Density  
Curves

Standard Normal


Sums of Normal  
Random Variables

## Normal Density Curves Cont'd

Same  $\mu$  but different  $\sigma^2$




# Normal Density Curves




- The parameter  $\mu$  determines the center of the distribution

# Normal Density Curves



- The parameter  $\mu$  determines the center of the distribution
- The parameter  $\sigma^2$  determines the spread of the distribution

# Normal Density Curves



- The parameter  $\mu$  determines the center of the distribution
- The parameter  $\sigma^2$  determines the spread of the distribution
- Also called bell-shaped distribution



# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value  $\Phi\left(\frac{x-\mu}{\sigma}\right)$  for  $-\infty < x < \infty$  from standard normal table

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value  $\Phi\left(\frac{x-\mu}{\sigma}\right)$  for  $-\infty < x < \infty$  from standard normal table



# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value  $\Phi\left(\frac{x-\mu}{\sigma}\right)$  for  $-\infty < x < \infty$  from standard normal table
- The expected value:  $\mathbb{E}[X] = \mu$

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value  $\Phi\left(\frac{x-\mu}{\sigma}\right)$  for  $-\infty < x < \infty$  from standard normal table
- The expected value:  $\mathbb{E}[X] = \mu$

# Characteristics of Normal Random Variables

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Parameters:  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value  $\Phi\left(\frac{x-\mu}{\sigma}\right)$  for  $-\infty < x < \infty$  from standard normal table
- The expected value:  $\mathbb{E}[X] = \mu$
- The variance:  $Var(X) = \sigma^2$

## Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

- Normal random variable  $X$  with mean  $\mu$  and standard deviation  $\sigma$  can convert to standard normal  $Z$  by the following :

$$Z = \frac{X - \mu}{\sigma}$$

## Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

- Normal random variable  $X$  with mean  $\mu$  and standard deviation  $\sigma$  can convert to standard normal  $Z$  by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by  $\Phi(z)$ , can be found from the standard normal table

## Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

- Normal random variable  $X$  with mean  $\mu$  and standard deviation  $\sigma$  can convert to standard normal  $Z$  by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by  $\Phi(z)$ , can be found from the standard normal table

## Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

- Normal random variable  $X$  with mean  $\mu$  and standard deviation  $\sigma$  can convert to standard normal  $Z$  by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by  $\Phi(z)$ , can be found from the standard normal table
- The probability  $\mathbb{P}(a \leq X \leq b)$  where  $X \sim N(\mu, \sigma^2)$  can be compute


$$\begin{aligned}\mathbb{P}(a \leq X \leq b) &= \mathbb{P}\left(\frac{a - \mu}{\sigma} \leq Z \leq \frac{b - \mu}{\sigma}\right) \\ &= \Phi\left(\frac{b - \mu}{\sigma}\right) - \Phi\left(\frac{a - \mu}{\sigma}\right)\end{aligned}$$

# Standard Normal (Z) Table

Example:

If  $z^* = 1.96$ , then

$P(-\infty \text{ to } z^*) = 0.9750$



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767




# Standard Normal (Z) Table Cont'd

Example:

If  $z^* = 1.96$ , then

$P(-\infty \text{ to } z^*) = 0.9750$



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	<b>0.9750</b>	0.9756	0.9761	0.9767

# Standard Normal (Z) Table Cont'd


Example:

If  $z^* = 1.96$ , then

$P(-\infty \text{ to } z^*) = 0.9750$

$$P(0 < Z < 1.96) = 0.9750 - 0.5$$

$$P(0 < Z < 1.96) = 0.4750$$



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	<b>0.9750</b>	0.9756	0.9761	0.9767

# Properties of $\Phi$

Normal Random  
Variables



Normal Density  
Curves

**Standard Normal**

Sums of Normal  
Random Variables

- $\Phi(0) = .50 \Rightarrow$  Mean and Median ( $50_{th}$  percentile) for standard normal are both 0

- $\Phi(0) = .50 \Rightarrow$  Mean and Median ( $50_{th}$  percentile) for standard normal are both 0

- $\Phi(0) = .50 \Rightarrow$  Mean and Median ( $50_{th}$  percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$

- $\Phi(0) = .50 \Rightarrow$  Mean and Median ( $50_{th}$  percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$


- $\Phi(0) = .50 \Rightarrow$  Mean and Median ( $50_{th}$  percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$
- $\mathbb{P}(Z > z) = 1 - \Phi(z) = \Phi(-z)$



## The Empirical Rules

The **Empirical Rules** provide a quick way to approximate certain probabilities for the Normal Distribution as the following table:

Interval	Percentage with interval
$\mu \pm \sigma$	68%
$\mu \pm 2\sigma$	95%
$\mu \pm 3\sigma$	99.7%



## Example

Let us examine  $Z$ . Find the following probabilities with respect to  $Z$ :

1  $Z$  is at most  $-1.75$  ▶

2  $Z$  is between  $-2$  and  $2$  inclusive ▶

3  $Z$  is less than  $.5$  ▶

## Example Cont'd

**Solution.**

$$\textcircled{1} \quad \mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401 \quad \textcircled{\leftarrow}$$

### Solution.

$$1 \quad \mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401 \quad \leftarrow$$

$$2 \quad \mathbb{P}(-2 \leq Z \leq 2) = \Phi(2) - \Phi(-2) = .9772 - .0228 = .9544 \quad \leftarrow$$

### Solution.

$$1 \quad \mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401$$

$$2 \quad \mathbb{P}(-2 \leq Z \leq 2) = \Phi(2) - \Phi(-2) = .9772 - .0228 = .9544$$

$$3 \quad \mathbb{P}(Z < .5) = \Phi(.5) = .6915$$

## Example

Suppose a STAT-8020 exam score follows a normal distribution with mean 78 and variance 36. Let  $X$  to denote the exam score, answer the following questions:

- 1 What is the probability that a randomly chosen test taker got a score greater than 84?
- 2 Suppose the passing score for this exam is 75. What is the probability that a randomly chosen test taker got a score greater than 84 given that she/he pass the exam?
- 3 Using the empirical rule to find the 84<sub>th</sub> percentile.

## Example

Find the following percentile with respect to  $Z$


1  $10_{th}$  percentile 


2  $55_{th}$  percentile 


3  $90_{th}$  percentile 

## Example Cont'd

### Solution.

1  $Z_{10} = -1.28$  

2  $Z_{55} = 0.13$  

3  $Z_{90} = 1.28$  



## Example

Let  $X$  be Normal with a mean of 20 and a variance of 49. Find the following probabilities and percentile:

- 1  $X$  is between 15 and 23 ▶
- 2  $X$  is more than 30 ▶
- 3  $X$  is more than 12 knowing it is less than 20 ▶
- 4 What is the value that is smaller than 20% of the distribution? ▶

## Example Cont'd

### Solution.

$$\textcircled{1} \quad \mathbb{P}(15 \leq X \leq 23) = \Phi\left(\frac{15-20}{7}\right) - \Phi\left(\frac{23-20}{7}\right) = \Phi(0.43) - \Phi(-0.71) = .6664 - .2389 = .4275 \quad \leftarrow$$

$$\textcircled{2} \quad \mathbb{P}(X > 30) = 1 - \mathbb{P}(X \leq 30) = 1 - \Phi\left(\frac{30-20}{7}\right) = 1 - .9236 = .0764 \quad \leftarrow$$

$$\textcircled{3} \quad \mathbb{P}(X > 12 | X < 20) = \frac{\mathbb{P}(12 < X < 20)}{\mathbb{P}(X < 20)} = \frac{\Phi(0) - \Phi(-1.14)}{\Phi(0)} = .7458 \quad \leftarrow$$

$$\textcircled{4} \quad Z_{80} = 0.84 \Rightarrow X_{80} = \mu + Z_{80} \times \sigma = 20 + 0.84 \times \sqrt{49} = 25.88 \quad \leftarrow$$

# Sums of Normal Random Variables

If  $X_i$   $1 \leq i \leq n$  are independent normal random variables with mean  $\mu_i$  and variance  $\sigma_i^2$ , respectively.

If  $X_i$   $1 \leq i \leq n$  are independent normal random variables with mean  $\mu_i$  and variance  $\sigma_i^2$ , respectively.

- Let  $S_n = \sum_{i=1}^n X_i$  then  $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$

If  $X_i$   $1 \leq i \leq n$  are independent normal random variables with mean  $\mu_i$  and variance  $\sigma_i^2$ , respectively.


- Let  $S_n = \sum_{i=1}^n X_i$  then  $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$


If  $X_i$   $1 \leq i \leq n$  are independent normal random variables with mean  $\mu_i$  and variance  $\sigma_i^2$ , respectively.


- Let  $S_n = \sum_{i=1}^n X_i$  then  $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$
- This can be applied for any integer  $n$

## Example

Let  $X_1$ ,  $X_2$ , and  $X_3$  be mutually independent, Normal random variables. Let their means and standard deviations be  $3k$  and  $k$  for  $k = 1, 2$ , and  $3$  respectively. Find the following distributions:

1  $\sum_{i=1}^3 X_i$  

2  $X_1 + 2X_2 - 3X_3$  

3  $X_1 + 5X_3$  

### Solution.

1  $\sum_{i=1}^3 X_i \sim N(\mu = 3 + 6 + 9 = 15, \sigma^2 = 1^2 + 2^2 + 3^2 = 14)$  ◀

2  $X_1 + 2X_2 - 3X_3 \sim N(\mu = 3 + 12 - 27 = -12, \sigma^2 = 1^2 + 4 \times 2^2 + 9 \times 3^2 = 98)$  ◀

3  $X_1 + 5X_3 \sim N(\mu = 3 + 45 = 48, \sigma^2 = 1^2 + 25 \times 3^2 = 226)$  ◀