

Lecture 11

Extreme Value Analysis

Readings: An Introduction to Statistical Modeling of Extreme Values, Stuart Coles, 2001

MATH 8090 Time Series Analysis
October 26 & October 28, 2021

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary
Extreme Value
Analysis

Whitney Huang
Clemson University

Agenda

1

Motivation

Motivation

2

Extreme Value Theorem & Block Maxima Method

EVT

Peaks-Over-
Threshold (POT)
Method

3

Peaks-Over-Threshold (POT) Method

Non-stationary
Extreme Value
Analysis

4

Non-stationary Extreme Value Analysis

Motivation

EVT

Peaks–Over–
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis



1 Motivation

2 Extreme Value Theorem & Block Maxima Method

3 Peaks–Over–Threshold (POT) Method

4 Non-stationary Extreme Value Analysis

Extreme Rainfall During Hurricane Harvey



Source: NASA (Left); National Weather Service (Right)

- “A storm forces Houston, the limitless city, to consider its limits” – The New York Times (8.31.17)


Motivation

EVT

Peaks-Over-Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

Environmental Extremes: Heatwaves, Storm Surges, etc.



- **Heat wave:** The 2003 European heat wave led to the hottest summer on record in Europe since 1540 that resulted in at least **30,000 deaths**
- **Storm Surge:** Hurricane Katrina produced the highest storm surge ever recorded (**27.8 feet**) on the U.S. coast

Motivation

EVT

Peaks-Over-Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

Why Study Extremes?

Although infrequent, extremes usually have large impact.

Goal: to quantify the tail behavior \Rightarrow often requires extrapolation.

Applications:

- Hydrology: flooding
- Climate: temperature, precipitation, wind, ...
- Finance
- Insurance/reinsurance
- Engineering: structural design, reliability

Motivation

EVT

Peaks-Over-Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

- How to estimate the magnitude of extreme events (e.g. 100-year rainfall)?
- How extremes vary in space?
- How extremes may change in future climate conditions?

Motivation

EVT

Peaks–Over–
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

1 Motivation

2 Extreme Value Theorem & Block Maxima Method

3 Peaks–Over–Threshold (POT) Method

4 Non-stationary Extreme Value Analysis


Usual vs Extremes

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis



Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Probability Framework

Let $X_1, \dots, X_n \stackrel{i.i.d.}{\sim} F$ and define $M_n = \max\{X_1, \dots, X_n\}$
Then the distribution function of M_n is

$$\begin{aligned}\mathbb{P}(M_n \leq x) &= \mathbb{P}(X_1 \leq x, \dots, X_n \leq x) \\ &= \mathbb{P}(X_1 \leq x) \times \dots \times \mathbb{P}(X_n \leq x) = F^n(x)\end{aligned}$$

Remark

$$F^n(x) \xrightarrow{n \rightarrow \infty} \begin{cases} 0 & \text{if } F(x) < 1 \\ 1 & \text{if } F(x) = 1 \end{cases}$$

⇒ the limiting distribution is degenerate.

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

Recall the Central Limit Theorem:

$$\frac{S_n - n\mu}{\sqrt{n}\sigma} \xrightarrow{d} N(0, 1),$$

where $S_n = \sum_{i=1}^n X_i$

⇒ rescaling is the key to obtain a non-degenerate distribution

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Asymptotic: Classical Limit Laws

Recall the Central Limit Theorem:

$$\frac{S_n - n\mu}{\sqrt{n}\sigma} \xrightarrow{d} N(0, 1),$$

where $S_n = \sum_{i=1}^n X_i$

⇒ rescaling is the key to obtain a non-degenerate distribution

Question: Can we get the limiting distribution of

$$\frac{M_n - b_n}{a_n}$$

for suitable sequence $\{a_n\} > 0$ and $\{b_n\}$?

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

CLT in Action

- ① Generate 100 (n) random numbers from an Exponential distribution (population distribution)
- ② Compute the sample mean of these 100 random numbers
- ③ Repeat this process 120 times

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

Motivation

EVT

Peaks-Over-Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

then G must be the same type of the following form:

$$G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]_+^{-\frac{1}{\xi}}\right\}$$

where $x_+ = \max(x, 0)$ and $G(x)$ is the distribution function of the **generalized extreme value distribution (GEV(μ, σ, ξ)**)

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

then G must be the same type of the following form:

$$G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]_+^{-\frac{1}{\xi}}\right\}$$

where $x_+ = \max(x, 0)$ and $G(x)$ is the distribution function of the **generalized extreme value distribution (GEV(μ, σ, ξ)**)

- μ and σ are location and scale parameters

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

then G must be the same type of the following form:

$$G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]_+^{-\frac{1}{\xi}}\right\}$$

where $x_+ = \max(x, 0)$ and $G(x)$ is the distribution function of the **generalized extreme value distribution (GEV(μ, σ, ξ)**)

- μ and σ are location and scale parameters
- ξ is a shape parameter determining the rate of tail decay, with

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

then G must be the same type of the following form:

$$G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]_+^{-\frac{1}{\xi}}\right\}$$

where $x_+ = \max(x, 0)$ and $G(x)$ is the distribution function of the **generalized extreme value distribution (GEV(μ, σ, ξ)**)

- μ and σ are location and scale parameters
- ξ is a shape parameter determining the rate of tail decay, with
 - $\xi > 0$ giving the heavy-tailed case (Fréchet)

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

then G must be the same type of the following form:

$$G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]_+^{-\frac{1}{\xi}}\right\}$$

where $x_+ = \max(x, 0)$ and $G(x)$ is the distribution function of the **generalized extreme value distribution (GEV(μ, σ, ξ)**)

- μ and σ are location and scale parameters
- ξ is a shape parameter determining the rate of tail decay, with
 - $\xi > 0$ giving the heavy-tailed case (**Fréchet**)
 - $\xi = 0$ giving the light-tailed case (**Gumbel**)

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Extremal Types Theorem (Fisher–Tippett 1928, Gnedenko 1943)

Define $M_n = \max\{X_1, \dots, X_n\}$ where $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} F$. If $\exists a_n > 0$ and $b_n \in \mathbb{R}$ such that, as $n \rightarrow \infty$, if

$$\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) \xrightarrow{d} G(x)$$

then G must be the same type of the following form:

$$G(x; \mu, \sigma, \xi) = \exp\left\{-\left[1 + \xi\left(\frac{x - \mu}{\sigma}\right)\right]_+^{-\frac{1}{\xi}}\right\}$$

where $x_+ = \max(x, 0)$ and $G(x)$ is the distribution function of the **generalized extreme value distribution (GEV(μ, σ, ξ)**)

- μ and σ are location and scale parameters
- ξ is a shape parameter determining the rate of tail decay, with
 - $\xi > 0$ giving the heavy-tailed case (**Fréchet**)
 - $\xi = 0$ giving the light-tailed case (**Gumbel**)
 - $\xi < 0$ giving the bounded-tailed case (**reversed Weibull**)

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Example: Exponential Maxima

Let $X \sim \text{Exp}(\lambda = 1)$. Set $a_n = 1$, $b_n = \log(n)$. We want to show $\frac{M_n - b_n}{a_n}$ converges to a GEV distribution, where $M_n = \max_{i=1}^n X_i$.

$$\begin{aligned}\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) &= \mathbb{P}(M_n \leq a_n x + b_n) \\ &= \mathbb{P}(M_n \leq x + \log(n)) \\ &= (1 - \exp(-x - \log(n)))^n \\ &= \left(1 - \frac{1}{n} \exp(-x)\right)^n \\ &\xrightarrow{n \rightarrow \infty} \exp(-\exp(x))\end{aligned}$$

It is the cdf of the standard Gumbel distribution

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Extremal Types Theorem in Action

- ① Generate 100 (n) random numbers from an Exponential distribution (population distribution)
- ② Compute the **sample maximum** of these 100 random numbers
- ③ Repeat this process 120 times

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Definition

A distribution G is said to be **max-stable** if

$$G^k(a_k x + b_k) = G(x), \quad k \in \mathbb{N}$$

for some constants $a_k > 0$ and b_k

- Taking powers of a distribution function results only in a change of location and scale
- A distribution is **max-stable** \iff it is a **GEV** distribution

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

- Quantiles of GEV

$$\begin{aligned} G(x_p) &= \exp \left\{ - \left[1 + \xi \left(\frac{x_p - \mu}{\sigma} \right) \right]_+^{-\frac{1}{\xi}} \right\} = 1 - p \\ \Rightarrow x_p &= \mu - \frac{\sigma}{\xi} \left[1 - \{-\log(1-p)\}^{-\xi} \right] \quad 0 < p < 1 \end{aligned}$$

- In the extreme value terminology, x_p is the return level associated with the return period $\frac{1}{p}$

Assume n is large enough so that

$$\begin{aligned}\mathbb{P}\left(\frac{M_n - b_n}{a_n} \leq x\right) &\approx \exp(-[1 + \xi x]^{-1/\xi}) \\ \Rightarrow \mathbb{P}(M_n \leq y) &\approx \exp\left(-\left[1 + \xi\left(\frac{y - b_n}{a_n}\right)^{-1/\xi}\right]\right) \\ &= \exp\left(-\left[1 + \xi\left(\frac{y - \mu}{\sigma}\right)\right]^{-1/\xi}\right)\end{aligned}$$

Then, we have a three-parameter estimation problem. μ, σ, ξ can be estimated via **maximum likelihood**

Motivation

EVT

Peaks-Over-Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

Clemson Daily Precipitation [Data Source: USHCN]

Extreme Value
Analysis




Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis


Daily Precip in Clemson



Block Maxima Method (Gumbel 1958)

1. Determine the block size and extract the block maxima

Daily Precip in Clemson



Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis


Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Block Maxima Method (Gumbel 1958)

1. Determine the block size and extract the block maxima




Block Maxima Method (Gumbel 1958)

2. Fit the GEV to the maximal and assess the fit

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary
Extreme Value Analysis

Block Maxima Method (Gumbel 1958)

2. Fit the GEV to the maximal and assess the fit


Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

Quantile Plot




Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Block Maxima Method (Gumbel 1958)

3. Perform inference for return levels, probabilities, etc. Two methods to quantify estimation uncertainty: 1) Delta-method (relies on asymptotic normality); 2) Profile likelihood



Motivation

EVT

Peaks–Over–
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

1 Motivation

2 Extreme Value Theorem & Block Maxima Method

3 Peaks–Over–Threshold (POT) Method


4 Non-stationary Extreme Value Analysis

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

Recall the Block Maxima Method




Question: Can we use data more efficiently?

Peaks-over-threshold (POT) method [Davison & Smith 1990]

1. Select a “sufficiently large” threshold u , extract the exceedances

Daily Precip in Clemson



Peaks-over-threshold (POT) method [Davison & Smith 1990]


2. Fit an appropriate model to exceedances

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis



If $M_n = \max_{i=1,\dots,n} X_i$ (for a large n) can be approximated by a GEV(μ, σ, ξ), then for sufficiently large u ,

$$\begin{aligned}\mathbb{P}(X_i > x + u | X_i > u) &= \frac{n\mathbb{P}(X_i > x + u)}{n\mathbb{P}(X_i > u)} \\ &\rightarrow \left(\frac{1 + \xi \frac{x+u-b_n}{a_n}}{1 + \xi \frac{u-b_n}{a_n}} \right)^{-\frac{1}{\xi}} \\ &= \left(1 + \frac{\xi x}{a_n + \xi(u - b_n)} \right)^{-\frac{1}{\xi}}\end{aligned}$$

⇒ Survival function of **generalized Pareto distribution**

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Pickands–Balkema–de Haan Theorem (1974, 1975)

If $M_n = \max_{1 \leq i \leq n} \{X_i\} \approx \text{GEV}(\mu, \sigma, \xi)$, then, for a “large” u (i.e., $u \rightarrow x_F = \sup\{x : F(x) < 1\}$),

$$\mathbb{P}(X > u) \approx \frac{1}{n} \left[1 + \xi \left(\frac{u - \mu}{\sigma} \right) \right]^{\frac{-1}{\xi}}$$

$F_u = \mathbb{P}(X - u < y | X > u)$ is well approximated by the generalized Pareto distribution (GPD). That is:

$$F_u(y) \xrightarrow{d} H_{\tilde{\sigma}, \xi}(y) \quad u \rightarrow x_F$$

where

$$H_{\tilde{\sigma}, \xi}(y) = \begin{cases} 1 - (1 + \xi y / \tilde{\sigma})^{-1/\xi} & \xi \neq 0; \\ 1 - \exp(-y / \tilde{\sigma}) & \xi = 0. \end{cases}$$

and $\tilde{\sigma} = \sigma + \xi(u - \mu)$

Motivation

EVT


Peaks-Over-Threshold (POT) Method

Non-stationary
Extreme Value
Analysis

How to Choose the Threshold?

Bias-variance tradeoff:


- Threshold too low \Rightarrow bias because of the model asymptotics being invalid
- Threshold too high \Rightarrow variance is large due to few data points



Task: To choose a u_0 s.t. the Mean Residual Life curve behaves linearly $\forall u > u_0$

Peaks-over-threshold (POT) method [Davison & Smith 1990]

2. Fit an appropriate model to exceedances and assess the fit



Motivation


EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

Peaks-over-threshold (POT) method [Davison & Smith 1990]

3. Perform inference for return levels, probabilities, etc



Temporal Dependence

Question: Is the GEV still the limiting distribution for block maxima of a stationary (but not independent) sequence $\{X_i\}$?

Answer: Yes, as long as mixing conditions hold. ([Leadbetter et al., 1983](#))

What does this mean for inference?

Block maximum approach: GEV still correct for marginal. Since block maximum data likely have negligible dependence, proceed as usual

Threshold exceedance approach: GPD is correct for the marginal. If extremes occur in clusters, estimation affected as likelihood assumes independence of threshold exceedances

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

Motivation

EVT

Peaks-Over-
Threshold (POT)
MethodNon-stationary
Extreme Value
Analysis

- To estimate the tail, EVT uses only extreme observations
- Shape parameter ξ is extremely important but hard to estimate
- Threshold exceedance approaches allow the user to retain more data than block-maximum approaches, thereby reducing the uncertainty with parameter estimates
- Temporal dependence in the data is more of an issue in threshold exceedance models. One can either decluster, or alternatively, adjust inference

Outline

1 Motivation

Motivation

EVT

Peaks–Over–
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

2 Extreme Value Theorem & Block Maxima Method

3 Peaks–Over–Threshold (POT) Method

4 Non-stationary Extreme Value Analysis

- $M_t \sim \text{GEV}(\mu(t), \sigma(t), \xi(t))$
- Typically assume “simple” structure for $\mu(t)$ and $\sigma(t)$, and $\xi(t)$ be a constant
- $\mu(t)$ and $\sigma(t)$ could depend on some relevant factors

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

- Extreme value theory provides a framework to model extreme values
 - GEV for fitting block maxima
 - GPD for fitting threshold exceedances
 - Return level for communicating risk
- Practical Issues:

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis

- Extreme value theory provides a framework to model extreme values
 - GEV for fitting block maxima
 - GPD for fitting threshold exceedances
 - Return level for communicating risk
- Practical Issues:** seasonality, temporal dependence, non-stationarity, ...

Motivation

EVT

Peaks-Over-Threshold (POT) Method

Non-stationary Extreme Value Analysis

For Further Reading

-  J. Beirlant, Y Goegebeur, J. Segers, and J Teugels
Statistics of Extremes: Theory and Applications.
Wiley, 2004.
-  L. de Haan, and A. Ferreira
Extreme Value Theory: An Introduction.
Springer, 2006.
-  S. I. Resnick
Heavy-Tail Phenomena: Probabilistic and Statistical Modeling. Springer, 2007.

Motivation

EVT

Peaks-Over-
Threshold (POT)
Method

Non-stationary
Extreme Value
Analysis