

Lecture 15



Interpolation of Spatial Data

DSA 8020 Statistical Methods II

April 19-23, 2021


Whitney Huang
Clemson University

Toy Examples of Spatial Interpolation



Question: What is your best guess of the value of the missing pixel, denoted as $Y(s_0)$, for each case?

Interpolating Paraná State Precipitation Data



Goal: To interpolate the values in the spatial domain

The Spatial Interpolation Problem

Given observations of a spatially varying quantity Y at n spatial locations

$$y(s_1), y(s_2), \dots, y(s_n), \quad s_i \in \mathcal{S}, i = 1, \dots, n$$

We want to estimate this quantity at any **unobserved location**

$$Y(s_0), \quad s_0 \in \mathcal{S}$$

Applications

- Mining: ore grade
- Climate: temperature, precipitation, ...
- Remote Sensing: CO₂ retrievals
- Environmental Science: air pollution levels, ...

Some History of Spatial Statistics

- Mining (Krige 1951)
Matheron (1960s),
Forestry (Matérn
1960)



- More recent work:
Cressie (1993) Stein
(1999)



1 Gaussian Process Spatial Model

2 Spatial Interpolation

3 Parameter estimation

The best guess (in a statistical sense) should be based on the conditional distribution $[Y(s_0) | Y = \mathbf{y}]$ where

$$\mathbf{y} = (y(s_1), \dots, y(s_n))^T$$

- Calculating this conditional distribution can be difficult
- Instead we use a **linear predictor**:

$$\hat{Y}(s_0) = \lambda_0 + \sum_{i=1}^n \lambda_i y(s_i)$$

- The best linear predictor is completely determined by the **mean** and **covariance** of $\{Y(s), s \in \mathcal{S}\}$, and the observations \mathbf{y}

We assume that the observed data $\{y(\mathbf{s}_i)\}_{i=1}^n$ is one partial realization of a (continuously indexed) spatial GP $\{Y(\mathbf{s})\}_{\mathbf{s} \in \mathcal{S}}$.

Model:

$$Y(\mathbf{s}) = m(\mathbf{s}) + \epsilon(\mathbf{s}), \quad \mathbf{s} \in \mathcal{S} \subset \mathbb{R}^d$$

where

- Mean function:

$$m(\mathbf{s}) = \mathbb{E}[Y(\mathbf{s})] = \mathbf{X}^T(\mathbf{s})\boldsymbol{\beta}$$

- Covariance function:

$$\{\epsilon(\mathbf{s})\}_{\mathbf{s} \in \mathcal{S}} \sim \text{GP}(0, K(\cdot, \cdot)), \quad K(\mathbf{s}_1, \mathbf{s}_2) = \text{Cov}(\epsilon(\mathbf{s}_1), \epsilon(\mathbf{s}_2))$$

In practice, the covariance must be estimated from the data $(y(s_1), \dots, y(s_n))^T$. We need to impose some structural assumptions

- Stationarity:

$$\begin{aligned} K(s_1, s_2) &= \text{Cov}(\epsilon(s_1), \epsilon(s_2)) = C(s_1 - s_2) \\ &= \text{Cov}(\epsilon(s_1 + h), \epsilon(s_2 + h)) \end{aligned}$$

- Isotropy:

$$K(s_1, s_2) = \text{Cov}(\epsilon(s_1), \epsilon(s_2)) = C(\|s_1 - s_2\|)$$

A Valid Covariance Function Must Be Positive Definite (p.d.)!


A covariance function is positive definite if

$$\sum_{i,j=1}^n a_i a_j C(\mathbf{s}_i - \mathbf{s}_j) \geq 0$$

for any finite locations $\mathbf{s}_1, \dots, \mathbf{s}_n$, and for any constants a_i ,
 $i = 1, \dots, n$

Question: what is the consequence if a covariance function is NOT p.d.? \Rightarrow **We can get a negative variance**

Question: How to guarantee a $C(\cdot)$ is p.d.?

- Using a **parametric covariance function** (see some examples in next slide)
- Using **Bochner's Theorem**  to construct a valid covariance function

Some Commonly Used Covariance Functions

- **Powered exponential:**

$$C(h) = \sigma^2 \exp\left(-\left(\frac{h}{\rho}\right)^\alpha\right), \quad \sigma^2 > 0, \rho > 0, 0 < \alpha \leq 2$$

- **Spherical:**

$$C(h) = \sigma^2 \left(1 - 1.5 \frac{h}{\rho} + 0.5 \left(\frac{h}{\rho}\right)^3\right) 1_{\{h \leq \rho\}}, \quad \sigma^2, \rho > 0$$

Note: it is only valid for 1, 2, and 3 dimensional spatial domain.

- **Matérn:**

$$C(h) = \sigma^2 \frac{(\sqrt{2\nu}h/\rho)^\nu \mathcal{K}_\nu(\sqrt{2\nu}h/\rho)}{\Gamma(\nu)2^{\nu-1}}, \quad \sigma^2 > 0, \rho > 0, \nu > 0$$

“Use the Matérn model” – Stein (1999, pp. 14)

1-D Realizations from Matérn Model with Fixed σ^2, ρ




Figure: courtesy of Rasmussen & Williams 2006

The larger ν is, the smoother the process is

2-D Realizations from Matérn Model with Fixed σ^2


Interpolation of
Spatial Data

CLEMSON
UNIVERSITY

Gaussian Process
Spatial Model

Spatial Interpolation

Parameter estimation



1 Gaussian Process Spatial Model

2 Spatial Interpolation

3 Parameter estimation

If

$$\begin{pmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{pmatrix} \sim N \left(\begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{pmatrix}, \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix} \right)$$

Then

$$[\mathbf{Y}_1 | \mathbf{Y}_2 = \mathbf{y}_2] \sim N(\boldsymbol{\mu}_{1|2}, \Sigma_{1|2})$$

where

$$\boldsymbol{\mu}_{1|2} = \boldsymbol{\mu}_1 + \Sigma_{12} \Sigma_{22}^{-1} (\mathbf{y}_2 - \boldsymbol{\mu}_2)$$

$$\Sigma_{1|2} = \Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21}$$

If $\{Y(\mathbf{s})\}_{\mathbf{s} \in \mathcal{S}}$ follows a GP, then

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} m_0 \\ \mathbf{m} \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & k^T \\ k & \Sigma \end{pmatrix} \right)$$

We have

$$[Y_0 | \mathbf{Y} = \mathbf{y}] \sim \mathcal{N}(m_{Y_0 | \mathbf{Y} = \mathbf{y}}, \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2)$$

where

$$m_{Y_0 | \mathbf{Y} = \mathbf{y}} = m_0 + k^T \Sigma^{-1} (\mathbf{y} - \mathbf{m})$$

$$\sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 = \sigma_0^2 - k^T \Sigma^{-1} k$$

Spatial Prediction of Paraná State Rainfall

Interpolation of
Spatial Data


CLEMSON
UNIVERSITY

Gaussian Process
Spatial Model


Spatial Interpolation

Parameter estimation

Prediction



Uncertainty



If $\{Y(\mathbf{s})\}_{\mathbf{s} \in \mathcal{S}}$ follows a GP, then

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} m_0 \\ \mathbf{m} \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & k^T \\ k & \Sigma \end{pmatrix} \right)$$

We have

$$[Y_0 | \mathbf{Y} = \mathbf{y}] \sim \mathcal{N}(m_{Y_0 | \mathbf{Y} = \mathbf{y}}, \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2)$$

where

$$\begin{aligned} m_{Y_0 | \mathbf{Y} = \mathbf{y}} &= m_0 + k^T \Sigma^{-1} (\mathbf{y} - \mathbf{m}) \\ \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 &= \sigma_0^2 - k^T \Sigma^{-1} k \end{aligned}$$

Question: what if we don't know $m_0, \mathbf{m}, \sigma_0^2, \Sigma$?

\Rightarrow We need to estimate the mean and covariance from the data \mathbf{y} .

Gaussian Process
Spatial Model

Spatial Interpolation

Parameter estimation

1 Gaussian Process Spatial Model




2 Spatial Interpolation

3 Parameter estimation

Recap: Gaussian Process

We assume that the observed data $\{y(s_i)\}_{i=1}^n$ is one partial realization of a (continuously indexed) spatial stochastic process $\{Y(s)\}_{s \in \mathcal{S}}$.

- Gaussian Processes $GP(m(\cdot), K(\cdot, \cdot))$ are widely used in modeling spatial stochastic processes
- Spatial statisticians often focus on the covariance function.
e.g. $K(h) = \sigma^2 \frac{(\sqrt{2\nu}h/\gamma)^\nu \mathcal{K}_\nu(\sqrt{2\nu}h/\gamma)}{\Gamma(\nu)2^{\nu-1}}$



Under the stationary and isotropic assumptions


Variogram:

$$\begin{aligned}2\gamma(\mathbf{s}_i, \mathbf{s}_j) &= \text{Var}(Y(\mathbf{s}_i) - Y(\mathbf{s}_j)) \\&= \text{E} \left\{ ((Y(\mathbf{s}_i) - \mu(\mathbf{s}_i)) - (Y(\mathbf{s}_j) - \mu(\mathbf{s}_j)))^2 \right\} \\&= \text{E} \left\{ (Y(\mathbf{s}_i) - Y(\mathbf{s}_j))^2 \right\} \\&= 2\gamma(\|\mathbf{s}_i - \mathbf{s}_j\|) = 2\gamma(h)\end{aligned}$$

Semivariogram and covariance function:

$$\gamma(h) = C(0) - C(h)$$


Semivariogram $\{\frac{1}{2}\text{Var}(\varepsilon(s_i) - \varepsilon(s_j))\}_{i,j}$



Source: `fields` vignette by Wiens and Krock, 2019

Estimation: Weighted Least Squares Method

$$\operatorname{argmin}_{\boldsymbol{\theta}} \sum_{u \in \mathcal{U}} \frac{N(h_u)}{[\gamma(h_u; \boldsymbol{\theta})]^2} [\hat{\gamma}(h_u) - \gamma(h_u; \boldsymbol{\theta})]^2$$



Maximum Likelihood Estimation (MLE)

Log-likelihood:

Given data $\mathbf{y} = (y(\mathbf{s}_1), \dots, y(\mathbf{s}_n))^T$

$$\ell_n(\boldsymbol{\beta}, \boldsymbol{\theta}; \mathbf{y}) \propto -\frac{1}{2} \log |\boldsymbol{\Sigma}_{\boldsymbol{\theta}}| - \frac{1}{2} (\mathbf{y} - \mathbf{X}^T \boldsymbol{\beta})^T [\boldsymbol{\Sigma}_{\boldsymbol{\theta}}]_{n \times n}^{-1} (\mathbf{y} - \mathbf{X}^T \boldsymbol{\beta})$$

where $\boldsymbol{\Sigma}_{\boldsymbol{\theta}}(i, j) = \sigma^2 \rho_{\rho, \nu}(\|\mathbf{s}_i - \mathbf{s}_j\|) + \tau^2 1_{\{\mathbf{s}_i = \mathbf{s}_j\}}, i, j = 1, \dots, n$

Maximum Likelihood Estimation (MLE)

Log-likelihood:

Given data $\mathbf{y} = (y(\mathbf{s}_1), \dots, y(\mathbf{s}_n))^T$

$$\ell_n(\boldsymbol{\beta}, \boldsymbol{\theta}; \mathbf{y}) \propto -\frac{1}{2} \log |\boldsymbol{\Sigma}_{\boldsymbol{\theta}}| - \frac{1}{2} (\mathbf{y} - \mathbf{X}^T \boldsymbol{\beta})^T [\boldsymbol{\Sigma}_{\boldsymbol{\theta}}]_{n \times n}^{-1} (\mathbf{y} - \mathbf{X}^T \boldsymbol{\beta})$$

where $\boldsymbol{\Sigma}_{\boldsymbol{\theta}}(i, j) = \sigma^2 \rho_{\rho, \nu}(\|\mathbf{s}_i - \mathbf{s}_j\|) + \tau^2 1_{\{\mathbf{s}_i = \mathbf{s}_j\}}, i, j = 1, \dots, n$

for any fixed $\boldsymbol{\theta}_0 \in \Theta$ the unique value of $\boldsymbol{\beta}$ that maximizes ℓ_n is given by

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}_0}^{-1} \mathbf{X})^{-1} \mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}_0} \mathbf{y}$$

Then we obtain the profile log likelihood

$$\ell_n(\boldsymbol{\theta}; \mathbf{y}) \propto -\frac{1}{2} \log |\boldsymbol{\Sigma}_{\boldsymbol{\theta}}| - \frac{1}{2} \mathbf{y}^T P(\boldsymbol{\theta}) \mathbf{y}$$

where

$$P(\boldsymbol{\theta}) = \boldsymbol{\Sigma}_{\boldsymbol{\theta}}^{-1} - \boldsymbol{\Sigma}_{\boldsymbol{\theta}}^{-1} \mathbf{X} (\mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}}^{-1} \mathbf{X})^{-1} \mathbf{X}^T \boldsymbol{\Sigma}_{\boldsymbol{\theta}}$$

Solve the maximization problem above to get the MLE

A complex-valued function C on \mathbb{R}^d is the covariance function for a weakly stationary mean square continuous complex-valued random process on \mathbb{R}^d if and only if it can be represented as

$$C(\mathbf{h}) = \int_{\mathbb{R}^d} \exp(i\omega^T \mathbf{h}) F(d\omega),$$

with F a positive finite measure. When F has a density with respect to Lebesgue measure, we have the spectral density f and

$$f(\omega) = \frac{1}{2\pi} \int_{\mathbb{R}^d} \exp(-i\omega^T \mathbf{h}) C(\mathbf{h}) d\mathbf{h}$$