

# Lecture 6

## Probability II

Text: Chapter 4

STAT 8010 Statistical Methods I  
September 8, 2020

Whitney Huang  
Clemson University

Probability II

CLEMSON  
UNIVERSITY

Union,  
Intersection, and  
Logical  
Relationships  
among Events

Complement Rule  
and General  
Addition Rule

Independence and  
Conditional  
Probability

Law of Total  
Probability

Bayes' Rule

6.1

Notes

---

---

---

---

---

---

---

### Agenda

- 1 Union, Intersection, and Logical Relationships among Events
- 2 Complement Rule and General Addition Rule
- 3 Independence and Conditional Probability
- 4 Law of Total Probability
- 5 Bayes' Rule

Probability II

CLEMSON  
UNIVERSITY

Union,  
Intersection, and  
Logical  
Relationships  
among Events

Complement Rule  
and General  
Addition Rule

Independence and  
Conditional  
Probability

Law of Total  
Probability

Bayes' Rule

6.2

Notes

---

---

---

---

---

---

---

### Probability Rules

- Any probability must be between 0 and 1 inclusively
- The sum of the probabilities for all the experimental outcomes must equal 1

If a probability model satisfies the two rules above, it is said to be **legitimate**

Probability II

CLEMSON  
UNIVERSITY

Union,  
Intersection, and  
Logical  
Relationships  
among Events

Complement Rule  
and General  
Addition Rule

Independence and  
Conditional  
Probability

Law of Total  
Probability

Bayes' Rule

6.3

Notes

---

---

---

---

---

---

---

Example

An experiment with three outcomes has been repeated 50 times, and it was learned that outcome 1 occurred 20 times, outcome 2 occurred 13 times, and outcome 3 occurred 17 times. Assign probabilities to the outcomes. What method did you use?

Solution.

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

64

Notes

---

---

---

---

---

---

---

Example

A decision maker subjectively assigned the following probabilities to the four possible outcomes of an experiment:

$$P(E_1) = 0.1 \ P(E_2) = 0.15 \ P(E_3) = 0.4 \ P(E_4) = 0.2$$

Are these probability assignments legitimate? Explain.

Solution.

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

65

Notes

---

---

---

---

---

---

---

Union, Intersection, and Logical Relationships among Events

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

66

Notes

---

---

---

---

---

---

---

Intersection and Union

- **Intersection:** the intersection of two events  $A$  and  $B$ , denoted by  $A \cap B$ , is the event that contains all outcomes of  $A$  that also belong to  $B \Rightarrow$  **AND**

Example: Let  $A = \{1, 2, 3\}$  and  $B = \{1, 2, 4, 5\}$ , then  $A \cap B = \{1, 2\}$

- **Union:** the union of two events  $A$  and  $B$ , denoted by  $A \cup B$ , is the event of all outcomes that belong to either  $A$  or  $B \Rightarrow$  **OR**

Example: Let  $A = \{1, 2, 3\}$  and  $B = \{1, 2, 4, 5\}$ , then  $A \cup B = \{1, 2, 3, 4, 5\}$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.7

Notes

---

---

---

---

---

---

---

Example

Suppose we flipped 3 fair coins. Let  $A$  be the event of **exactly 2 tails**. Let  $B$  be the event that the **first 2 tosses are tails**. Let  $C$  be the event that **all 3 tosses are tails**. What are  $A \cap B$ ,  $A \cup C$ , and  $(A \cap B) \cup C$ ?

**Solution.**

$$A = \{(T, T, H), (T, H, T), (H, T, T)\}$$
$$B = \{(T, T, T), (T, T, H)\}$$
$$C = \{T, T, T\}$$

- $A \cap B = \{(T, T, H)\}$
- $A \cup C = \{(T, T, H), (T, H, T), (H, T, T), (T, T, T)\}$
- $(A \cap B) \cup C = \{(T, T, H) \cup \{T, T, T\} = \{(T, T, H), (T, T, T)\}$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.8

Notes

---

---

---

---

---

---

---

Logical Relationships among Events

- **Mutually exclusive:** refers to two (or more) events that cannot both occur when the random experiment is formed.

$$A \cap B = \emptyset$$

- **Exhaustive:** refers to event(s) that comprise the sample space.

$$A \cup B = \Omega$$

- **Partition:** events that are both mutually exclusive and exhaustive.

$$A \cap B = \emptyset \quad \text{and} \quad A \cup B = \Omega$$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.9

Notes

---

---

---

---

---

---

---

# Complement Rule and General Addition Rule

Probability II
CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events
Complement Rule and General Addition Rule
Independence and Conditional Probability
Law of Total Probability
Bayes' Rule

6.10

Notes

---

---

---


---

---

---

---

## Complement



Probability II
CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events
Complement Rule and General Addition Rule
Independence and Conditional Probability
Law of Total Probability
Bayes' Rule

6.11

Notes

---

---

---

---

---

---

---

## Complement Rule

- 1 By the definition of complement

$$A \cup A^c = \Omega$$

- 2 Apply the probability operator

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(\Omega) = 1$$

- 3 Since  $A$  and  $A^c$  are mutually exclusive

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(A) + \mathbb{P}(A^c)$$

- 4 Hence we get 

$\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$

Probability II
CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events
Complement Rule and General Addition Rule
Independence and Conditional Probability
Law of Total Probability
Bayes' Rule

6.12

Notes

---

---

---

---

---

---

---

Example

Suppose we rolled a fair, six-sided die 10 times. Let  $T$  be the event that we roll at least 1 three. If one were to calculate  $T$  you would need to find the probability of 1 three, 2 threes,  $\dots$ , and 10 threes and add them all up. However, you can use the complement rule to calculate  $\mathbb{P}(T)$

Solution.

Let  $X$  be the times that we rolled a 3, then  
 $\mathbb{P}(T) = \mathbb{P}(X \geq 1) =$   
 $\mathbb{P}(X = 1) + \mathbb{P}(X = 2) + \dots + \mathbb{P}(X = 10)$   
need to compute 10 probabilities  
If we apply the complement rule  
 $\mathbb{P}(T) = 1 - \mathbb{P}(T^c) = 1 - \mathbb{P}(X = 0)$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.13

Notes

---

---

---

---

---


---

---

---

Venn Diagram

A Venn diagram is a diagram that shows all possible logical relations between a finite collection of events.



Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.14

Notes

---

---

---

---

---

---


---

---

General Addition Rule

The general addition rule is a way of finding the probability of a union of 2 events. It is

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$



Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.15

Notes

---

---

---

---

---

---

---

---

Example

Three of the major commercial computer operating systems are Windows, Mac OS, and Red Hat Linux Enterprise. A Computer Science professor selects 50 of her students and asks which of these three operating systems they use. The results for the 50 students are summarized below.

- 30 students use Windows
- 16 students use at least two of the operating systems
- 9 students use all three operating systems
- 18 students use Mac OS
- 46 students use at least one of the operating systems
- 11 students use both Windows and Linux
- 11 students use both Windows and Mac OS

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.16

Notes

---

---

---


---

---

---

---

Example cont'd



Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.17

Notes

---

---

---

---

---

---

---

Independence and Conditional Probability

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.18

Notes

---

---

---

---

---

---

---

## Independence: A Motivating Example

### Example

You toss a fair coin and it comes up "Heads" three times. What is the chance that the next toss will also be a "Head"?



Notes

---

---

---

---

---

---

---

## Independence and Conditional Probability

### Conditional Probability


Let  $A$  and  $B$  be events. The probability that event  $B$  occurs **given** (knowing) that event  $A$  occurs is called a **conditional probability** and is denoted by  $P(B|A)$ . The formula of conditional probability is

$$P(B|A) = \frac{P(B \cap A)}{P(A)}$$

### Independent events

Suppose  $P(A) > 0$ ,  $P(B) > 0$ . We say that event  $B$  is **independent** of event  $A$  if the occurrence of event  $A$  does not affect the probability that event  $B$  occurs.

$$P(B|A) = P(B) \Rightarrow P(B \cap A) = P(B)P(A)$$



Notes

---

---

---

---

---

---

---

## Law of Partitions & Multiplication Rule

### Law of partitions

Let  $A_1, A_2, \dots, A_k$  form a partition of  $\Omega$ . Then, for all events  $B$ ,

$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(A_i \cap B)$$


### Multiplication rule

- 2 events:

$$\mathbb{P}(B \cap A) = \mathbb{P}(A) \times \mathbb{P}(B|A) = \mathbb{P}(B) \times \mathbb{P}(A|B)$$

- More than 2 events:

$$\begin{aligned} \mathbb{P}(\cap_{i=1}^n A_i) &= \mathbb{P}(A_1) \times \mathbb{P}(A_2|A_1) \times \mathbb{P}(A_3|A_1 \cap A_2) \\ &\quad \times \dots \times \mathbb{P}(A_n|A_{n-1} \cap \dots \cap A_1) \end{aligned}$$



Notes

---

---

---

---

---

---

---

Law of Total Probability

Let  $A_1, A_2, \dots, A_k$  form a partition of  $\Omega$ . Then, for all events  $B$ ,

$$\begin{aligned} \mathbb{P}(B) &= \sum_{i=1}^k \underbrace{\mathbb{P}(A_i \cap B)}_{\text{Law of partitions}} \\ &= \sum_{i=1}^k \underbrace{\mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}_{\text{Multiplication rule}} \end{aligned}$$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.22

Notes

---

---

---

---

---

---

---

Example

Suppose that two factories supply light bulbs to the market. Factory X's bulbs work for over 5000 hours in 99% of cases, whereas factory Y's bulbs work for over 5000 hours in 95% of cases. It is known that factory X supplies 60% of the total bulbs available and Y supplies 40% of the total bulbs available. What is the chance that a purchased bulb will work for longer than 5000 hours?

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.23

Notes

---

---

---

---

---

---

---

Bayes' Rule: Motivating example

The Monty Hall Problem

There was an old television show called Let's Make a Deal, whose original host was named Monty Hall. The set-up is as follows. You are on a game show and you are given the choice of three doors. Behind one door is a car, behind the others are goats. You pick a door, and the host, who knows what is behind the doors, opens another door (not your pick) which has a goat behind it. Then he asks you if you want to change your original pick. The question we ask you is, "Is it to your advantage to switch your choice?"

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.24

Notes

---

---

---

---

---

---

---



The Monty Hall Problem



Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.25

Notes

---

---

---

---

---

---

---

---

The Monty Hall Problem Solution



Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.26

Notes

---

---

---

---

---

---

---

---

Bayes' Rule

General form

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

Let  $A_1, A_2, \dots, A_k$  form a partition of the sample space. Then for every event  $B$  in the sample space,

$$\mathbb{P}(A_j|B) = \frac{\mathbb{P}(B|A_j) \times \mathbb{P}(A_j)}{\sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}, \quad j = 1, 2, \dots, k$$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.27

Notes

---

---

---

---

---

---

---

---

Example

Let us assume that a specific disease is only present in 5 out of every 1,000 people. Suppose that the test for the disease is accurate 99% of the time a person has the disease and 95% of the time that a person lacks the disease. What is the probability that the person has the disease given that they tested positive?

Solution.

$$\mathbb{P}(D|+) = \frac{\mathbb{P}(D \cap +)}{\mathbb{P}(+)} = \frac{.005 \times .99}{.005 \times .99 + .995 \times .05} = \frac{.00495}{.0547} = .0905$$

The reason we get such a surprising result is because the disease is so rare that the number of false positives greatly outnumbers the people who truly have the disease.

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.28

Notes

---

---

---

---

---

---

---

---

Review of Probability (we learned so far)

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection, Mutually Exclusive, Exhaustive, Partition
- Venn Diagram
- Independence and Conditional Probability

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.29

Notes

---

---

---

---

---

---

---

---

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$  for any event  $A$ ,  $\mathbb{P}(\emptyset) = 0$ ,  $\mathbb{P}(\Omega) = 1$
- Complement rule:  $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule:  
 $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:  
 $\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$
- Conditional probability:  $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- Law of total probability:  
 $\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(B \cap A_i) = \sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)$
- Independence: if  $A$  and  $B$  are independent, then  
 $\mathbb{P}(A|B) = \mathbb{P}(A)$ ,  $\mathbb{P}(B|A) = \mathbb{P}(B)$ , and  
 $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$

Probability II

CLEMSON UNIVERSITY

Union, Intersection, and Logical Relationships among Events

Complement Rule and General Addition Rule

Independence and Conditional Probability

Law of Total Probability

Bayes' Rule

6.30

Notes

---

---

---

---

---

---

---

---