

Lecture 36

Simple Linear Regression

STAT 8010 Statistical Methods I
November 20, 2019

Whitney Huang
Clemson University

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis

Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.1

Notes

Agenda

- 1 What is regression analysis
- 2 Simple Linear Regression (SLR)
- 3 Parameter Estimation in SLR

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis

Simple Linear Regression (SLR)


Parameter Estimation in SLR

36.2

Notes

What is Regression Analysis?

Regression analysis: A set of statistical procedures for estimating the relationship between **response variable** and **predictor variable(s)**



We will focus on **simple linear regression** in the next few lectures

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis


Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.3

Notes

Scatterplot: Is Linear Trend Reasonable?



```
> cov(x, y)
[1] 3.042785
> cor(x, y)
[1] 0.9622804
```

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis

Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.4

Notes

Simple Linear Regression (SLR)

- Y: dependent (response) variable; X: independent (predictor) variable
- In SLR we **assume** there is a **linear relationship** between X and Y:
$$Y = \beta_0 + \beta_1 X + \varepsilon$$
 - We will need to estimate β_0 (intercept) and β_1 (slope)
 - Then we can use the estimated regression equation to
 - make predictions
 - study the relationship between response and predictor
 - control the response
 - Yet we need to quantify our uncertainty regarding the linear relationship

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis


Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.5

Notes

Regression equation: $Y = \beta_0 + \beta_1 X$



- β_0 : $E[Y]$ when $X = 0$
- β_1 : $E[\Delta Y]$ when X increases by 1

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis

Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.6

Notes

Assumptions about the Random Error ε

In order to estimate β_0 and β_1 , we make the following assumptions about ε

- $E[\varepsilon_i] = 0$
- $\text{Var}[\varepsilon_i] = \sigma^2$
- $\text{Cov}[\varepsilon_i, \varepsilon_j] = 0, \quad i \neq j$

Therefore, we have

$$E[Y_i] = \beta_0 + \beta_1 X_i, \text{ and}$$
$$\text{Var}[Y_i] = \sigma^2$$

The regression line $\beta_0 + \beta_1 X$ represents the **conditional expectation curve** whereas σ^2 measures the magnitude of the **variation** around the regression curve



Notes

Estimation: Method of Least Square

For the given observations $(x_i, y_i)_{i=1}^n$, choose β_0 and β_1 to minimize the *sum of squared errors*:

$$L(\beta_0, \beta_1) = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

Solving the above minimization problem requires some knowledge from Calculus....

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{(X_i - \bar{X})^2}$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X}$$

We also need to **estimate** σ^2

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{n-2}, \text{ where } \hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i$$



Notes

Properties of Least Squares Estimates

- **Gauss-Markov** theorem states that in a linear regression these least squares estimators

1. **Are unbiased**, i.e.,

- $E[\hat{\beta}_1] = \beta_1; E[\hat{\beta}_0] = \beta_0$
- $E[\hat{\sigma}^2] = \sigma^2$

2. Have **minimum variance** among all unbiased linear estimators

Note that we do not make any distributional assumption on ε_i



Notes

Example: Maximum Heart Rate vs. Age

The maximum heart rate MaxHeartRate of a person is often said to be related to age Age by the equation:

MaxHeartRate = 220 – Age.

Suppose we have 15 people of varying ages are tested for their maximum heart rate (bpm) (link to the “dataset”: <http://whitneyhuang83.github.io/maxHeartRate.csv>)

- 1 Compute the estimates for the regression coefficients
- 2 Compute the fitted values
- 3 Compute the estimate for σ

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis

Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.10

Notes

Estimate the Parameters β_1 , β_0 , and σ^2

Y_i and X_i are the Maximum Heart Rate and Age of the i^{th} individual

- To obtain $\hat{\beta}_1$
 - 1 Compute $\bar{Y} = \frac{\sum_{i=1}^n Y_i}{n}$, $\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$
 - 2 Compute $Y_i - \bar{Y}$, $X_i - \bar{X}$, and $(X_i - \bar{X})^2$ for each observation
 - 3 Compute $\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})$ divided by $\sum_{i=1}^n (X_i - \bar{X})^2$
- $\hat{\beta}_0$: Compute $\bar{Y} - \hat{\beta}_1 \bar{X}$
- $\hat{\sigma}^2$
 - 1 Compute the fitted values:
 $\hat{Y}_i = \hat{\beta}_0 + \hat{\beta}_1 X_i$, $i = 1, \dots, n$
 - 2 Compute the **residuals** $e_i = Y_i - \hat{Y}_i$, $i = 1, \dots, n$
 - 3 Compute the **residual sum of squares (RSS)** $= \sum_{i=1}^n (Y_i - \hat{Y}_i)^2$ and divided by $n - 2$ (why?)

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis

Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.11

Notes

Let's Do the Calculations

$$\bar{X} = \frac{15}{15} \frac{18 + 23 + \dots + 39 + 37}{15} = 37.33$$
$$\bar{Y} = \frac{15}{15} \frac{202 + 186 + \dots + 183 + 178}{15} = 180.27$$

X	18	23	25	35	65	54	34	66	72	19	23	42	18	39	37
Y	202	186	187	180	156	169	174	172	153	199	193	174	198	183	178
	-19.33	-14.33	-12.33	-2.33	-27.67	-16.67	-3.33	-16.67	-34.67	-16.33	-14.33	-4.67	-19.33	-1.67	-9.33
	21.73	6.73	6.73	-0.27	-24.27	-11.27	-6.27	-6.27	-27.27	18.73	12.73	-6.27	17.73	2.73	-2.27
	-420.18	-42.18	-83.04	0.82	-671.38	-167.78	20.89	-154.31	-945.24	-343.44	-182.51	-29.24	-342.84	4.56	0.76
	373.78	205.44	152.11	5.44	765.44	277.78	11.11	348.44	1201.78	336.11	295.44	21.78	373.78	5.78	0.11
	195.69	191.70	190.11	182.13	158.20	166.97	182.93	165.38	152.61	194.89	191.70	176.54	195.69	178.94	180.53

- $\hat{\beta}_1 = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2} = -0.7977$
- $\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X} = 210.0485$
- $\hat{\sigma}^2 = \frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{13} = 20.9563 \Rightarrow \hat{\sigma} = 4.5778$

Simple Linear Regression

CLEMSON UNIVERSITY

What is regression analysis



Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.12

Notes

Let's Double Check

Output from  ( Studio)

```
> fit <- lm(MaxHeartRate ~ Age)
> summary(fit)


Call:
lm(formula = MaxHeartRate ~ Age)

Residuals:
    Min       1Q   Median       3Q      Max
-8.9258 -2.5383  0.3879  3.1867  6.6242

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 210.04846   2.86694   73.27  < 2e-16 ***
Age         -0.79773    0.06996  -11.40 3.85e-08 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.578 on 13 degrees of freedom
Multiple R-squared:  0.9091,    Adjusted R-squared:  0.9021
F-statistic: 130 on 1 and 13 DF,  p-value: 3.848e-08
```

Simple Linear Regression



What is regression analysis


Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.13


Notes

Linear Regression Fit



Question: Is linear relationship between max heart rate and age reasonable? ⇒ [Residual Analysis](#)

Simple Linear Regression



What is regression analysis

Simple Linear Regression (SLR)

Parameter Estimation in SLR

36.14

Notes

Notes
