

# Lecture 7

## Probability III

Text: Chapter 4

STAT 8010 Statistical Methods I  
September 10, 2020

Whitney Huang  
Clemson University



Notes

---

---

---

---

---

---

---

### Agenda

- 1 Law of Total Probability
- 2 Bayes' Rule
- 3 Random Variables



Notes

---

---

---

---

---

---

---

### Review: Independence and Conditional Probability

#### Conditional Probability

Let  $A$  and  $B$  be events. The probability that event  $B$  occurs **given** (knowing) that event  $A$  occurs is called a **conditional probability** and is denoted by  $P(B|A)$ . The formula of conditional probability is

$$P(B|A) = \frac{P(B \cap A)}{P(A)}$$

#### Independent events

Suppose  $P(A) > 0$ ,  $P(B) > 0$ . We say that event  $B$  is **independent** of event  $A$  if the occurrence of event  $A$  does not affect the probability that event  $B$  occurs.

$$P(B|A) = P(B) \Rightarrow P(B \cap A) = P(B)P(A)$$



Notes

---

---

---

---

---

---

---

Law of Partitions & Multiplication Rule

Law of partitions

Let  $A_1, A_2, \dots, A_k$  form a partition of  $\Omega$ . Then, for all events  $B$ ,

$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(A_i \cap B)$$

Multiplication rule

- 2 events:

$$\mathbb{P}(B \cap A) = \mathbb{P}(A) \times \mathbb{P}(B|A) = \mathbb{P}(B) \times \mathbb{P}(A|B)$$

- More than 2 events:

$$\begin{aligned} \mathbb{P}(\cap_{i=1}^n A_i) &= \mathbb{P}(A_1) \times \mathbb{P}(A_2|A_1) \times \mathbb{P}(A_3|A_1 \cap A_2) \\ &\quad \times \dots \times \mathbb{P}(A_n|A_{n-1} \cap \dots \cap A_1) \end{aligned}$$

Probability III

CLEMSON  
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

74

Notes

---

---

---

---

---

---

---

Law of Total Probability

Let  $A_1, A_2, \dots, A_k$  form a partition of  $\Omega$ . Then, for all events  $B$ ,

$$\begin{aligned} \mathbb{P}(B) &= \sum_{i=1}^k \underbrace{\mathbb{P}(A_i \cap B)}_{\text{Law of partitions}} \\ &= \sum_{i=1}^k \underbrace{\mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}_{\text{Multiplication rule}} \end{aligned}$$

Probability III

CLEMSON  
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

75

Notes

---

---

---

---

---

---

---

Example

Suppose that two factories supply light bulbs to the market. Factory X's bulbs work for over 5000 hours in 99% of cases, whereas factory Y's bulbs work for over 5000 hours in 95% of cases. It is known that factory X supplies 60% of the total bulbs available and Y supplies 40% of the total bulbs available. What is the chance that a purchased bulb will work for longer than 5000 hours?

Probability III

CLEMSON  
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

76

Notes

---

---

---

---

---

---

---

Bayes' Rule: Motivating example

The Monty Hall Problem

There was an old television show called Let's Make a Deal, whose original host was named Monty Hall. The set-up is as follows. You are on a game show and you are given the choice of three doors. Behind one door is a car, behind the others are goats. You pick a door, and the host, who knows what is behind the doors, opens another door (not your pick) which has a goat behind it. Then he asks you if you want to change your original pick. The question we ask you is, "Is it to your advantage to switch your choice?"

Probability III

CLEMSON  
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.7

Notes

---

---

---

---

---

---

---

The Monty Hall Problem



Probability III

CLEMSON  
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.8

Notes

---

---

---

---

---

---

---

The Monty Hall Problem Solution

Probability III

CLEMSON  
UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.9

Notes

---

---

---

---

---

---

---

Bayes' Rule

General form

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

Let  $A_1, A_2, \dots, A_k$  form a partition of the sample space. Then for every event  $B$  in the sample space,

$$\mathbb{P}(A_j|B) = \frac{\mathbb{P}(B|A_j) \times \mathbb{P}(A_j)}{\sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}, \quad j = 1, 2, \dots, k$$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.10

Notes

---

---

---

---

---

---

---

Example

Let us assume that a specific disease is only present in 5 out of every 1,000 people. Suppose that the test for the disease is accurate 99% of the time a person has the disease and 95% of the time that a person lacks the disease. What is the probability that the person has the disease given that they tested positive?

Solution.

$$\mathbb{P}(D|+) = \frac{\mathbb{P}(D \cap +)}{\mathbb{P}(+)} = \frac{.005 \times .99}{.005 \times .99 + .995 \times .05} = \frac{.00495}{.0547} = .0905$$

The reason we get such a surprising result is because the disease is so rare that the number of false positives greatly outnumbers the people who truly have the disease.

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.11

Notes

---

---

---

---

---

---

---

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$  for any event  $A$ ,  $\mathbb{P}(\emptyset) = 0$ ,  $\mathbb{P}(\Omega) = 1$
- Complement rule:  $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule:  
 $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:  
 $\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$
- Conditional probability:  $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- Law of total probability:  
 $\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(B \cap A_i) = \sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)$
- Independence: if  $A$  and  $B$  are independent, then  
 $\mathbb{P}(A|B) = \mathbb{P}(A)$ ,  $\mathbb{P}(B|A) = \mathbb{P}(B)$ , and  
 $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.12

Notes

---

---

---

---

---

---


---

Random Variables

A **random variable** is a real-valued function whose domain is the sample space of a random experiment. In other words, a random variable is a function

$$X : \Omega \mapsto \mathbb{R}$$

where  $\Omega$  is the sample space of the random experiment under consideration and  $\mathbb{R}$  represents the set of all real numbers.



Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.13

Notes

---

---

---

---

---

---


---

Discrete and Continuous Random Variables

There are two main types of quantitative random variables (r.v.s): **discrete** and **continuous**. A discrete r.v. often involves a count of something.

Discrete random variable

A random variable  $X$  is called a discrete random variable if the outcome of the random variable is limited to a countable set of real numbers (usually integers).



Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.14

Notes

---

---

---

---

---

---

---

Example

The following is a chart describing the number of siblings each student in a particular class has.

Siblings	Frequency	Relative Frequency
0	8	.200
1	17	.425
2	11	.275
3	3	.075
4	1	.025
Total	40	1

Let's define the event  $A$  as the event that a randomly chosen student has 2 or more siblings. What is  $\mathbb{P}(A)$ ?

**Solution.**

$$\begin{aligned} \mathbb{P}(A) &= \mathbb{P}(X \geq 2) = \mathbb{P}(X = 2) + \mathbb{P}(X = 3) + \mathbb{P}(X = 4) \\ &= .275 + .075 + .025 = .375 \end{aligned}$$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.15

Notes

---

---

---

---

---

---

---

Probability Mass Function

Let  $X$  be a discrete random variable. Then the probability mass function (pmf) of  $X$  is the real-valued function defined on  $\mathbb{R}$  by

$$p_X(x) = \mathbb{P}(X = x)$$

The capital letter,  $X$ , is used to denote random variable. Lowercase letter,  $x$ , is used to denote possible values of the random variable.

$p_X(x)$ : The probability that the discrete random variable  $X$  is exactly equal to  $x$ .

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.16

Notes

---

---

---

---

---

---

---

Probability Mass Function Example

Flip a fair coin 3 times. Let  $X$  denote the number of heads tossed in the 3 flips. Create a pmf for  $X$

Solution.

The random variable  $X$  maps any outcome to an integer (e.g.  $X((T, T, T)) = 0, X((H, H, T)) = 2$ )

$x$	0	1	2	3
$p_X(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.17

Notes

---

---

---

---

---

---

---

Properties of a PMF

- $0 \leq p_X(x) \leq 1, x \in \{0, 1, 2, \dots\}$
- $\sum_x p_X(x) = 1$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.18

Notes

---

---

---

---

---

---

---

Example

Let  $X$  be a random variable with pmf defined as follows:

$$p_X(x) = \begin{cases} k(5-x) & \text{if } x = 0, 1, 2, 3, 4 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find the value of  $k$  that makes  $p_X(x)$  a legitimate pmf.
- 2 What is the probability that  $X$  is between 1 and 3 inclusive?
- 3 If  $X$  is not 0, what is the probability that  $X$  is less than 3?

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.19

Notes

---

---

---

---

---

---

---

Mean of Discrete Random Variables

The mean of a discrete r.v.  $X$ , denoted by  $\mathbb{E}[X]$ , is defined by

$$\mathbb{E}[X] = \sum_x x \times p_X(x)$$

**Remark:**  
The mean of a discrete r.v. is a weighted average of its possible values, and the weight used is its probability. Sometimes we refer to the expected value as the **expectation (expected value)**, or the **first moment**.

For any function, say  $g(X)$ , we can also find an expectation of that function. It is

$$\mathbb{E}[g(X)] = \sum_x g(x) \times p_X(x)$$

Example

$$\mathbb{E}[X^2] = \sum_x x^2 \times p_X(x)$$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.20

Notes

---

---

---

---

---

---

---

Properties of Mean

Let  $X$  and  $Y$  be discrete r.v.s defined on the same sample space and having finite expectation (i.e.  $\mathbb{E}[X], \mathbb{E}[Y] < \infty$ ). Let  $a$  and  $b$  be constants. Then the following hold:

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[aX + b] = a \times \mathbb{E}[X] + b$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.21

Notes

---

---

---

---

---

---

---

Number of Siblings Example Revisited

Siblings (X)	Frequency	Relative Frequency
0	8	.200
1	17	.425
2	11	.275
3	3	.075
4	1	.025
Total	40	1

Find the expected value of the number of siblings

Solution.

$$\mathbb{E}[X] = \sum_x xp_X(x) = 0 \times .200 + 1 \times .425 + 2 \times .275 + 3 \times .075 + 4 \times .025 = 1.3$$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.22

Notes

---

---

---

---

---

---

---

---

Variance/Standard Deviation of Discrete r.v.'s

The **variance** of a (discrete) r.v., denoted by  $Var(X)$ , is a measure of the spread, or variability, in the r.v.  $Var(X)$  is defined by

$$Var(X) = \mathbb{E}[(X - \mathbb{E}[x])^2]$$

or

$$Var(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$$

The **standard deviation**, denoted by  $sd(X)$ , is the square root of its variance

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.23

Notes

---

---

---

---

---

---

---

---

Properties of Variance

Let  $c$  be a constant. Then the following hold:

- $Var(cX) = c^2 \times Var(X)$
- $Var(X + c) = Var(X)$

Probability III

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

7.24

Notes

---

---

---

---

---

---

---

---



Example

Suppose  $X$  and  $Y$  are random variables with  $\mathbb{E}[X] = 3$ ,  $\mathbb{E}[Y] = 4$  and  $\text{Var}(X) = 4$ . Find:

- 1  $\mathbb{E}[2X + 1]$
- 2  $\mathbb{E}[X - Y]$
- 3  $\mathbb{E}[X^2]$
- 4  $\mathbb{E}[X^2 - 4]$
- 5  $\mathbb{E}[(X - 4)^2]$
- 6  $\text{Var}(2X - 4)$

Probability III

CLEMSON  
UNIVERSITY

Law of Total  
Probability

Bayes' Rule

Random Variables

7.25

Notes

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---