

Lecture 15


Interpolation of Spatial Data

DSA 8020 Statistical Methods II
April 19-23, 2021

Whitney Huang
Clemson University

Notes


Toy Examples of Spatial Interpolation



Question: What is your best guess of the value of the missing pixel, denoted as $Y(s_0)$, for each case?

Notes

Interpolating Paraná State Precipitation Data



Goal: To interpolate the values in the spatial domain

Notes

The Spatial Interpolation Problem

Given observations of a spatially varying quantity Y at n spatial locations

$y(s_1), y(s_2), \dots, y(s_n), \quad s_i \in \mathcal{S}, i = 1, \dots, n$

We want to estimate this quantity at any **unobserved location**

$Y(s_0), \quad s_0 \in \mathcal{S}$

Applications

- Mining: ore grade
- Climate: temperature, precipitation, ...
- Remote Sensing: CO₂ retrievals
- Environmental Science: air pollution levels, ...

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.4

Notes

Some History of Spatial Statistics

- Mining (Kriging 1951) Matheron (1960s), Forestry (Matérn 1960)
- More recent work: Cressie (1993) Stein (1999)



Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.5

Notes

Outline

- 1 Gaussian Process Spatial Model
- 2 Spatial Interpolation
- 3 Parameter estimation

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.6

Notes

Linear Interpolation

The best guess (in a statistical sense) should be based on the conditional distribution $[Y(s_0) | Y = y]$ where

$$y = (y(s_1), \dots, y(s_n))^T$$

- Calculating this conditional distribution can be difficult
- Instead we use a **linear predictor**:

$$\hat{Y}(s_0) = \lambda_0 + \sum_{i=1}^n \lambda_i y(s_i)$$

- The best linear predictor is completely determined by the **mean** and **covariance** of $\{Y(s), s \in S\}$, and the observations y



Notes

Gaussian Process (GP) Spatial Model

We assume that the observed data $\{y(s_i)\}_{i=1}^n$ is one partial realization of a (continuously indexed) spatial GP $\{Y(s)\}_{s \in S}$.

Model:

$$Y(s) = m(s) + \epsilon(s), \quad s \in S \subset \mathbb{R}^d$$


where

- **Mean function:**

$$m(s) = E[Y(s)] = X^T(s)\beta$$

- **Covariance function:**

$$\{\epsilon(s)\}_{s \in S} \sim \text{GP}(0, K(\cdot, \cdot)), \quad K(s_1, s_2) = \text{Cov}(\epsilon(s_1), \epsilon(s_2))$$



Notes

Assumptions on Covariance Function


In practice, the covariance must be estimated from the data $(y(s_1), \dots, y(s_n))^T$. We need to impose some structural assumptions

- **Stationarity:**

$$\begin{aligned} K(s_1, s_2) &= \text{Cov}(\epsilon(s_1), \epsilon(s_2)) = C(s_1 - s_2) \\ &= \text{Cov}(\epsilon(s_1 + h), \epsilon(s_2 + h)) \end{aligned}$$

- **Isotropy:**

$$K(s_1, s_2) = \text{Cov}(\epsilon(s_1), \epsilon(s_2)) = C(\|s_1 - s_2\|)$$



Notes

A Valid Covariance Function Must Be Positive Definite (p.d.)!

A covariance function is positive definite if

$$\sum_{i,j=1}^n a_i a_j C(\mathbf{s}_i - \mathbf{s}_j) \geq 0$$

for any finite locations $\mathbf{s}_1, \dots, \mathbf{s}_n$, and for any constants $a_i, i = 1, \dots, n$

Question: what is the consequence if a covariance function is NOT p.d.? \Rightarrow **We can get a negative variance**

Question: How to guarantee a $C(\cdot)$ is p.d.?

- Using a **parametric covariance function** (see some examples in next slide)
- Using **Bochner's Theorem** to construct a valid covariance function

Notes

Some Commonly Used Covariance Functions

- Powered exponential:**

$$C(h) = \sigma^2 \exp\left(-\left(\frac{h}{\rho}\right)^\alpha\right), \quad \sigma^2 > 0, \rho > 0, 0 < \alpha \leq 2$$

- Spherical:**

$$C(h) = \sigma^2 \left(1 - 1.5 \frac{h}{\rho} + 0.5 \left(\frac{h}{\rho}\right)^3\right) 1_{\{h \leq \rho\}}, \quad \sigma^2, \rho > 0$$

Note: it is only valid for 1, 2, and 3 dimensional spatial domain.

- Matérn:**

$$C(h) = \sigma^2 \frac{(\sqrt{2\nu}h/\rho)^\nu K_\nu(\sqrt{2\nu}h/\rho)}{\Gamma(\nu)2^{\nu-1}}, \quad \sigma^2 > 0, \rho > 0, \nu > 0$$

"Use the Matérn model" – Stein (1999, pp. 14)

Notes

1-D Realizations from Matérn Model with Fixed σ^2, ρ





Figure: courtesy of Rasmussen & Williams 2006

The larger ν is, the smoother the process is

Notes

2-D Realizations from Matérn Model with Fixed σ^2



Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.13

Notes

Outline

- 1 Gaussian Process Spatial Model
- 2 Spatial Interpolation
- 3 Parameter estimation

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.14

Notes

Conditional Distribution of Multivariate Normal

If
$$\begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix} \sim N \left(\begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix} \right)$$

Then
$$[Y_1|Y_2 = y_2] \sim N(\mu_{1|2}, \Sigma_{1|2})$$

where
$$\begin{aligned} \mu_{1|2} &= \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(y_2 - \mu_2) \\ \Sigma_{1|2} &= \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} \end{aligned}$$

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.15

Notes

GP-Based Spatial Interpolation: Kriging

If $\{Y(s)\}_{s \in \mathcal{S}}$ follows a GP, then

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim N \left(\begin{pmatrix} m_0 \\ \mathbf{m} \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & k^T \\ k & \Sigma \end{pmatrix} \right)$$

We have

$$[Y_0 | \mathbf{Y} = \mathbf{y}] \sim N \left(m_{Y_0 | \mathbf{Y} = \mathbf{y}}, \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 \right)$$

where

$$\begin{aligned} m_{Y_0 | \mathbf{Y} = \mathbf{y}} &= m_0 + k^T \Sigma^{-1} (\mathbf{y} - \mathbf{m}) \\ \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 &= \sigma_0^2 - k^T \Sigma^{-1} k \end{aligned}$$

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model


Spatial Interpolation

Parameter estimation

15.16

Notes

Spatial Prediction of Paraná State Rainfall



Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.17

Notes

GP-Based Spatial Interpolation: Kriging

If $\{Y(s)\}_{s \in \mathcal{S}}$ follows a GP, then

$$\begin{pmatrix} Y_0 \\ \mathbf{Y} \end{pmatrix} \sim N \left(\begin{pmatrix} m_0 \\ \mathbf{m} \end{pmatrix}, \begin{pmatrix} \sigma_0^2 & k^T \\ k & \Sigma \end{pmatrix} \right)$$

We have

$$[Y_0 | \mathbf{Y} = \mathbf{y}] \sim N \left(m_{Y_0 | \mathbf{Y} = \mathbf{y}}, \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 \right)$$

where

$$\begin{aligned} m_{Y_0 | \mathbf{Y} = \mathbf{y}} &= m_0 + k^T \Sigma^{-1} (\mathbf{y} - \mathbf{m}) \\ \sigma_{Y_0 | \mathbf{Y} = \mathbf{y}}^2 &= \sigma_0^2 - k^T \Sigma^{-1} k \end{aligned}$$

Question: what if we don't know $m_0, \mathbf{m}, \sigma_0^2, \Sigma$?

⇒ We need to estimate the mean and covariance from the data \mathbf{y} .

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.18

Notes

Outline

- 1 Gaussian Process Spatial Model
- 2 Spatial Interpolation
- 3 Parameter estimation

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation


15.19

Notes

Recap: Gaussian Process

We assume that the observed data $\{y(s_i)\}_{i=1}^n$ is one partial realization of a (continuously indexed) spatial stochastic process $\{Y(s)\}_{s \in \mathcal{S}}$.

- Gaussian Processes $GP(m(\cdot), K(\cdot, \cdot))$ are widely used in modeling spatial stochastic processes
- Spatial statisticians often focus on the covariance function. e.g. $K(h) = \sigma^2 \frac{(\sqrt{2\nu}h/\gamma)^\nu \mathcal{K}_\nu(\sqrt{2\nu}h/\gamma)}{\Gamma(\nu)2^{\nu-1}}$



Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.20

Notes

Variogram, Semivariogram, and Covariance Function

Under the stationary and isotropic assumptions

Variogram:

$$\begin{aligned} 2\gamma(s_i, s_j) &= \text{Var}(Y(s_i) - Y(s_j)) \\ &= E\left\{((Y(s_i) - \mu(s_i)) - (Y(s_j) - \mu(s_j)))^2\right\} \\ &= E\left\{(Y(s_i) - Y(s_j))^2\right\} \\ &= 2\gamma(\|s_i - s_j\|) = 2\gamma(h) \end{aligned}$$

Semivariogram and covariance function:

$$\gamma(h) = C(0) - C(h)$$

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model


Spatial Interpolation

Parameter estimation

15.21

Notes

Semivariogram $\{\frac{1}{2}\text{Var}(\varepsilon(s_i) - \varepsilon(s_j))\}_{i,j}$




Source: fields vignette by Wiens and Krock, 2019

Notes

Estimation: Weighted Least Squares Method

$$\underset{\theta}{\operatorname{argmin}} \sum_{u \in \mathcal{U}} \frac{N(h_u)}{[\gamma(h_u; \theta)]^2} [\hat{\gamma}(h_u) - \gamma(h_u; \theta)]^2$$



Notes

Maximum Likelihood Estimation (MLE)

Log-likelihood:

Given data $\mathbf{y} = (y(s_1), \dots, y(s_n))^T$

$$\ell_n(\beta; \mathbf{y}) \propto -\frac{1}{2} \log |\Sigma_\theta| - \frac{1}{2} (\mathbf{y} - \mathbf{X}^T \beta)^T [\Sigma_\theta]_{n \times n}^{-1} (\mathbf{y} - \mathbf{X}^T \beta)$$

where $\Sigma_\theta(i, j) = \sigma^2 \rho_{\rho, \nu}(\|s_i - s_j\|) + \tau^2 1_{\{s_i = s_j\}}, i, j = 1, \dots, n$

for any fixed $\theta_0 \in \Theta$ the unique value of β that maximizes ℓ_n is given by

$$\hat{\beta} = (\mathbf{X}^T \Sigma_{\theta_0}^{-1} \mathbf{X})^{-1} \mathbf{X}^T \Sigma_{\theta_0}^{-1} \mathbf{y}$$

Then we obtain the profile log likelihood

$$\ell_n(\theta; \mathbf{y}) \propto -\frac{1}{2} \log |\Sigma_\theta| - \frac{1}{2} \mathbf{y}^T P(\theta) \mathbf{y}$$

where

$$P(\theta) = \Sigma_\theta^{-1} - \Sigma_\theta^{-1} \mathbf{X} (\mathbf{X}^T \Sigma_\theta^{-1} \mathbf{X})^{-1} \mathbf{X}^T \Sigma_\theta^{-1}$$

Solve the maximization problem above to get the MLE

Notes

Bochner's Theorem

A complex-valued function C on \mathbb{R}^d is the covariance function for a weakly stationary mean square continuous complex-valued random process on \mathbb{R}^d if and only if it can be represented as

$$C(\mathbf{h}) = \int_{\mathbb{R}^d} \exp(i\omega^T \mathbf{h}) F(d\omega),$$

with F a positive finite measure. When F has a density with respect to Lebesgue measure, we have the spectral density f and

$$f(\omega) = \frac{1}{2\pi} \int_{\mathbb{R}^d} \exp(-i\omega^T \mathbf{h}) C(\mathbf{h}) d\mathbf{h}$$

Interpolation of Spatial Data

CLEMSON UNIVERSITY

Gaussian Process Spatial Model

Spatial Interpolation

Parameter estimation

15.25

Notes

Notes

Notes
