

Lecture 8

Introduction to Probability II

Text: Chapter 4

STAT 8010 Statistical Methods I
September 6, 2019

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Whitney Huang
Clemson University

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

- 1 Union and Intersection
- 2 Logical Relationships among Events
- 3 Complement Rule and General Addition Rule
- 4 Venn Diagram

- **Intersection:** the intersection of two events A and B , denoted by $A \cap B$, is the event that contains all outcomes of A that also belong to $B \Rightarrow$ **AND**

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

- **Intersection:** the intersection of two events A and B , denoted by $A \cap B$, is the event that contains all outcomes of A that also belong to $B \Rightarrow$ **AND**

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 4, 5\}$, then
 $A \cap B = \{1, 2\}$

- **Intersection:** the intersection of two events A and B , denoted by $A \cap B$, is the event that contains all outcomes of A that also belong to $B \Rightarrow$ **AND**

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 4, 5\}$, then
 $A \cap B = \{1, 2\}$

- **Union:** the union of two events A and B , denoted by $A \cup B$, is the event of all outcomes that belong to either A or $B \Rightarrow$ **OR**

- **Intersection:** the intersection of two events A and B , denoted by $A \cap B$, is the event that contains all outcomes of A that also belong to $B \Rightarrow$ **AND**

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 4, 5\}$, then
 $A \cap B = \{1, 2\}$

- **Union:** the union of two events A and B , denoted by $A \cup B$, is the event of all outcomes that belong to either A or $B \Rightarrow$ **OR**

Example: Let $A = \{1, 2, 3\}$ and $B = \{1, 2, 4, 5\}$, then
 $A \cup B = \{1, 2, 3, 4, 5\}$

Example

Suppose we flipped 3 fair coins. Let A be the event of **exactly 2 tails**. Let B be the event that the **first 2 tosses are tails**. Let C be the event that **all 3 tosses are tails**. What are $A \cap B$, $A \cup C$, and $(A \cap B) \cup C$?

Solution.

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Example

Suppose we flipped 3 fair coins. Let A be the event of **exactly 2 tails**. Let B be the event that the **first 2 tosses are tails**. Let C be the event that **all 3 tosses are tails**. What are $A \cap B$, $A \cup C$, and $(A \cap B) \cup C$?

Solution.

$$A = \{(T, T, H), (T, H, T), (H, T, T)\}$$

$$B = \{(T, T, T), (T, T, H)\}$$

$$C = \{T, T, T\}$$

$$\textcircled{1} \quad A \cap B = \{(T, T, H)\}$$

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Example

Suppose we flipped 3 fair coins. Let A be the event of **exactly 2 tails**. Let B be the event that the **first 2 tosses are tails**. Let C be the event that **all 3 tosses are tails**. What are $A \cap B$, $A \cup C$, and $(A \cap B) \cup C$?

Solution.

$$A = \{(T, T, H), (T, H, T), (H, T, T)\}$$

$$B = \{(T, T, T), (T, T, H)\}$$

$$C = \{T, T, T\}$$

$$1 \quad A \cap B = \{(T, T, H)\}$$

$$2 \quad A \cup C = \{(T, T, H), (T, H, T), (H, T, T), (T, T, T)\}$$

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Example

Suppose we flipped 3 fair coins. Let A be the event of **exactly 2 tails**. Let B be the event that the **first 2 tosses are tails**. Let C be the event that **all 3 tosses are tails**. What are $A \cap B$, $A \cup C$, and $(A \cap B) \cup C$?

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Solution.

$$A = \{(T, T, H), (T, H, T), (H, T, T)\}$$

$$B = \{(T, T, T), (T, T, H)\}$$

$$C = \{T, T, T\}$$

1 $A \cap B = \{(T, T, H)\}$

2 $A \cup C = \{(T, T, H), (T, H, T), (H, T, T), (T, T, T)\}$

3 $(A \cap B) \cup C = \{(T, T, H)\} \cup \{(T, T, T)\} = \{(T, T, H), (T, T, T)\}$

Logical Relationships among Events

- **Mutually exclusive:** refers to two (or more) events that cannot both occur when the random experiment is formed.

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Logical Relationships among Events

- **Mutually exclusive:** refers to two (or more) events that cannot both occur when the random experiment is formed.

$$A \cap B = \emptyset$$

- **Exhaustive:** refers to event(s) that comprise the sample space.

Logical Relationships among Events

- **Mutually exclusive:** refers to two (or more) events that cannot both occur when the random experiment is formed.

$$A \cap B = \emptyset$$

- **Exhaustive:** refers to event(s) that comprise the sample space.

$$A \cup B = \Omega$$

- **Partition:** events that are both mutually exclusive and exhaustive.

Logical Relationships among Events

- **Mutually exclusive:** refers to two (or more) events that cannot both occur when the random experiment is formed.

$$A \cap B = \emptyset$$

- **Exhaustive:** refers to event(s) that comprise the sample space.

$$A \cup B = \Omega$$

- **Partition:** events that are both mutually exclusive and exhaustive.

$$A \cap B = \emptyset \quad \text{and} \quad A \cup B = \Omega$$


Complement

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram



Complement Rule

- 1 By the definition of complement

$$A \cup A^c = \Omega$$

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Complement Rule

- 1 By the definition of complement

$$A \cup A^c = \Omega$$

- 2 Apply the probability operator

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(\Omega) = 1$$

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Complement Rule

- 1 By the definition of complement

$$A \cup A^c = \Omega$$

- 2 Apply the probability operator

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(\Omega) = 1$$

- 3 Since A and A^c are mutually exclusive

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(A) + \mathbb{P}(A^c)$$

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Complement Rule

- 1 By the definition of complement

$$A \cup A^c = \Omega$$

- 2 Apply the probability operator

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(\Omega) = 1$$

- 3 Since A and A^c are mutually exclusive

$$\mathbb{P}(A \cup A^c) = \mathbb{P}(A) + \mathbb{P}(A^c)$$

- 4 Hence we get $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$

Example

Suppose we rolled a fair, six-sided die 10 times. Let T be the event that we roll at least 1 three. If one were to calculate T you would need to find the probability of 1 three, 2 threes, \dots , and 10 threes and add them all up. However, you can use the complement rule to calculate $\mathbb{P}(T)$

Solution.

Let X be the times that we rolled a 3, then

$$\mathbb{P}(T) = \mathbb{P}(X \geq 1) = \underbrace{\mathbb{P}(X = 1) + \mathbb{P}(X = 2) + \dots + \mathbb{P}(X = 10)}_{\text{need to compute 10 probabilities}}$$

Example

Suppose we rolled a fair, six-sided die 10 times. Let T be the event that we roll at least 1 three. If one were to calculate T you would need to find the probability of 1 three, 2 threes, \dots , and 10 threes and add them all up. However, you can use the complement rule to calculate $\mathbb{P}(T)$

Solution.

Let X be the times that we rolled a 3, then

$$\mathbb{P}(T) = \mathbb{P}(X \geq 1) = \underbrace{\mathbb{P}(X = 1) + \mathbb{P}(X = 2) + \dots + \mathbb{P}(X = 10)}_{\text{need to compute 10 probabilities}}$$

If we apply the complement rule

$$\mathbb{P}(T) = 1 - \mathbb{P}(T^c) = 1 - \mathbb{P}(X = 0)$$

Venn Diagram


A Venn diagram is a diagram that shows all possible logical relations between a finite collection of events.

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule


Venn Diagram



General Addition Rule

The general addition rule is a way of finding the probability of a union of 2 events. It is $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

VENN DIAGRAM!



Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram

Example

Three of the major commercial computer operating systems are Windows, Mac OS, and Red Hat Linux Enterprise. A Computer Science professor selects 50 of her students and asks which of these three operating systems they use. The results for the 50 students are summarized below.

- 30 students use Windows
- 16 students use at least two of the operating systems
- 9 students use all three operating systems
- 18 students use Mac OS
- 46 students use at least one of the operating systems
- 11 students use both Windows and Linux


Example cont'd

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram



Conditional Probability

Let A and B be events. The probability that event A occurs **given** (knowing) that event B occurs is called a **conditional probability** and is denoted by $P(A|B)$. The formula of conditional probability is


$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Union and Intersection

Logical Relationships
among Events

Complement Rule and
General Addition Rule

Venn Diagram



In a conditional probability problem, the sample space is "reduced" to the "space" of the given outcome (e.g. if given B , we now just care about the probability of A occurring "inside" of B)

Summary

In this lecture, we learned

- Union , Intersection
- **Logical Relationships among Events** Mutually Exclusive, Exhaustive, Partition
- Complement Rule and General Addition Rule
- Venn Diagram