

# Lecture 13

## Inference on Two Population Means

Text: Chapter 6

STAT 8010 Statistical Methods I

February 27, 2020

Whitney Huang  
Clemson University



### Notes

---

---

---

---

---

---

---

### Agenda

1 Inference on Two Population Means: Two-Sample t Confidence Intervals/Tests

2 Paired T-Test



### Notes

---

---

---

---

---

---

---

### Comparing Two Population Means

- We often interested in comparing two groups (e.g.)
  - Does a particular pesticide increase the yield of corn per acre?
  - Do men and women in the same occupation have different salaries?
- The common ingredient in these questions: They can be answered by conducting statistical inferences of two populations using two (independent) samples, one from each of two populations



### Notes

---

---

---

---

---

---

---

## Notation

- Parameters:
  - Population means:  $\mu_1, \mu_2$
  - Population standard deviations:  $\sigma_1, \sigma_2$
- Statistics:
  - Sample means:  $\bar{X}_1, \bar{X}_2$
  - Sample standard deviations:  $s_1, s_2$
  - Sample sizes:  $n_1, n_2$



## Notes

---

---

---

---

---

---

---

## Statistical Inference for $\mu_1 - \mu_2$

- Point estimate:  $\bar{X}_1 - \bar{X}_2$
- Interval estimate: Need to figure out  $\sigma_{\bar{X}_1 - \bar{X}_2}$
- Hypothesis Testing:
  - Upper-tailed test:  $H_0 : \mu_1 - \mu_2 = 0$  vs.  $H_a : \mu_1 - \mu_2 > 0$
  - Lower-tailed test:  $H_0 : \mu_1 - \mu_2 = 0$  vs.  $H_a : \mu_1 - \mu_2 < 0$
  - Two-tailed test:  $H_0 : \mu_1 - \mu_2 = 0$  vs.  $H_a : \mu_1 - \mu_2 \neq 0$



## Notes

---

---

---

---

---

---

---

## Confidence Intervals for $\mu_1 - \mu_2$

If we are willing to **assume**  $\sigma_1 = \sigma_2$ , then we can “pool” these two (independent) samples together to estimate the common  $\sigma$  using  $s_p$ :

$$s_p = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}}$$

We can then derive the (estimated) standard error of  $\bar{X}_1 - \bar{X}_2$ , which takes the following form

$$\sqrt{\frac{s_p^2}{n_1} + \frac{s_p^2}{n_2}} = s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

With CLT (assuming sample sizes are sufficiently large), we obtain the  $(1 - \alpha) \times 100\%$  CI for  $\mu_1 - \mu_2$ :

$$\underbrace{\bar{X}_1 - \bar{X}_2}_{\text{point estimate}} \pm \underbrace{t_{\alpha/2, n_1 + n_2 - 1} s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}_{\text{margin of error}}$$



## Notes

---

---

---

---

---

---

---

Confidence Intervals for  $\mu_1 - \mu_2$ : What if  $\sigma_1 \neq \sigma_2$ ?

- We will use  $s_1^2, s_2^2$  as the estimates for  $\sigma_1^2$  and  $\sigma_2^2$  to obtain the standard error:

$$\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

- The formula for the degrees of freedom is somewhat complicated:

$$\frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{(s_1^2/n_1)^2}{n_1-1} + \frac{(s_2^2/n_2)^2}{n_2-1}}$$

- We can then construct the  $(1 - \alpha) \times 100\%$  CI for  $\mu_1 - \mu_2$ :

$$\underbrace{\bar{X}_1 - \bar{X}_2}_{\text{point estimate}} \pm \underbrace{t(\alpha/2, \text{df calculated from above}) \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}_{\text{margin of error}}$$

Notes

---

---

---

---

---

---

---

To Pool ( $\sigma_1 = \sigma_2$ ) or Not to Pool ( $\sigma_1 \neq \sigma_2$ )?

We could perform the following test:

- $H_0 : \sigma_1^2/\sigma_2^2 = 1$  vs.  $\sigma_1^2/\sigma_2^2 \neq 1$
- Test statistic:  $F^* = s_1^2/s_2^2$ . Under  $H_0$ ,  $F^* \sim F(n_1 - 1, n_2 - 1)$
- For a given  $\alpha$ , we reject  $H_0$  if the P-value  $< \alpha$  (or  $F_{obs} > F_{\alpha}(n_1, n_2)$ )
- If we fail to reject  $H_0$ , then we will use  $s_p$  as an estimate for  $\sigma$  and we have  $s_{\bar{X}_1 - \bar{X}_2} = s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$ . Otherwise, we use  $s_{\bar{X}_1 - \bar{X}_2} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$

Notes

---

---

---

---

---

---

---

Example

An experiment was conducted to evaluate the effectiveness of a treatment for tapeworm in the stomachs of sheep. A random sample of 24 worm-infected lambs of approximately the same age and health was randomly divided into two groups. Twelve of the lambs were injected with the drug (treatment group) and the remaining twelve were left untreated (control group). After a 6-month period, the worm counts were recorded:

Treatment	18	43	28	50	16	32	13	35	38	33	6	7
Control	40	54	26	63	21	37	39	23	48	58	28	39

Notes

---

---

---


---

---

---

---

## Plot the Two Samples



- $n_1 = n_2 = 12 \Rightarrow$  sample size is perhaps not large enough for CLT to work. But the boxplots suggest the distributions are symmetric with no outliers
- The untreated lambs (control group) appear to have higher average worm counts than the treated lambs (treatment group). But do we have enough evidence?

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests

Paired T-Test

13.10

## Notes

---

---

---

---

---

---

---

---

## Example Cont'd

```
> apply(dat, 2, mean)
treatment control
26.58333 39.66667
> apply(dat, 2, sd)
treatment control
14.36193 13.85859
> var.test(treatment, control)
```

F test to compare two variances

```
data: treatment and control
F = 1.074, num df = 11, denom df = 11, p-value = 0.9079
alternative hypothesis: true ratio of variances is not equal to 1
95 percent confidence interval:
 0.3091686 3.7306092
sample estimates:
ratio of variances
 1.073959
```

We fail to reject  $\sigma_1 = \sigma_2 = \sigma$ . Therefore we will use  $s_p$ , the pooled standard deviation, as an estimate for  $\sigma$

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests

Paired T-Test

13.11

## Notes

---

---

---

---

---

---

---

---

## Example Cont'd

- Place a 95% confidence interval on  $\mu_1 - \mu_2$  to assess the size of the difference in the two population means
- Test whether the mean number of tapeworms in the stomachs of the treated lambs is less than the mean for untreated lambs. Use an  $\alpha = 0.05$  test

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests

Paired T-Test

13.12

## Notes

---

---

---

---

---

---

---

---

Another Example

A simple random sample with sample size 37 is taken and are subjected to a treatment ( $\bar{X}_1 = 19.45, s_1 = 4.3$ ). A Simple random sample with sample size 31 is taken and given a placebo ( $\bar{X}_2 = 18.2, s_2 = 2.2$ ). At the 10% level can we say that the means are different between the two groups?

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.13

Notes

---

---

---

---

---

---

---

Paired T-Test: Motivating Example

Insurance handlers are concerned about the high estimates they are receiving for auto repairs from garage I compared to garage II. To verify their suspicions, each of 15 cars recently involved in an accident was taken to both garages for separate estimates of repair costs. The estimates from the two garages are given in the following table

Garage I	Garage II	Garage I	Garage II	Garage I	Garage II
17.6	17.3	20.2	19.1	19.5	18.4
11.3	11.5	13.0	12.7	16.3	15.8
15.3	14.9	16.2	15.3	12.2	12.0
14.8	14.2	21.3	21.0	22.1	21.0
16.9	16.1	17.6	16.7	18.4	17.5

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.14

Notes

---

---

---

---

---

---

---

Example Cont'd

Suppose we perform a two-sample test

Sample statistics:  
 $\bar{X}_1 = 16.85, \bar{X}_2 = 16.23, s_1 = 3.20, s_2 = 2.94$

- $H_0 : \mu_1 - \mu_2 = 0$  vs.  $H_a : \mu_1 - \mu_2 > 0$
- $t_{obs} = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}} = \frac{16.85 - 16.23}{\sqrt{\frac{3.2^2}{15} + \frac{2.94^2}{15}}} = \frac{0.62}{1.12} = 0.55$
- Critical value for rejection region:  $t_{0.05, df=27} = 1.70$
- Since  $t_{obs}$  is not in the rejection region. We fail to reject  $H_0$  at 0.05 level.

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.15

Notes

---

---

---


---

---

---

---

Boxplots and R Output



```
Welch Two Sample t-test

data: GarageI and GarageII
t = 0.54616, df = 27.797, p-value = 0.2947
alternative hypothesis: true difference in means is greater than 0
95 percent confidence interval:
-1.29749      Inf
sample estimates:
mean of x mean of y
16.84667 16.23333
```

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.16

Notes

---

---


---

---

---

---

---



For all but one of the 15 cars, the estimates from garage I were higher than that from garage II.

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.17

Notes

---

---

---

---

---

---

---

Analyzing Matched Pairs

- Matched pairs are **dependent samples** where each unit in the first sample is directly linked with a unit in the second sample
- This could occur in several situations, for example, before/after study, study on twins, pairing subjects based on similar characteristics
- We need different strategy for testing two dependent samples ⇒ **Paired T-Tests**

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.18

Notes

---

---

---

---

---

---

---

Paired T-Tests

- $H_0 : \mu_{diff} = 0$  vs.  $H_a : \mu_{diff} > 0$  (Upper-tailed);  $\mu_{diff} < 0$  (Lower-tailed);  $\mu_{diff} \neq 0$  (Two-tailed)
- Test statistic:  $t^* = \frac{\bar{x}_{diff} - 0}{\frac{s_{diff}}{\sqrt{n}}}$ . If  $\mu_{diff} = 0$ , then  $t^* \sim t_{df=n-1}$
- Use rejection region method or P-value method to make a decision

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.19

Notes

---

---

---

---

---

---

---

Car Repair Example Revisited

Garage I - Garage II	Garage I - Garage II	Garage I - Garage II
17.6 - 17.3 = 0.3	20.2 - 19.1 = 1.1	19.5 - 18.4 = 1.1
11.3 - 11.5 = -0.2	13.0 - 12.7 = 0.3	16.3 - 15.8 = 0.5
15.3 - 14.9 = 0.4	16.2 - 15.3 = 0.9	12.2 - 12.0 = 0.2
14.8 - 14.2 = 0.6	21.3 - 21.0 = 0.3	22.1 - 21.0 = 1.1
16.9 - 16.1 = 0.8	17.6 - 16.7 = 0.9	18.4 - 17.5 = 0.9

- 1 First, compute the difference in paired samples
- 2 Compute the sample mean and standard deviation for the differences
- 3 Then perform a one sample t-test

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.20

Notes

---

---

---

---

---

---

---

Car Repair Example Cont'd

$\bar{X}_{diff} = 0.61, s_{diff} = 0.39$

- 1  $H_0 : \mu_{diff} = 0$  vs.  $H_a : \mu_{diff} > 0$
- 2  $t_{obs} = \frac{0.61}{\frac{0.39}{\sqrt{15}}} = 6.03$
- 3 Critical value for rejection region:  $t_{0.05, df=14} = 1.76 \Rightarrow$  reject  $H_0$
- 4 We do have enough evidence that the true mean repair cost difference for the garage I and II is greater than 0

Inference on Two Population Means

CLEMSON UNIVERSITY

Inference on Two Population Means:  
Two-Sample t  
Confidence  
Intervals/Tests  
Paired T-Test

13.21

Notes

---

---

---


---

---

---

---

Boxplot and R Output



Paired t-test

data: GarageI and GarageII  
t = 6.0234, df = 14, p-value = 1.563e-05  
alternative hypothesis: true difference in means  
is greater than 0  
95 percent confidence interval:  
0.4339886 Inf  
sample estimates:  
mean of the differences  
0.6133333

Notes

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---