

DSA 8020 R Session 4: Multiple Linear Regression III

Whitney

January 24, 2021

Contents

Model Selection	1
Best Subset Selection	1
Backward Selection	5
Stepwise Selection	6
Model Diagnostics	7
Residual Plot	7
Residual Histogram/QQplot	8
Leverage	10
Studentized Residuals	11
Jackknife Residuals	12
Identifying Influential Observations: DFFITS	13
Transformation	14

```
library(faraway)
data(gala)
galaNew <- gala[, -2]
```

Model Selection

Best Subset Selection

```
#install.packages(c("tidyverse", "caret", "leaps"))
library(tidyverse)
library(caret)
library(leaps)
models <- regsubsets(Species ~ ., data = galaNew)
summary(models)

## Subset selection object
## Call: regsubsets.formula(Species ~ ., data = galaNew)
## 5 Variables (and intercept)
```

```
##          Forced in Forced out
## Area          FALSE      FALSE
## Elevation     FALSE      FALSE
## Nearest       FALSE      FALSE
## Scrutz        FALSE      FALSE
## Adjacent      FALSE      FALSE
## 1 subsets of each size up to 5
## Selection Algorithm: exhaustive
##          Area Elevation Nearest Scrutz Adjacent
## 1 ( 1 ) " " " *"      " "      " "      " "
## 2 ( 1 ) " " " *"      " "      " "      " *"
## 3 ( 1 ) " " " *"      " "      " *"      " *"
## 4 ( 1 ) " *" " *"      " "      " *"      " *"
## 5 ( 1 ) " *" " *"      " *"      " *"      " *
```


```
res.sum <- summary(models)
```

```
criteria <- data.frame(
  Adj.R2 = res.sum$adjr2,
  Cp = res.sum$cp,
  BIC = res.sum$bic)
```


```
criteria
```

```
##      Adj.R2      Cp      BIC
## 1 0.5291255 20.599003 -16.84525
## 2 0.7181425  2.897184 -29.93078
## 3 0.7258462  3.193068 -28.49317
## 4 0.7283816  4.000075 -26.54733
## 5 0.7170651  6.000000 -23.14622
```


```
plot(2:6, criteria$Cp, las = 1, xlab = "p", ylab = "Cp",
     pch = 16, col = "gray", ylim = c(1, max(criteria$Cp)))
abline(0, 1)
```



```
plot(2:6, criteria$Adj.R2, las = 1, xlab = "p", ylab = "", pch = 16, col = "gray",
     main = expression(R['adj']^2))
points(5, criteria$Adj.R2[4], col = "blue", pch = 16)
```



```
plot(2:6, criteria$BIC, las = 1, xlab = "p", ylab = "", pch = 16, col = "gray", main = "BIC")
points(3, criteria$BIC[2], col = "blue", pch = 16)
```



Backward Selection

```
full <- lm(Species ~ ., data = galaNew)
step(full, direction = "backward")
```

```
## Start:  AIC=251.93
## Species ~ Area + Elevation + Nearest + Scruz + Adjacent
##
##           Df Sum of Sq    RSS    AIC
## - Nearest   1         0 89232 249.93
## - Area       1      4238 93469 251.33
## - Scruz      1      4636 93867 251.45
## <none>                89231 251.93
## - Adjacent   1     66406 155638 266.62
## - Elevation  1    131767 220998 277.14
```

```
##
## Step: AIC=249.93
## Species ~ Area + Elevation + Scrutz + Adjacent
##
##           Df Sum of Sq    RSS    AIC
## - Area      1      4436  93667 249.39
## <none>                89232 249.93
## - Scrutz     1      7544  96776 250.37
## - Adjacent   1     72312 161544 265.74
## - Elevation  1    139445 228677 276.17
##
## Step: AIC=249.39
## Species ~ Elevation + Scrutz + Adjacent
##
##           Df Sum of Sq    RSS    AIC
## - Scrutz     1      6336 100003 249.35
## <none>                93667 249.39
## - Adjacent   1     69860 163527 264.11
## - Elevation  1    275784 369451 288.56
##
## Step: AIC=249.35
## Species ~ Elevation + Adjacent
##
##           Df Sum of Sq    RSS    AIC
## <none>                100003 249.35
## - Adjacent   1     73251 173254 263.84
## - Elevation  1    280817 380820 287.47
##
##
## Call:
## lm(formula = Species ~ Elevation + Adjacent, data = galaNew)
##
## Coefficients:
## (Intercept)    Elevation    Adjacent
##      1.43287      0.27657     -0.06889
```

Stepwise Selection

```
step(full, direction = "both")
```


```
## Start: AIC=251.93
## Species ~ Area + Elevation + Nearest + Scrutz + Adjacent
##
##           Df Sum of Sq    RSS    AIC
## - Nearest   1         0  89232 249.93
## - Area       1      4238  93469 251.33
## - Scrutz     1      4636  93867 251.45
## <none>                89231 251.93
## - Adjacent   1     66406 155638 266.62
## - Elevation  1    131767 220998 277.14
##
## Step: AIC=249.93
```

```
## Species ~ Area + Elevation + Scruz + Adjacent
##
##           Df Sum of Sq   RSS   AIC
## - Area      1      4436  93667 249.39
## <none>                89232 249.93
## - Scruz      1      7544  96776 250.37
## + Nearest    1         0  89231 251.93
## - Adjacent   1     72312 161544 265.74
## - Elevation  1    139445 228677 276.17
##
## Step:  AIC=249.39
## Species ~ Elevation + Scruz + Adjacent
##
##           Df Sum of Sq   RSS   AIC
## - Scruz      1      6336 100003 249.35
## <none>                93667 249.39
## + Area      1      4436  89232 249.93
## + Nearest    1      198  93469 251.33
## - Adjacent   1     69860 163527 264.11
## - Elevation  1    275784 369451 288.56
##
## Step:  AIC=249.35
## Species ~ Elevation + Adjacent
##
##           Df Sum of Sq   RSS   AIC
## <none>                100003 249.35
## + Scruz      1      6336  93667 249.39
## + Area      1      3227  96776 250.37
## + Nearest    1     1550  98453 250.88
## - Adjacent   1     73251 173254 263.84
## - Elevation  1    280817 380820 287.47
##
##
## Call:
## lm(formula = Species ~ Elevation + Adjacent, data = galaNew)
##
## Coefficients:
## (Intercept)      Elevation      Adjacent
##      1.43287       0.27657      -0.06889
```

Model Diagnostics

Residual Plot


```
mod <- lm(Species ~ Elevation + Adjacent, data = galaNew)
plot(mod$fitted.values, mod$residuals, pch = 16, col = "blue")
abline(h = 0, col = "red")
```




Residual Histogram/QQplot

```
par(las = 1)
hist(mod$residuals, 12, prob = T,
     col = "lightblue", border = "gray")
xg <- seq(-200, 200, 1)
yg <- dnorm(xg, 0, 60.86)
lines(xg, yg)
```


Histogram of mod\$residuals




```
plot(qnorm(1:30 / 31, 0, 60.86), sort(mod$residuals), pch = 16,  
     col = "gray", xlab = "Normal Quantiles", ylab = "Residuals")  
abline(0, 1)
```



Leverage

```
step_gala <- step(full, trace = F)
X <- model.matrix(step_gala)
H <- X %*% solve((t(X) %*% X)) %*% t(X)
lev <- hat(X)
high_lev <- which(lev >= 2 * 3 / 30)
attach(gala)

par(las = 1)
plot(Elevation, Adjacent, cex = sqrt(5 * lev), col = "blue", ylim = c(0, 5000))
points(Elevation[high_lev], Adjacent[high_lev], col = "red", pch = 16,
       cex = sqrt(5 * lev[high_lev]))
```




Studentized Residuals

```
gs <- summary(step_gala)
gs$sig
```

```
## [1] 60.85898
```

```
studRes <- gs$res / (gs$sig * sqrt(1 - lev))
plot(step_gala$fitted.values, studRes, pch = 16, cex = 0.8, col = "blue",
      ylab = expression(r[i]), main = "Studentized Residuals", xlab = "")
abline(h = 0, lty = 2, col = "gray")
```

Studentized Residuals




Jackknife Residuals

```
jack <- rstudent(step_gala)

par(las = 1)
plot(jack, pch = 16, cex = 0.8, col = "blue", main = " Jackknife Residuals ",
      xlab = "", ylab = "")
abline(h = 0, lty = 2, col = "gray")
```


Jackknife Residuals



Identifying Influential Observations: DFFITS

```
library(olsrr)
ols_plot_dffits(step_gala)
```


Influence Diagnostics for Species



Transformation

```
par(las = 1)
plot(step_gala$fitted.values, step_gala$residuals,
     pch = 16, cex = 0.8, col = "blue", main = " Residuals ",
     xlab = expression(hat(Y)), ylab = expression(e))
abline(h = 0, lty = 2, col = "gray")
```

Residuals



```
sqrt_fit <- lm(sqrt(Species) ~ Elevation + Adjacent)

par(las = 1)
plot(sqrt_fit$fitted.values, sqrt_fit$residuals,
     pch = 16, cex = 0.8, col = "blue", main = "Residuals ",
     xlab = expression(hat(Y)), ylab = expression(e))
abline(h = 0, lty = 2, col = "gray")
```

Residuals

