

Lecture 3

Completely Randomized Designs: Model, Estimation, Inference

STAT 8050 Design and Analysis of Experiments
January 16, 2020

Whitney Huang
Clemson University

Let Y_{ij} be the random variable that represents the response for the j^{th} experimental unit to treatment i . Also, let $\mu_i = E(Y_{ij})$ be the mean response for the i^{th} treatment. We have

$$Y_{ij} = \mu_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

where ϵ_{ij} is the random variable representing error associated with Y_{ij} with $E(\epsilon_{ij}) = 0$. This is called a **means model**.

Let Y_{ij} be the random variable that represents the response for the j^{th} experimental unit to treatment i . Also, let $\mu_i = E(Y_{ij})$ be the mean response for the i^{th} treatment. We have

$$Y_{ij} = \mu_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

where ϵ_{ij} is the random variable representing error associated with Y_{ij} with $E(\epsilon_{ij}) = 0$. This is called a **means model**.

Alternatively, we could let $\mu_i = \mu + \alpha_i$, which leads to

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i.$$

This is called an **effects model**

In both the **means model** and the **effects model**. We further assume

$$\epsilon_{ij} \sim N(0, \sigma^2),$$

and ϵ_{ij} 's are independent to each other.

This yields

$$Y_{ij} \sim N(\mu + \alpha_i, \sigma^2)$$

Effects Model

$$Y_{ij} \sim N(\mu_i, \sigma^2)$$

Means Model

In both the **means model** and the **effects model**. We further assume

$$\epsilon_{ij} \sim N(0, \sigma^2),$$

and ϵ_{ij} 's are independent to each other.

This yields

$$Y_{ij} \sim N(\mu + \alpha_i, \sigma^2)$$

Effects Model

$$Y_{ij} \sim N(\mu_i, \sigma^2)$$

Means Model

Note: We make the common variance assumption here

The model

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

- is overparameterized

The model

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

- is overparameterized

The model

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

- is overparameterized
- is nonidentifiable

The model

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

- is overparameterized
- is nonidentifiable

The model

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

- is overparameterized
- is nonidentifiable

Example

Suppose $g = 2$, then we have to estimate μ , α_1 , and α_2 .

$$\begin{aligned} \mu = 10, \alpha_1 = -1, \alpha_2 = 1, \\ \text{and } \mu = 11, \alpha_1 = -2, \alpha_2 = 0. \end{aligned}$$

The model

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad i = 1, \dots, g, \quad j = 1, \dots, n_i,$$

- is overparameterized
- is nonidentifiable

Example

Suppose $g = 2$, then we have to estimate μ, α_1 , and α_2 .

$$\begin{aligned} \mu = 10, \alpha_1 = -1, \alpha_2 = 1, \\ \text{and } \mu = 11, \alpha_1 = -2, \alpha_2 = 0. \end{aligned}$$

\Rightarrow each yield $Y_{1j} \sim N(9, \sigma^2)$ and $Y_{2j} \sim N(11, \sigma^2)$

- $\bar{Y}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ — i^{th} treatment mean

- $\bar{Y}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ — i^{th} treatment mean

- $\bar{Y}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ – i^{th} treatment mean
- $Y_{i\cdot} = \sum_{j=1}^{n_i} Y_{ij}$ – Total for i^{th} treatment

- $\bar{Y}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ – i^{th} treatment mean
- $Y_{i\cdot} = \sum_{j=1}^{n_i} Y_{ij}$ – Total for i^{th} treatment

- $\bar{Y}_i = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ – i^{th} treatment mean
- $Y_i = \sum_{j=1}^{n_i} Y_{ij}$ – Total for i^{th} treatment
- $Y_{..} = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij} = \sum_{i=1}^J Y_{i.}$ – Total of all observations

- $\bar{Y}_i = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ – i^{th} treatment mean
- $Y_i = \sum_{j=1}^{n_i} Y_{ij}$ – Total for i^{th} treatment
- $Y_{..} = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij} = \sum_{i=1}^J Y_{i.}$ – Total of all observations

- $\bar{Y}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} Y_{ij}$ – i^{th} treatment mean
- $Y_{i\cdot} = \sum_{j=1}^{n_i} Y_{ij}$ – Total for i^{th} treatment
- $Y_{\cdot\cdot} = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij} = \sum_{i=1}^j Y_{i\cdot}$ – Total of all observations
- $\bar{Y}_{\cdot\cdot} = \frac{1}{N} \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}$ – Grand mean of all observations where $N = \sum_{i=1}^g n_i$

Least Squares Estimation

To estimate $\mu, \alpha_1, \dots, \alpha_g$, we find the values for these parameters that minimize

$$\sum_{i=1}^g \sum_{j=1}^{n_i} e_{ij}^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - (\mu + \alpha_i))^2.$$

To estimate $\mu, \alpha_1, \dots, \alpha_g$, we find the values for these parameters that minimize

$$\sum_{i=1}^g \sum_{j=1}^{n_i} e_{ij}^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - (\mu + \alpha_i))^2.$$

To obtain the estimates, we have a system of $g + 1$ equations with $g + 1$ unknowns. Unfortunately, we only have g treatment means that can be used to solve this system of equations \Rightarrow **no unique solution exists for $\hat{\mu}, \hat{\alpha}_1, \dots, \hat{\alpha}_g$**

To estimate $\mu, \alpha_1, \dots, \alpha_g$, we find the values for these parameters that minimize

$$\sum_{i=1}^g \sum_{j=1}^{n_i} e_{ij}^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - (\mu + \alpha_i))^2.$$

To obtain the estimates, we have a system of $g + 1$ equations with $g + 1$ unknowns. Unfortunately, we only have g treatment means that can be used to solve this system of equations \Rightarrow **no unique solution exists for $\hat{\mu}, \hat{\alpha}_1, \dots, \hat{\alpha}_g$**

Typically constraints are used to obtain solutions and hence estimators.

Note: Different software uses different constraints

Constraints

| Constraint | $\hat{\mu}$ | $\hat{\alpha}_i$ | $\hat{\mu} + \hat{\alpha}_i$ | $\hat{\alpha}_i - \hat{\alpha}_{i'}$ |
|---------------------------------------|-------------|------------------|------------------------------|--------------------------------------|
| $\hat{\alpha}_g = 0$ | | | | |
| $\hat{\mu} = 0$ | | | | |
| $\sum_{i=1}^g n_i \hat{\alpha}_i = 0$ | | | | |

| Constraint | $\hat{\mu}$ | $\hat{\alpha}_i$ | $\hat{\mu} + \hat{\alpha}_i$ | $\hat{\alpha}_i - \hat{\alpha}_{i'}$ |
|---------------------------------------|-------------|------------------|------------------------------|--------------------------------------|
| $\hat{\alpha}_g = 0$ | | | | |
| $\hat{\mu} = 0$ | | | | |
| $\sum_{i=1}^g n_i \hat{\alpha}_i = 0$ | | | | |

$\hat{\mu}$ and $\hat{\alpha}_i$ depends upon the constraint used. $\hat{\mu} + \hat{\alpha}_i$ and $\hat{\alpha}_i - \hat{\alpha}_{i'}$ are **invariant** to the constraint used.

| Constraint | $\hat{\mu}$ | $\hat{\alpha}_i$ | $\hat{\mu} + \hat{\alpha}_i$ | $\hat{\alpha}_i - \hat{\alpha}_{i'}$ |
|---------------------------------------|-------------|------------------|------------------------------|--------------------------------------|
| $\hat{\alpha}_g = 0$ | | | | |
| $\hat{\mu} = 0$ | | | | |
| $\sum_{i=1}^g n_i \hat{\alpha}_i = 0$ | | | | |

$\hat{\mu}$ and $\hat{\alpha}_i$ depends upon the constraint used. $\hat{\mu} + \hat{\alpha}_i$ and $\hat{\alpha}_i - \hat{\alpha}_{i'}$ are **invariant** to the constraint used.

Note: If we use the **means model**, $\hat{\mu}_i = \bar{Y}_{i\cdot}$, and we do not have these issues here, but we will have other issues later on.

The total variation is represented by total sum of squares SS_T :

$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2$$

This quantity can be decomposed to variation between treatments (SS_{TRT}) and variation within treatment (SS_E):

$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2 = \underbrace{\sum_{i=1}^g n_i (\bar{Y}_{i.} - \bar{Y}_{..})^2}_{SS_{TRT}} + \underbrace{\sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{i.})^2}_{SS_E}$$

Equivalent Computational Formulae

Completely
Randomized
Designs: Model,
Estimation, Inference



$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \frac{Y_{..}^2}{N}$$

Equivalent Computational Formulae

Completely
Randomized
Designs: Model,
Estimation, Inference



$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \frac{Y_{..}^2}{N}$$

$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \frac{Y_{..}^2}{N}$$

$$SS_{TRT} = \sum_{i=1}^g n_i (\bar{Y}_{i.} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \sum_{i=1}^g \frac{Y_{i.}^2}{n_i}$$

$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \frac{Y_{..}^2}{N}$$

$$SS_{TRT} = \sum_{i=1}^g n_i (\bar{Y}_{i.} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \sum_{i=1}^g \frac{Y_{i.}^2}{n_i}$$

Equivalent Computational Formulae

Completely
Randomized
Designs: Model,
Estimation, Inference



$$SS_T = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \frac{Y_{..}^2}{N}$$

$$SS_{TRT} = \sum_{i=1}^g n_i (\bar{Y}_{i.} - \bar{Y}_{..})^2 = \sum_{i=1}^g \sum_{j=1}^{n_i} Y_{ij}^2 - \sum_{i=1}^g \frac{Y_{i.}^2}{n_i}$$

$$SS_E = \sum_{i=1}^g \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_{i.})^2 = \sum_{i=1}^g \frac{Y_{i.}^2}{n_i} - \frac{Y_{..}^2}{N}$$

Mean Squares

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ **total degrees of freedom** (Why?)

Mean Squares

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ **total degrees of freedom** (Why?)

Mean Squares

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ total degrees of freedom (Why?)
- We have ____ treatment degrees of freedom \Rightarrow

Mean Squares

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ total degrees of freedom (Why?)
- We have ____ treatment degrees of freedom \Rightarrow

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ total degrees of freedom (Why?)
- We have ____ treatment degrees of freedom \Rightarrow

$$MS_{TRT} = \frac{SS_{TRT}}{g - 1}$$

- We have ____ error degrees of freedom \Rightarrow

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ total degrees of freedom (Why?)
- We have ____ treatment degrees of freedom \Rightarrow

$$MS_{TRT} = \frac{SS_{TRT}}{g - 1}$$

- We have ____ error degrees of freedom \Rightarrow

Dividing mean squares by their associated degrees of freedom yield “variance-like” quantities called **mean squares**.

- We have $N - 1$ total degrees of freedom (Why?)
- We have ____ treatment degrees of freedom \Rightarrow

$$MS_{TRT} = \frac{SS_{TRT}}{g - 1}$$

- We have ____ error degrees of freedom \Rightarrow

$$MS_E = \frac{SS_E}{N - g}$$

Note that

$$MS_E = \frac{1}{N - g} \underbrace{\sum_{i=1}^g (n_i - 1) s_i^2}_{SS_E}$$

provides an **unbiased** estimator of σ^2 **regardless of whether the treatment population means differ or not.**

Note that

$$MS_E = \frac{1}{N - g} \underbrace{\sum_{i=1}^g (n_i - 1) s_i^2}_{SS_E}$$

provides an **unbiased** estimator of σ^2 **regardless of whether the treatment population means differ or not.**

Also, it can be shown that

$$MS_{TRT} = \frac{1}{g - 1} \underbrace{\sum_{i=1}^g n_i (\bar{Y}_i - \bar{Y}_{..})^2}_{SS_{TRT}}$$

is an **unbiased** estimator of σ^2 **if all treatment population means are equal.**

If

$$H_0 : \alpha_1 = \alpha_2 = \cdots = \alpha_g = 0$$

is true, then MS_{TRT} and MS_E will be “similar”. Otherwise, they will be different. We can show that

$$E(MS_{TRT}) = \sigma^2 + \sum_{i=1}^g n_i \alpha_i^2 / (g - 1) \geq \sigma^2 = E(MS_E)$$

\Rightarrow if H_0 is false, MS_{TRT} will tend to be larger than MS_E .

| Source | df | SS | MS | EMS |
|-----------|---------|------------|-----------------------------------|--|
| Treatment | $g - 1$ | SS_{TRT} | $MS_{TRT} = \frac{SS_{TRT}}{g-1}$ | $\sigma^2 + \frac{\sum_{i=1}^g n_i \alpha_i^2}{g-1}$ |
| Error | $N - g$ | SS_E | $MS_E = \frac{SS_E}{N-g}$ | σ^2 |
| Total | $N - 1$ | SS_T | | |

Testing for treatment effects

$$H_0 : \alpha_i = 0 \quad \text{for all } i$$

$$H_a : \alpha_i \neq 0 \quad \text{for some } i$$

Test statistics: $F = \frac{MS_{TRT}}{MS_E}$. Under H_0 , the test statistic follows an F-distribution with $g - 1$ and $N - g$ degrees of freedom

Reject H_0 if

$$F_{obs} > F_{g-1, N-g; \alpha}$$

for an α -level test, $F_{g-1, N-g; \alpha}$ is the $100 \times (1 - \alpha)\%$ percentile of a **central F-distribution** with $g - 1$ and $N - g$ degrees of freedom.

Reject H_0 if

$$F_{obs} > F_{g-1, N-g; \alpha}$$

for an α -level test, $F_{g-1, N-g; \alpha}$ is the $100 \times (1 - \alpha)\%$ percentile of a **central F-distribution** with $g - 1$ and $N - g$ degrees of freedom.

P-value

The **P-value** of the F-test is the probability of obtaining F at least as extreme as F_{obs} , that is, $P(F > F_{obs})$.

Reject H_0 if

$$F_{obs} > F_{g-1, N-g; \alpha}$$

for an α -level test, $F_{g-1, N-g; \alpha}$ is the $100 \times (1 - \alpha)\%$ percentile of a **central F-distribution** with $g - 1$ and $N - g$ degrees of freedom.

P-value

The **P-value** of the F-test is the probability of obtaining F at least as extreme as F_{obs} , that is, $P(F > F_{obs})$.

We reject H_0 if P-value $< \alpha$.

F Distribution and the F-Test

Consider the observed F test statistic: $F_{obs} = \frac{MS_{TRT}}{MS_E}$

F Distribution and the F-Test

Consider the observed F test statistic: $F_{obs} = \frac{MS_{TRT}}{MS_E}$

- Should be “near” 1 if the treatment means are equal

F Distribution and the F-Test

Consider the observed F test statistic: $F_{obs} = \frac{MS_{TRT}}{MS_E}$

- Should be “near” 1 if the treatment means are equal

F Distribution and the F-Test

Consider the observed F test statistic: $F_{obs} = \frac{MS_{TRT}}{MS_E}$

- Should be “near” 1 if the treatment means are equal
- Should be “larger than” 1 if treatment means are not equal

F Distribution and the F-Test

Consider the observed F test statistic: $F_{obs} = \frac{MS_{TRT}}{MS_E}$


- Should be “near” 1 if the treatment means are equal
- Should be “larger than” 1 if treatment means are not equal

F Distribution and the F-Test

Consider the observed F test statistic: $F_{obs} = \frac{MS_{TRT}}{MS_E}$

- Should be “near” 1 if the treatment means are equal
- Should be “larger than” 1 if treatment means are not equal

⇒ We use the null distribution $F \sim F_{df_1=g-1, df_2=N-g}$ to quantify if F_{obs} is large enough to reject H_0



Example

An experiment was conducted to determine if experience has an effect on the time it takes for mice to run a maze. Four treatment groups, consisting of mice having been trained on the maze one, two, three and four times were run through the maze and their times recorded. Three mice were originally assigned to each group, but it was discovered that some lab assistants, in an attempt to win a bet, gave one mouse a stimulant and another mouse a sedative. These mice were removed from the analysis.

| Training runs | 1 | 2 | 3 | 4 |
|---------------|-------|-------|-------|-----|
| Times | 11, 9 | 7,8,9 | 6,5,7 | 5,3 |
| y_i | 20 | 24 | 18 | 8 |
| n_i | 2 | 3 | 3 | 2 |
| s_i^2 | | | | |

Example Cont'd

| Training runs | 1 | 2 | 3 | 4 |
|---------------|-------|-------|-------|-----|
| Times | 11, 9 | 7,8,9 | 6,5,7 | 5,3 |
| y_i | 20 | 24 | 18 | 8 |
| n_i | 2 | 3 | 3 | 2 |
| s_i^2 | | | | |

- Write down the model.

Example Cont'd

| Training runs | 1 | 2 | 3 | 4 |
|---------------|-------|-------|-------|-----|
| Times | 11, 9 | 7,8,9 | 6,5,7 | 5,3 |
| y_i | 20 | 24 | 18 | 8 |
| n_i | 2 | 3 | 3 | 2 |
| s_i^2 | | | | |

- Write down the model.

Example Cont'd

| Training runs | 1 | 2 | 3 | 4 |
|---------------|-------|-------|-------|-----|
| Times | 11, 9 | 7,8,9 | 6,5,7 | 5,3 |
| y_i | 20 | 24 | 18 | 8 |
| n_i | 2 | 3 | 3 | 2 |
| s_i^2 | | | | |

- Write down the model.
- Fill out the ANOVA table and test whether the time to run the maze is affected by training. Use a significant level of .05.