

Lecture 5

Multivariate Linear Regression

Readings: Johnson & Wichern 2007, Chapter 7; DSA 8020

Lectures 1-4 [\[Link\]](#); Zelterman, 2015, Chapter 9

DSA 8070 Multivariate Analysis

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Whitney Huang
Clemson University

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

- 1 Motivation
- 2 Model and Assumptions
- 3 Parameter Estimation
- 4 Inference and Prediction
- 5 Motor Trend Car Road Tests Data Analysis

Example: Motor Trend Car Road Tests

```
> head(mtcars)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1
Hornet 4 Drive	21.4	6	258	110	3.08	3.215	19.44	1	0	3	1
Hornet Sportabout	18.7	8	360	175	3.15	3.440	17.02	0	0	3	2
Valiant	18.1	6	225	105	2.76	3.460	20.22	1	0	3	1

Suppose we would like to study the (linear) relationship between mpg, disp, hp, wt (responses) and cyl, am, carb (predictors)

Multiple regression predicts one outcome; multivariate regression predicts several simultaneously

Why Multivariate Regression Instead of Separate Regressions?

- **Estimation:**

- Coefficient estimates are the same as running separate regressions

- **Inference:** The real gain comes from joint modeling

- Test hypotheses across multiple outcomes simultaneously
- Accounts for correlations among responses \Rightarrow more powerful and accurate tests

- **Examples:**

- Does a predictor affect all outcomes jointly?
- Multivariate analog of ANOVA (MANOVA)

The multiple linear regression model has the form:

$$y_i = \beta_0 + \sum_{j=1}^p \beta_j x_{ij} + \varepsilon_i, \quad i = 1, \dots, n,$$

where

- y_i is the **response** for the i -th observation
- x_{ij} is the j -th **predictor** for the i -th observation
- β_0 and β_j 's are the **regression intercept** and **slopes** for the response, respectively
- ε_i is the **error** term for the response of the i -th observation

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

The Multivariate Linear Regression Model: Scalar Form

The multivariate (multiple) linear regression model has the form:

$$y_{ik} = \beta_{0k} + \sum_{j=1}^p \beta_{jk} x_{ij} + \varepsilon_{ik}, \quad i = 1, \dots, n, \quad k = 1, \dots, d,$$

where

- y_{ik} is the k -th **response** for the i -th observation
- x_{ij} is the j -th **predictor** for the i -th observation
- β_{0k} and β_{jk} 's are the **regression intercept** and **slopes** for k -th response, respectively
- ε_{ik} is the **error** term for the k -th response of the i -th observation

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

The Multivariate Linear Regression Model: Assumptions

The assumptions of the model are:

- Relationship between $\{x_j\}_{j=1}^p$ and Y_k is **linear** for each $k \in \{1, \dots, d\}$
- $(\varepsilon_{i1}, \dots, \varepsilon_{id})^T \stackrel{i.i.d.}{\sim} N(\mathbf{0}, \Sigma)$ is an **unobserved random vector**
- $[Y_{ik} | x_{i1}, \dots, x_{ip}] \sim N(\beta_{0k} + \sum_{j=1}^p \beta_{jk} x_{ij}, \sigma_{kk})$ for each $k \in \{1, \dots, d\}$

The Multivariate Linear Regression Model: Matrix Form

The multivariate multiple linear regression model has the form

$$Y = XB + E,$$

where

- $Y = [y_1, \dots, y_d]$ is the $n \times d$ **response matrix**, where $y_k = (y_{1k}, \dots, y_{nk})^T$ is the k -th response vector
- $X = [1, x_1, \dots, x_p]$ is the $n \times (p + 1)$ **design matrix**
- $B = [\beta_1, \dots, \beta_d]$ is the $(p + 1) \times d$ **matrix of regression coefficients**
- $E = [\epsilon_1, \dots, \epsilon_d]$ is the $n \times d$ **error matrix**

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Matrix form writes the multivariate linear regression model for all $n \times d$ points simultaneously as

$$\mathbf{Y} = \mathbf{X}\mathbf{B} + \mathbf{E}$$

$$\begin{bmatrix} y_{11} & \cdots & y_{1d} \\ y_{21} & \cdots & y_{2d} \\ \vdots & \ddots & \vdots \\ y_{n1} & \cdots & y_{nd} \end{bmatrix} = \begin{bmatrix} 1 & \cdots & x_{1p} \\ 1 & \cdots & x_{2p} \\ \vdots & \ddots & \vdots \\ 1 & \cdots & x_{np} \end{bmatrix} \begin{bmatrix} \beta_{01} & \cdots & \beta_{0d} \\ \beta_{11} & \cdots & \beta_{1d} \\ \vdots & \ddots & \vdots \\ \beta_{p1} & \cdots & \beta_{pd} \end{bmatrix} + \begin{bmatrix} \varepsilon_{11} & \cdots & \varepsilon_{1d} \\ \varepsilon_{21} & \cdots & \varepsilon_{2d} \\ \vdots & \ddots & \vdots \\ \varepsilon_{n1} & \cdots & \varepsilon_{nd} \end{bmatrix}$$

Assuming that n subjects are **independent**, we have

$$\bullet \varepsilon_k \sim N(0, \sigma_{kk}), \quad k \in \{1, \dots, d\}$$

$$\bullet \varepsilon_i \stackrel{i.i.d.}{\sim} N(\mathbf{0}, \Sigma), \quad i = 1, \dots, n$$

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

The **ordinary least squares** OLS estimate is

$$\operatorname{argmin}_{\mathbf{B} \in \mathbb{R}^{(p+1) \times d}} \|\mathbf{Y} - \mathbf{X}\mathbf{B}\|^2 = \operatorname{argmin}_{\mathbf{B} \in \mathbb{R}^{(p+1) \times d}} \sum_{i=1}^n \sum_{k=1}^d \left(y_{ik} - \beta_{0k} - \sum_{j=1}^p \beta_{jk} x_{ij} \right)^2,$$

where $\|\cdot\|$ denotes the Frobenius norm.

- $\text{OLS}(\mathbf{B}) = \|\mathbf{Y} - \mathbf{X}\mathbf{B}\|^2 = \text{tr}(\mathbf{Y}^T \mathbf{Y}) - 2\text{tr}(\mathbf{Y}^T \mathbf{X}\mathbf{B}) + \text{tr}(\mathbf{B}^T \mathbf{X}^T \mathbf{X} \mathbf{B})$
- $\frac{\partial \text{OLS}(\mathbf{B})}{\partial \mathbf{B}} = -2\mathbf{X}^T \mathbf{Y} + 2\mathbf{X}^T \mathbf{X} \mathbf{B}$

The OLS estimate has the form

$$\hat{\mathbf{B}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} \Rightarrow \hat{\beta}_k = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}_k, \quad k \in \{1, \dots, d\}$$

Expected Value of Least Squares Coefficients

The expected value of the estimated coefficients is given by

$$\begin{aligned}\mathbb{E}(\hat{B}) &= \mathbb{E}[(X^T X)^{-1} X^T Y] \\ &= (X^T X)^{-1} X^T \mathbb{E}(Y) \\ &= (X^T X)^{-1} X^T X B \\ &= B\end{aligned}$$

$\Rightarrow \hat{B}$ is an unbiased estimator of B

- Fitted values are given by

$$\hat{Y} = X\hat{B},$$

$$\text{i.e., } \hat{y}_{ik} = \hat{\beta}_{0k} + \sum_{j=1}^p \hat{\beta}_{jk} x_{ij}, \quad i = 1, \dots, n, \quad k = 1, \dots, d$$

- Residuals are given by

$$\hat{E} = Y - \hat{Y},$$

$$\text{i.e., } \hat{e}_{ik} = y_{ik} - \hat{y}_{ik}, \quad i = 1, \dots, n, \quad k = 1, \dots, d$$

Just like in univariate linear regression we can write the fitted values as

$$\begin{aligned}\hat{Y} &= X\hat{B} \\ &= X(X^T X)^{-1} X^T Y \\ &= HY,\end{aligned}$$

where $H = X(X^T X)^{-1} X^T$ is the **hat matrix**

$\Rightarrow H$ projects y_k onto the column space of X for $k \in \{1, \dots, d\}$

Partitioning the Total Variation

We can partition the total covariation in $\{\mathbf{y}_i\}_{i=1}^n$ (SSCP_{Tot}) as

$$\begin{aligned}\text{SSCP}_{\text{tot}} &= \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})^T (\mathbf{y}_i - \bar{\mathbf{y}}) \\&= \sum_{i=1}^n (\mathbf{y}_i - \hat{\mathbf{y}}_i + \hat{\mathbf{y}}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \hat{\mathbf{y}}_i + \hat{\mathbf{y}}_i - \bar{\mathbf{y}})^T \\&= \underbrace{\sum_{i=1}^n (\hat{\mathbf{y}}_i - \bar{\mathbf{y}})(\hat{\mathbf{y}}_i - \bar{\mathbf{y}})^T}_{\text{SSCP}_{\text{Reg}}} + \underbrace{\sum_{i=1}^n (\mathbf{y}_i - \hat{\mathbf{y}}_i)(\mathbf{y}_i - \hat{\mathbf{y}}_i)^T}_{\text{SSCP}_{\text{Err}}} \\&\quad + \underbrace{2 \sum_{i=1}^n (\hat{\mathbf{y}}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \hat{\mathbf{y}}_i)^T}_{=0} \\&= \text{SSCP}_{\text{Reg}} + \text{SSCP}_{\text{Err}}\end{aligned}$$

The corresponding **degrees of freedom** are $d(n-1)$ for SSCP_{Tot} ; dp for SSCP_{Reg} ; and $d(n-p-1)$ for SSCP_{Err}

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

The estimated error covariance matrix is

$$\begin{aligned}\hat{\Sigma} &= \frac{\sum_{i=1}^n (\mathbf{y}_i - \hat{\mathbf{y}}_i)(\mathbf{y}_i - \hat{\mathbf{y}}_i)^T}{n - p - 1} \\ &= \frac{\text{SSCP}_{Err}}{n - p - 1}\end{aligned}$$

- $\hat{\Sigma}$ is an unbiased estimate of Σ
- The estimate $\hat{\Sigma}$ is the mean SSCP_{Err}

Sampling Distributions of \hat{B} , \hat{Y} , and \hat{E}

We would need to figure out the **sampling distributions** of estimator and predictor in order to draw inference

Given the model assumptions, we have

$$\text{vec}(\hat{B}) \sim N(\text{vec}(B), \Sigma \otimes (X^T X)^{-1})$$

$$\text{vec}(\hat{Y}) \sim N(\text{vec}(XB), \Sigma \otimes H)$$

$$\text{vec}(\hat{E}) \sim N(0, \Sigma \otimes (I - H)),$$

where $\text{vec}(\cdot)$ is the vectorization operator and \otimes is the Kronecker product

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Inference about Multiple $\hat{\beta}_{jk}$

Assume that $q < p$ and want to test if a reduced model is sufficient:

$$H_0 : B_2 = \mathbf{0}_{p-q} \times d, \quad \text{versus} \quad H_a : B_2 \neq \mathbf{0}_{p-q} \times d,$$

where

$$B = \begin{bmatrix} B_1 \\ B_2 \end{bmatrix}$$

is the partitioned of the coefficient vector

We can compare the SSCP_{Err} for the **full model**:

$$y_{ik} = \beta_{0k} + \sum_{j=1}^p \beta_{jk} x_{ij} + \varepsilon_{ik}, \quad k = 1, \dots, d$$

and the **reduced model**:

$$y_{ik} = \beta_{0k} + \sum_{j=1}^q \beta_{jk} x_{ij} + \varepsilon_{ik}, \quad k = 1, \dots, d$$

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Some Test Statistics

Let $\tilde{E} = n\tilde{\Sigma}$ denote the SSCP_{Err} matrix from the **full model**,
and let $\tilde{H} = n(\tilde{\Sigma}_1 - \tilde{\Sigma})$ denote the hypothesis SSCP_{Err} matrix
Some test statistics for

$$H_0 : B_2 = \mathbf{0}_{p-q} \times d, \quad \text{versus} \quad H_a : B_2 \neq \mathbf{0}_{p-q} \times d :$$

- Wilks's Lambda

$$\Lambda^* = \frac{|\tilde{E}|}{|\tilde{H} + \tilde{E}|}$$

Reject H_0 if Λ^* is “small”

- Hotelling-Lawley Trace

$$T_0^2 = \text{tr}(\tilde{H}\tilde{E}^{-1})$$

Reject H_0 if T_0^2 is “large”

- Pillai's Trace

$$V = \text{tr}(\tilde{H}(\tilde{H} + \tilde{E})^{-1})$$

Reject H_0 if V is “large”

We would like to estimate the **expected value of the response** for a given predictor $\mathbf{x}_h = (1, x_{h1}, \dots, x_{hp})$.

Note that we have

$$\hat{\mathbf{y}}_h \sim N(\mathbf{B}^T \mathbf{x}_h, \mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h \Sigma)$$

We can exploit the duality between interval estimation and hypothesis testing. That is, we can test

$$H_0 : \mathbb{E}(\mathbf{y}_h) = \mathbf{y}_h^* \text{ versus } H_a : \mathbb{E}(\mathbf{y}_h) \neq \mathbf{y}_h^*$$

The $100(1 - \alpha)\%$ confidence region is the collection of \mathbf{y}_h^* values that fail to reject H_0 at α level

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Test statistics:

$$T^2 = \left(\frac{\hat{B}^T \mathbf{x}_h - B^T \mathbf{x}_h}{\sqrt{\mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h}} \right)^T \hat{\Sigma}^{-1} \left(\frac{\hat{B}^T \mathbf{x}_h - B^T \mathbf{x}_h}{\sqrt{\mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h}} \right)$$
$$\stackrel{H_0}{\sim} \frac{d(n-p-1)}{n-p-d} F_{d,n-p-d}$$

Therefore, the $100(1 - \alpha)\%$ simultaneous **confidence interval** for y_{hk} is

$$\hat{y}_{hk} \pm \sqrt{\frac{d(n-p-1)}{n-p-d} F_{d,n-p-d}} \sqrt{\mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h} \hat{\sigma}_{kk},$$

$$k \in \{1, \dots, d\}$$

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Here we want to predict the **observed value of response** for a given predictor

- **Note:** interested in actual \hat{y}_h instead of $\mathbb{E}(\hat{y}_h)$
- Given $\mathbf{x}_h = (1, x_{h1}, \dots, x_{hp})$, the fitted value is still $\hat{y}_h = \hat{\mathbf{B}}^T \mathbf{x}_h$

We can exploit the duality between interval estimation and hypothesis testing. That is, we can test

$$H_0 : \mathbf{y}_h = \mathbf{y}_h^* \text{ versus } H_a : \mathbf{y}_h \neq \mathbf{y}_h^*$$

The $100(1 - \alpha)\%$ prediction interval is the collection of \mathbf{y}_h^* values that fail to reject H_0 at α level

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Test statistics:

$$T^2 = \left(\frac{\hat{B}^T \mathbf{x}_h - B^T \mathbf{x}_h}{\sqrt{1 + \mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h}} \right)^T \hat{\Sigma}^{-1} \left(\frac{\hat{B}^T \mathbf{x}_h - B^T \mathbf{x}_h}{\sqrt{1 + \mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h}} \right)$$
$$\stackrel{H_0}{\sim} \frac{d(n-p-1)}{n-p-d} F_{d, n-p-d}$$

Therefore, the $100(1 - \alpha)\%$ simultaneous **prediction interval** for y_{hk} is

$$\hat{y}_{hk} \pm \sqrt{\frac{d(n-p-1)}{n-p-d} F_{d, n-p-d}} \sqrt{(1 + \mathbf{x}_h^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_h)} \hat{\sigma}_{kk},$$

$$k \in \{1, \dots, d\}$$

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

Motor Trend Car Road Tests Data Analysis

Study the linear relationship between mpg, disp, hp, wt (responses) and cyl, am, carb (predictors) in the mtcars dataset



Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis



- **Model:** `lm(Y ~ cyl + am + carb, data = mtcars)`
- **Key findings:**
 - `mpg`: decreases with more cylinders, increases with manual transmission
 - `disp`: strongly influenced by `cyl`
 - `hp`: influenced by `cyl` and `carb`
 - `wt`: influenced by `cyl`, `am`, `carb`
- **Note:** Multivariate regression produces the *same point estimates* as running separate regressions for each response

SSCP decomposition:

$$SSCP_{Tot} = SSCP_{Reg} + SSCP_{Err}$$

- Estimated error covariance matrix ($\hat{\Sigma}$)

	mpg	disp	hp	wt
mpg	7.8680094	-53.27166	-19.7015979	-0.6575443
disp	-53.2716607	2504.87095	425.1328988	18.1065416
hp	-19.7015979	425.13290	577.2703337	0.4662491
wt	-0.6575443	18.10654	0.4662491	0.2573503
	mpg	disp	hp	wt
mpg	1.0000000	-0.3794645	-0.29233405	-0.46209388
disp	-0.3794645	1.0000000	0.35354314	0.71314929
hp	-0.2923340	0.3535431	1.00000000	0.03825304
wt	-0.4620939	0.7131493	0.03825304	1.00000000

⇒ Captures residual dependencies among responses

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
PredictionMotor Trend Car Road
Tests Data Analysis

Do We Need “cyl”?

```
> mvlm0 <- lm(Y ~ am + carb, data = mtcars)
> anova(mvlm, mvlm0, test = "Wilks")
```

Analysis of Variance Table

Model 1: $Y \sim \text{cyl} + \text{am} + \text{carb}$

Model 2: $Y \sim \text{am} + \text{carb}$

	Res.Df	Df	Gen.var.	Wilks	approx F	num Df	Df
1	27		29.862				
2	29	2	43.692	0.16395	8.8181		8

	den	Df	Pr(>F)
1			
2	48		2.525e-07 ***

Interpretation: cylinder count explains variation across responses

Default `predict()` lacks multivariate CI/PI, so we use an R function from Prof. Helwig at the University of Minnesota

```
> newdata <- data.frame(cyl = factor(6, levels = c(4, 6, 8)),  
+                        am = 1, carb = 4)  
> # confidence interval  
> pred.mlm(mvmlm, newdata)  
      mpg      disp      hp      wt  
fit 21.51824 159.2707 136.98500 2.631108  
lwr 16.65593  72.5141  95.33649 1.751736  
upr 26.38055 246.0273 178.63351 3.510479  
> # prediction interval  
> pred.mlm(mvmlm, newdata, interval = "prediction")  
      mpg      disp      hp      wt  
fit 21.518240 159.27070 136.98500 2.6311076  
lwr  9.680053 -51.95435  35.58397 0.4901152  
upr 33.356426 370.49576 238.38603 4.7720999
```

Motivation

Model and
Assumptions

Parameter Estimation

Inference and
Prediction

Motor Trend Car Road
Tests Data Analysis

In this lecture, we learned about Multivariate Linear Regression

- Model and Assumptions
- Parameter Estimation
- Inference and Prediction