


# Conditional Approaches to Estimating Concurrent Wind and Precipitation Extremes

Whitney Huang

Joint work with Francis Zwiers and Adam Monahan



Advances and Challenges in Wind Modeling and its Applications  
August 12, 2020



University  
of Victoria



samsi  
NSF Duke NCSU UNC

# Outline of the talk

- ▶ Concurrent extremes: **simultaneous** occurrence of extreme values for **multiple** climate variables
- ▶ Conditional approaches for estimating concurrent extremes:
  - ▶ **Quantile regression**
  - ▶ **Conditional extreme value models**
- ▶ Estimating concurrent extremes using a large ensemble
  - ▶ Estimating simultaneous extreme wind speed and precipitation
  - ▶ 35-member Regional Climate Model initial-condition ensemble

## Some examples of concurrent extreme events



Credit: Shutterstock



Source: [www.standardmedia.co.ke](http://www.standardmedia.co.ke)




## Estimating the magnitude of concurrent extremes

Consider the bivariate case, i.e.,  $\mathbf{X} = (X_1, X_2)^T$

- ▶ There is no natural ordering to define an extreme value for multivariate data
- ▶ Traditional multivariate extreme value analysis mainly focus on modeling component-wise maximum  $\Rightarrow$  may lead to “wrong” events 😞
- ▶ It is important to account for “event simultaneity” for modeling concurrent extremes  $\Rightarrow$  we do this by conditioning on one variable being extremes

# Classical multivariate extreme value analysis

Modeling **componentwise maxima** using multivariate extreme value distribution (extreme-value marginals + tail copula)





**Issue:** Ignore the event simultaneity

# Componentwise maxima vs. concomitants of maxima

**Red:** (annual max precip, annual max wind speed)

**Blue:** (annual max precip, concurrent wind speed)


**Green:** (annual max wind speed, concurrent precip)



## An illustration of conditional approach

Let  $X_1$  and  $X_2$  be daily precipitation and wind speed


1. Condition on  $X_1$  being “extreme” e.g., annual maximum



## An illustration of conditional approach

Let  $X_1$  and  $X_2$  be daily precipitation and wind speed

1. Condition on  $X_1$  being “extreme” e.g., annual maximum
2. Model the conditional distribution of  $X_2$  given  $X_1$  being “extreme”



## Estimating conditional distribution: Quantile regression (QR) [Koenker and Bassett, 1978]

- ▶ Extending regression analysis by estimating conditional quantiles:  $Q_{X_2}(\tau|X_1 = x_1) = \inf\{x_2 : F(x_2|x_1) \geq \tau\}, \tau \in (0, 1)$
- ▶ Estimate  $Q_{X_2}(\tau|x_1)$  at a finite number of quantile levels  $\tau_1, \tau_2, \dots, \tau_J$
- ▶ Could suffer from the issue of quantile curves crossing  $\Rightarrow$  We use Monotone Composite Quantile Regression Neural Network [MCQRNN, Cannon, 2018] to (non-parametrically) estimate quantile curves **simultaneously**

## Estimating conditional distribution: Conditional extreme value (CEV) models [Heffernan and Tawn, 04]


- ▶ **Marginal modeling:**
  1. Estimate marginal distributions of  $X_1$  and  $X_2$
  2. Transform  $(X_1, X_2)^T$  to Laplace marginals  $(Y_1, Y_2)^T$
- ▶ **Dependence modeling:**

Assume for large  $u$ ,

$[(Y_2 - a(Y_1)) / (b(Y_1)) \leq z | Y_1 > u] \sim G(z)$ , where  $a(y) = \alpha y$  and  $b(y) = y^\beta$ ,  $\alpha \in [-1, 1]$ ,  $\beta \in (-\infty, 1)$

## A cartoon illustration of the CEV dependence modeling

Assume for large  $u$ ,  $[(Y_2 - a(Y_1)) / (b(Y_1)) \leq z | Y_1 > u] \sim G(z)$ ,  
where  $a(y) = \alpha y$  and  $b(y) = y^\beta$ ,  $\alpha \in [-1, 1]$ ,  $\beta \in (-\infty, 1)$



- ▶  $Y_2 = \alpha Y_1 + Y_1^\beta Z$ ,  
 $Z = \frac{Y_2 - \alpha Y_1}{Y_1^\beta} \sim G$
- ▶  $\alpha$  and  $\beta$  are estimated by making a parametric assumption of  $Y_2$
- ▶  $G$  estimated nonparametrically

## “Data”


Output from CanRCM4, Canadian Regional Climate Model 4, 1950 – 2100

- ▶ 35-member initial-condition ensemble
- ▶ Using CMIP5 historical forcings (1950-2005) and then RCP 8.5 (2006-2100)
- ▶ North American region,  $0.44^{\circ}$  horizontal grid resolution ( $\sim 50$  km)


Each run in ensemble produces (nearly) statistically independent realizations of climate system, which allows us to:

- ▶ provide more accurate estimates and more realistic modeling in climate extremes
- ▶ assess how well statistical procedures work

# The three grid cells considered in this talk




# Estimating conditional quantiles using MCQRNN and CEV



- ▶ DJF max precipitation ↑ concurrent wind speed ↑
- ▶ QR and CEV yield reasonably close wind speed high quantile estimates
- ▶ Conditional quantiles are substantially larger than their unconditional counterparts



# Seasonal conditional upper quantiles



- ▶ Wind speeds and precipitation are larger during SON and DJF
- ▶ Both methods produce close high quantile estimates up to the 50-year return level but can diverge beyond that
- ▶ Conditional quantiles are substantially larger than their unconditional counterparts



# Resample ensemble runs to obtain uncertainty estimates

Here we show quantile function estimates with  $\tau = 0.9$



# A quick assessment of statistical model performance

- ▶ We treat the fitted conditional quantile function at  $\tau = 0.9$  using all 35 ensemble members as the “truth”
- ▶ We assess the model performance by fitting CEV and MCQRNN for each ensemble member



## Summary & Discussion

- ▶ We explore conditional approaches to estimate the concurrent wind and precipitation extremes
- ▶ Large climate model ensemble is a power tool for studying climate extremes
- ▶ **Ongoing work**
  - ▶ Nonstationary extension account for both seasonality and long term trend for marginal and dependence structures
  - ▶ Spatial extension to borrow strength across space to improve estimation of concurrent extremes
  - ▶ Estimating conditional upper tail of precipitation conditioning on extreme wind speed

## Summary & Discussion

- ▶ We explore conditional approaches to estimate the concurrent wind and precipitation extremes
- ▶ Large climate model ensemble is a power tool for studying climate extremes
- ▶ **Ongoing work**
  - ▶ Nonstationary extension account for both seasonality and long term trend for marginal and dependence structures
  - ▶ Spatial extension to borrow strength across space to improve estimation of concurrent extremes
  - ▶ Estimating conditional upper tail of precipitation conditioning on extreme wind speed

**Thank you for your attention!**

**ArXiv:** <https://arxiv.org/pdf/2006.08720.pdf>.