

Lecture 4

Multivariate Normal Distribution, Copula, and Nonparametric Density Estimation

Readings: Zelterman, 2015 Chapters 5, 6, 7, Izeman, 2008
Chapter 4.1, 4.3, 4.5

DSA 8070 Multivariate Analysis
September 12 - September 16, 2022

Whitney Huang
Clemson University

Agenda

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation



Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation

- 1 **Multivariate Normal Distribution**
- 2 **Geometry of the Multivariate Normal Density**
- 3 **Copula**
- 4 **Nonparametric Density Estimation**

The Multivariate Normal Distribution

Just as the **univariate normal distribution** tends to be the most important distribution in **univariate statistics**, the **multivariate normal distribution** is the most important distribution in **multivariate statistics**


- **Mathematical Simplicity:** It is easy to obtain multivariate methods based on the multivariate normal distribution
- **Central Limit Theorem:** *The **sample mean vector** is going to be approximately **multivariate normally distributed** when the sample size is sufficiently large*
- Many natural phenomena may be modeled using this distribution (perhaps after transformation)

Review: Univariate Normal Distributions

The probability density function of the normal distribution is

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right\},$$

where μ and σ^2 are its **mean** and **variance**, respectively.





$\left(\frac{x-\mu}{\sigma}\right)^2 = (x-\mu)(\sigma^2)^{-1}(x-\mu)$ is the squared statistical distance between x and μ in standard deviation units

Multivariate Normal Distributions

If we have a p -dimensional random vector that is distributed according to a **multivariate normal distribution** with mean vector $\boldsymbol{\mu} = (\mu_1, \mu_2, \dots, \mu_p)^T$ and covariance matrix $\boldsymbol{\Sigma} = \{(\sigma_{ij})\}$, the probability density function is

$$f(\mathbf{x}) = \frac{1}{2\pi^{\frac{p}{2}} |\boldsymbol{\Sigma}|^{\frac{1}{2}}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\}.$$



Review: Central Limit Theorem (CLT)

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation



The **sampling distribution** of the **mean** will become approximately **normally distributed** as the **sample size** becomes larger, **irrespective of the shape of the population distribution!**

Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation

Let X_1, X_2, \dots, X_n $\overset{i.i.d.}{\sim} F$ with $\mu = E[X_i]$ and $\sigma^2 = \text{Var}[X_i]$. Then $\bar{X}_n = \frac{\sum_{i=1}^n X_i}{n} \xrightarrow{d} N(\mu, \frac{\sigma^2}{n})$ as $n \rightarrow \infty$.

CLT In Action

- 1 Generate 100 (n) random numbers from an Exponential distribution (population distribution)
- 2 Compute the **sample mean** of these 100 random numbers
- 3 Repeat this process 120 times

Properties of the Multivariate Normal Distribution

- If $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then any subset of \mathbf{X} also has a multivariate normal distribution

Example: Each single variable $X_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, \dots, p$

- If $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then any linear combination of the variables has a univariate normal distribution

Example: If $Y = \mathbf{a}^T \mathbf{X}$. Then $Y \sim N(\mathbf{a}^T \boldsymbol{\mu}, \mathbf{a}^T \boldsymbol{\Sigma} \mathbf{a})$

- Any conditional distribution for a subset of the variables conditional on known values for another subset of variables is a multivariate distribution

Example:

$$\mathbf{X}_1 | \mathbf{X}_2 = \mathbf{x}_2 \sim N(\boldsymbol{\mu}_1 + \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1}(\mathbf{x}_2 - \boldsymbol{\mu}_2), \boldsymbol{\Sigma}_{11} - \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} \boldsymbol{\Sigma}_{21})$$

Example: Linear Combination of the Cholesterol Measurements [source: Penn State Univ. STAT 505]

Cholesterol levels were taken 0, 2, and 4 days following the heart attack on n patients. The mean vector is:

	Variable	Mean
$\bar{\mathbf{x}} =$	X_1 (0-day)	259.5
	X_2 (2-day)	230.8
	X_3 (4-day)	221.5

and the covariance matrix

$$\mathbf{S} = \begin{bmatrix} 2276 & 1508 & 813 \\ 1508 & 2206 & 1349 \\ 813 & 1349 & 1865 \end{bmatrix}$$

Suppose we are interested in $\Delta = X_2 - X_1$, the difference between the 2-day and the 0-day measurements. We can write the linear combination of interest as

$$\Delta = \mathbf{a}^T \mathbf{X} = \begin{bmatrix} -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix}$$

Cholesterol Measurements Example Cont'd

- The mean value for the difference Δ is

$$\begin{bmatrix} -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 259.5 \\ 230.8 \\ 221.5 \end{bmatrix} = -28.7$$

- The variance for Δ is

$$\begin{aligned} & \begin{bmatrix} -1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2276 & 1508 & 813 \\ 1508 & 2206 & 1349 \\ 813 & 1349 & 1865 \end{bmatrix} \begin{bmatrix} -1 & 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} -768 & 698 & 536 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \\ &= 1466 \end{aligned}$$


- If we assume these three variables together follows a multivariate normal distribution, then Δ follows a univariate normal distribution

Bivariate Normal Distribution

Let's focus bivariate normal distributions first as we can visualize them to facilitate our understanding. Suppose we have X_1 and X_2 jointly follows a bivariate normal distribution:

$$\begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \sim N \left[\begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix} \right]$$

Let's fix $\mu_1 = \mu_2 = 0$ and $\sigma_1^2 = \sigma_2^2 = 1$



Exponent of Multivariate Normal Distribution

Recall the multivariate normal density:

$$f(\mathbf{x}) = \frac{1}{2\pi^{\frac{p}{2}} |\Sigma|^{\frac{1}{2}}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\}.$$

This density function only depends on \mathbf{x} through the **squared Mahalanobis distance**: $(\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})$


- For bivariate normal, we get an **ellipse** whose equation is $(\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) = c^2$ which gives all $\mathbf{x} = (x_1, x_2)$ pairs with constant density
- These ellipses are call contours and all are centered around $\boldsymbol{\mu}$
- A **constant probability contour** equals

$$= \text{all } \mathbf{x} \text{ such that } (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) = c^2$$

= surface of ellipsoid centered at $\boldsymbol{\mu}$

Multivariate Normality and Outliers

The variable $d^2 = (X - \mu)^T \Sigma^{-1} (X - \mu)$ has a chi-square distribution with p degrees of freedom, i.e., $d^2 \sim \chi_p^2$ if $X \sim N(\mu, \Sigma) \Rightarrow$ we can exploit this result to check **multivariate normality** and to detect **outliers**




- Sort $(x_i - \bar{x})^T S^{-1} (x_i - \bar{x})$ in an increasing order to get **sample quantiles**
- Calculate the **theoretical quantiles** using the **chi-square quantiles** with $p = \frac{i-0.5}{n}$, $i = 1, \dots, n$
- Plot sample quantile against theoretical quantiles

Eigenvalues and Eigenvectors of Σ and the Geometry of the Multivariate Normal Density

Multivariate Normal Distribution, Copula, and Nonparametric Density Estimation

CLEMSON UNIVERSITY

Let $X \sim N(\mu, \Sigma)$, where $\mu = (10, 5)^T$ and $\Sigma = \begin{bmatrix} 64 & 16 \\ 16 & 9 \end{bmatrix}$. The 95% probability contour is shown below



Multivariate Normal Distribution

Geometry of the Multivariate Normal Density

Copula

Nonparametric Density Estimation

Next, we talk about how to “draw” this contour

- The solid ellipsoid of values x satisfy

$$(x - \mu)^T \Sigma^{-1} (x - \mu) \leq c^2 = \chi_{df=p, \alpha}^2$$

Here we have $p = 2$ and $\alpha = 0.05 \Rightarrow c = \sqrt{\chi_{2,0.05}^2} = 2.4478$

- Major axis: $\mu \pm c\sqrt{\lambda_1}e_1$, where (λ_1, e_1) is the first eigenvalue/eigenvector of Σ .

$$\Rightarrow \lambda_1 = 68.316, \quad e_1 = \begin{bmatrix} -0.9655 \\ -0.2604 \end{bmatrix}$$

- Minor axis: $\mu \pm c\sqrt{\lambda_2}e_2$, where (λ_2, e_2) is the second eigenvalue/eigenvector of Σ .

$$\Rightarrow \lambda_2 = 4.684, \quad e_2 = \begin{bmatrix} 0.2604 \\ -0.9655 \end{bmatrix}$$

Graph of 95% Probability Contour

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation


CLEMSON
UNIVERSITY

Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation



Example: Wechsler Adult Intelligence Scale [source: Penn State Univ. STAT 505]

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation



We have data (`wechslet.txt`) on 37 subjects ($n = 37$) taking the Wechsler Adult Intelligence Test, which consists four different components: 1) Information; 2) Similarities; 3) Arithmetic; 4) Picture Completion.

Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation

- 1 Calculate the sample mean vector \bar{x} and covariance matrix S
- 2 Compute the eigenvalues and eigenvectors of S and give a geometry interpretation
- 3 Diagnostic the multivariate normal assumption

Beyond Normality: Copula [Sklar, 1959; Joe, 1997]

A **copula** is a **multivariate cumulative distribution function** for which the marginal probability distribution of each variable is uniform on the interval $[0, 1]$


$$\begin{aligned} F(x_1, \dots, x_p) &= \mathbb{P}\mathbb{r}(X_1 \leq x_1, \dots, X_p \leq x_p) \\ &= \mathbb{P}\mathbb{r}(F_1^{-1}(U_1) \leq x_1, \dots, F_p^{-1}(U_p) \leq x_p) \\ &= \mathbb{P}\mathbb{r}(U_1 \leq F_1(x_1), \dots, U_p \leq F_p(x_p)) \\ &= C(F_1(x_1), \dots, F_p(x_p)) \end{aligned}$$

- Copulas are used to model the **dependence** between random variables
- Copula approach has become popular in many areas, e.g., quantitative finance as it allows for **separate modeling of marginal distributions and dependence structure**

An Illustration of Bivariate Gaussian Copula

Left: Normal marginals + Gaussian Copula ($\rho = 0.7$)


Right: Exponential marginals + Gaussian Copula ($\rho = 0.7$)



More Examples

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation

CLEMSON
UNIVERSITY



Multivariate Normal
Distribution


Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation

Old Faithful Geyser Data

Waiting time between eruptions and the duration of the eruption for the Old Faithful geyser in Yellowstone NP



Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation

CLEMSON
UNIVERSITY

Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation

Hisograms of Old Faithful Data

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation




CLEMSON
UNIVERSITY

Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation



Kernel Density Estimates of Old Faithful

Multivariate Normal
Distribution, Copula,
and Nonparametric
Density Estimation

CLEMSON
UNIVERSITY

Multivariate Normal
Distribution

Geometry of the
Multivariate Normal
Density

Copula

Nonparametric Density
Estimation

