

Lecture 7

Binomial, Hypergeometric R.V.s & Continuous Random Variables

Text: Chapter 4

STAT 8010 Statistical Methods I

January 30, 2020

Whitney Huang
Clemson University

Agenda

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



- 1 **Binomial Random Variables**
- 2 **Hypergeometric Random Variable**
- 3 **Continuous Random Variables**
- 4 **Normal Distributions**

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in n Bernoulli trials, where the probability of success in one trial is p . Let X be a Binomial r.v.

- The definition of X : # of successes in n trials of Bernoulli trials.

Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in n Bernoulli trials, where the probability of success in one trial is p . Let X be a Binomial r.v.

- The definition of X : # of successes in n trials of Bernoulli trials.
- The support: $0, 1, \dots, n$

Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in n Bernoulli trials, where the probability of success in one trial is p . Let X be a Binomial r.v.

- The definition of X : # of successes in n trials of Bernoulli trials.
- The support: $0, 1, \dots, n$
- Its parameter(s) and definition(s): p : the probability of success on 1 trial; n is the sample size

Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in n Bernoulli trials, where the probability of success in one trial is p . Let X be a Binomial r.v.

- The definition of X : **# of successes in n trials of Bernoulli trials.**
- The support: **$0, 1, \dots, n$**
- Its parameter(s) and definition(s): **p : the probability of success on 1 trial; n is the sample size**
- The probability mass function (pmf):

$$p_X(x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$$

Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in n Bernoulli trials, where the probability of success in one trial is p . Let X be a Binomial r.v.

- The definition of X : **# of successes in n trials of Bernoulli trials.**
- The support: **$0, 1, \dots, n$**
- Its parameter(s) and definition(s): **p : the probability of success on 1 trial; n is the sample size**
- The probability mass function (pmf):

$$p_X(x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$$

- The expected value:

$$\mathbb{E}[X] = np$$

Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in n Bernoulli trials, where the probability of success in one trial is p . Let X be a Binomial r.v.

- The definition of X : **# of successes in n trials of Bernoulli trials.**
- The support: **$0, 1, \dots, n$**
- Its parameter(s) and definition(s): **p : the probability of success on 1 trial; n is the sample size**
- The probability mass function (pmf):

$$p_X(x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$$

- The expected value:

$$\mathbb{E}[X] = np$$

- The variance:

$$\text{Var}(X) = np(1-p)$$

Example

To test for Extrasensory perception (ESP), we have 4 cards. They will be shuffled and one randomly selected each time, and you are to guess which card is selected. This is repeated 10 times. Suppose you do not have ESP. Let R be the number of times you guess a card correctly. What are the distribution and parameter(s) of R ? What is the expected value of R ? Furthermore, suppose that you get certified as having ESP if you score at least an 8 on the test. What is the probability that you get certified as having ESP?

Example

To test for Extrasensory perception (ESP), we have 4 cards. They will be shuffled and one randomly selected each time, and you are to guess which card is selected. This is repeated 10 times. Suppose you do not have ESP. Let R be the number of times you guess a card correctly. What are the distribution and parameter(s) of R ? What is the expected value of R ? Furthermore, suppose that you get certified as having ESP if you score at least an 8 on the test. What is the probability that you get certified as having ESP?

Solution.

Example

To test for Extrasensory perception (ESP), we have 4 cards. They will be shuffled and one randomly selected each time, and you are to guess which card is selected. This is repeated 10 times. Suppose you do not have ESP. Let R be the number of times you guess a card correctly. What are the distribution and parameter(s) of R ? What is the expected value of R ? Furthermore, suppose that you get certified as having ESP if you score at least an 8 on the test. What is the probability that you get certified as having ESP?

Solution.

$$\begin{aligned} R &\sim \text{Binomial}(n = 10, p = \frac{1}{4} = .25) \\ \mathbb{E}[R] &= n \times p = 2.5 \\ \mathbb{P}(X \geq 8) &= .000416 \end{aligned}$$

Example

Suppose that 95% of consumers can recognize Coke in a blind taste test. Assume consumers are independent of one another. The company randomly selects 4 consumers for a taste test. Let X be the number of consumers who recognize Coke.

- 1 What is the probability that X is at least 1?
- 2 What is the probability that X is at most 3?

Binomial and Hypergeometric r.v.s

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



The binomial distribution describes the probability of k successes in n trials **with replacement**.

We want a distribution to describe the probability of k successes in n trials **without replacement** from a finite population of size N containing exactly K successes.

⇒ **Hypergeometric Distribution**

Important applications are **quality control** and statistical **estimation of population proportions**. The hypergeometric r.v. is the equivalent of a Binomial r.v. except that sampling is done **without replacement**.

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions


An Example of Hypergeometric r.v.

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables

CLEMSON
UNIVERSITY

Probability:

What is the probability to get 1 red and 2 black balls?



Statistics:

What percentage of balls in the box are red?

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Let X be a hypergeometric r.v.

- The definition of X : # of successes in n trials of a random experiment, where sampling is done without replacement (or trials are dependent)

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Hypergeometric r.v.s

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Let X be a hypergeometric r.v.

- The definition of X : # of successes in n trials of a random experiment, where sampling is done without replacement (or trials are dependent)
- The support: $k \in \{\max(0, n + K - N), \dots, \min(n, K)\}$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Let X be a hypergeometric r.v.

- The definition of X : # of successes in n trials of a random experiment, where sampling is done without replacement (or trials are dependent)
- The support: $k \in \{\max(0, n + K - N), \dots, \min(n, K)\}$
- Its parameter(s) and definition(s): N : the population size, n : the sample size, and K : number of success in the population

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Let X be a hypergeometric r.v.

- The definition of X : # of successes in n trials of a random experiment, where sampling is done without replacement (or trials are dependent)
- The support: $k \in \{\max(0, n + K - N), \dots, \min(n, K)\}$
- Its parameter(s) and definition(s): N : the population size, n : the sample size, and K : number of success in the population
- The probability mass function (pmf):
$$p_X(k) = \frac{\binom{K}{k} \times \binom{N-K}{n-k}}{\binom{N}{n}}$$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Let X be a hypergeometric r.v.

- The definition of X : # of successes in n trials of a random experiment, where sampling is done without replacement (or trials are dependent)
- The support: $k \in \{\max(0, n + K - N), \dots, \min(n, K)\}$
- Its parameter(s) and definition(s): N : the population size, n : the sample size, and K : number of success in the population
- The probability mass function (pmf): $p_X(k) = \frac{\binom{K}{k} \times \binom{N-K}{n-k}}{\binom{N}{n}}$
- The expected value: $\mathbb{E}[X] = n \frac{K}{N}$

Let X be a hypergeometric r.v.

- The definition of X : # of successes in n trials of a random experiment, where sampling is done without replacement (or trials are dependent)
- The support: $k \in \{\max(0, n + K - N), \dots, \min(n, K)\}$
- Its parameter(s) and definition(s): N : the population size, n : the sample size, and K : number of success in the population
- The probability mass function (pmf): $p_X(k) = \frac{\binom{K}{k} \times \binom{N-K}{n-k}}{\binom{N}{n}}$
- The expected value: $\mathbb{E}[X] = n \frac{K}{N}$
- The variance: $\text{Var}(X) = n \frac{K}{N} \frac{N-K}{N} \frac{N-n}{N-1}$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Example

There are 100 identical looking 52" TVs at Best Buy in Anderson, SC. Let 10 of them be defective. Suppose we want to buy 8 of the aforementioned TVs (at random). What is the probability that we don't get any defective TVs?

Solution.

Example

There are 100 identical looking 52" TVs at Best Buy in Anderson, SC. Let 10 of them be defective. Suppose we want to buy 8 of the aforementioned TVs (at random). What is the probability that we don't get any defective TVs?


Solution.

Let D be the number of defective TVs in the sample.

$$D \sim \text{Hyp}(N = 100, n = 8, K = 10)$$

$$\mathbb{P}(D = 0) = \frac{\binom{10}{0} \binom{90}{8}}{\binom{100}{8}} = 0.4166$$

From Discrete to Continuous Random Variables



Probability Mass Functions vs. Probability Density Functions

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables

CLEMSON
UNIVERSITY


Binomial Random
Variables

Hypergeometric
Random Variable


Continuous Random
Variables

Normal Distributions

Pmf for Binomial($n=10, p=0.3$)



Pdf for Normal(mean=5, sd=1.5)



Remarks:

- pmf assigns probabilities to each possible values of a discrete random variable
- pdf describes the relative likelihood for a continuous random variable to take on a given interval

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^{x=b} p_X(x)$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^{x=b} p_X(x)$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^{x=b} p_X(x)$

For continuous distributions, the properties for probability density functions (Pdfs) are similar:

- $f_X(x) \geq 0$ for all possible values of x

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^{x=b} p_X(x)$

For continuous distributions, the properties for probability density functions (Pdfs) are similar:

- $f_X(x) \geq 0$ for all possible values of x
- $\int_{-\infty}^{\infty} f_X(x) dx = 1$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Probability Mass Functions v.s. Probability Density Functions cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^{x=b} p_X(x)$

For continuous distributions, the properties for probability density functions (Pdfs) are similar:

- $f_X(x) \geq 0$ for all possible values of x
- $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- $\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Cumulative Distribution Functions (cdfs) for Continuous Distribution

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

- The cdf $F_X(x)$ is defined as $F_X(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f_X(x) dx$
- we use cdf to calculate probabilities of a continuous random variable within an interval, i.e. $\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx = \int_{-\infty}^b f_X(x) dx - \int_{-\infty}^a f_X(x) dx = F_X(b) - F_X(a)$

Remark: $\mathbb{P}(X = x) = \int_x^x f_X(x) dx = 0$ for all possible values of x

Expected Value and Variance

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Expected Value and Variance

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Expected Value and Variance

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Expected Value and Variance

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[cX] = c\mathbb{E}[X]$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Expected Value and Variance

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[cX] = c\mathbb{E}[X]$
- $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Expected Value and Variance

Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[cX] = c\mathbb{E}[X]$
- $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$
- $\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \int_{-\infty}^{\infty} x^2f_X(x) dx - \left(\int_{-\infty}^{\infty} xf_X(x) dx\right)^2$

Expected Value and Variance

Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[cX] = c\mathbb{E}[X]$
- $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$
- $\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \int_{-\infty}^{\infty} x^2 f_X(x) dx - \left(\int_{-\infty}^{\infty} xf_X(x) dx\right)^2$
- $\text{Var}(cX) = c^2 \text{Var}(X)$

Expected Value and Variance

Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x xp_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[cX] = c\mathbb{E}[X]$
- $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$
- $\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \int_{-\infty}^{\infty} x^2 f_X(x) dx - \left(\int_{-\infty}^{\infty} xf_X(x) dx\right)^2$
- $\text{Var}(cX) = c^2 \text{Var}(X)$
- $\text{Var}(X - c) = \text{Var}(X)$

Example

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf
- 2 Find the expected value and variance of X

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf
- 2 Find the expected value and variance of X

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf
- 2 Find the expected value and variance of X
- 3 What is the probability that X is within .5 inches of the expected diameter?

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf
- 2 Find the expected value and variance of X
- 3 What is the probability that X is within .5 inches of the expected diameter?

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf
- 2 Find the expected value and variance of X
- 3 What is the probability that X is within .5 inches of the expected diameter?
- 4 Find $F_X(x)$

Characteristics of the Normal random variable:
Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$

Characteristics of the Normal random variable:
Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$
- Its parameter(s) and definition(s): μ : mean and σ^2 : variance

Characteristics of the Normal random variable:
Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$
- Its parameter(s) and definition(s): μ : mean and σ^2 : variance
- The probability density function (pdf): $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ for $-\infty < x < \infty$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Characteristics of the Normal random variable:
Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$
- Its parameter(s) and definition(s): μ : mean and σ^2 : variance
- The probability density function (pdf): $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ for $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value $\Phi\left(\frac{x-\mu}{\sigma}\right)$ for $-\infty < x < \infty$ from standard normal table

Characteristics of the Normal random variable:

Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$
- Its parameter(s) and definition(s): μ : mean and σ^2 : variance
- The probability density function (pdf): $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ for $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value $\Phi\left(\frac{x-\mu}{\sigma}\right)$ for $-\infty < x < \infty$ from standard normal table
- The expected value: $\mathbb{E}[X] = \mu$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Characteristics of the Normal random variable:


Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$
- Its parameter(s) and definition(s): μ : mean and σ^2 : variance
- The probability density function (pdf): $\frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\mu)^2}{2\sigma^2}}$ for $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value $\Phi\left(\frac{x-\mu}{\sigma}\right)$ for $-\infty < x < \infty$ from standard normal table
- The expected value: $\mathbb{E}[X] = \mu$
- The variance: $\text{Var}(X) = \sigma^2$

Normal Density Curves

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables

CLEMSON
UNIVERSITY



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables


Normal Distributions

- The parameter μ determines the center of the distribution

Normal Density Curves

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables

CLEMSON
UNIVERSITY



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables


Normal Distributions

- The parameter μ determines the center of the distribution
- The parameter σ^2 determines the spread of the distribution

Normal Density Curves

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables

CLEMSON
UNIVERSITY



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

- The parameter μ determines the center of the distribution
- The parameter σ^2 determines the spread of the distribution
- Also called bell-shaped distribution

Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



- Normal random variable X with mean μ and standard deviation σ can convert to standard normal Z by the following :

$$Z = \frac{X - \mu}{\sigma}$$

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



- Normal random variable X with mean μ and standard deviation σ can convert to standard normal Z by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by $\Phi(z)$, can be found from the standard normal table

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



- Normal random variable X with mean μ and standard deviation σ can convert to standard normal Z by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by $\Phi(z)$, can be found from the standard normal table

Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

- Normal random variable X with mean μ and standard deviation σ can convert to standard normal Z by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by $\Phi(z)$, can be found from the standard normal table
- The probability $\mathbb{P}(a \leq X \leq b)$ where $X \sim N(\mu, \sigma^2)$ can be compute

$$\begin{aligned}\mathbb{P}(a \leq X \leq b) &= \mathbb{P}\left(\frac{a - \mu}{\sigma} \leq Z \leq \frac{b - \mu}{\sigma}\right) \\ &= \Phi\left(\frac{b - \mu}{\sigma}\right) - \Phi\left(\frac{a - \mu}{\sigma}\right)\end{aligned}$$

Properties of Φ

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

- $\Phi(0) = .50 \Rightarrow$ Mean and Median (50_{th} percentile) for standard normal are both 0

- $\Phi(0) = .50 \Rightarrow$ Mean and Median (50_{th} percentile) for standard normal are both 0

Properties of Φ

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

- $\Phi(0) = .50 \Rightarrow$ Mean and Median (50_{th} percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$

Properties of Φ

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

- $\Phi(0) = .50 \Rightarrow$ Mean and Median (50_{th} percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$

- $\Phi(0) = .50 \Rightarrow$ Mean and Median (50_{th} percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$
- $\mathbb{P}(Z > z) = 1 - \Phi(z) = \Phi(-z)$

Example

Let us examine Z . Find the following probabilities with respect to Z :

1 Z is at most -1.75 ▶

2 Z is between -2 and 2 inclusive ▶

3 Z is less than $.5$ ▶

Example Cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Solution.

$$\textcircled{1} \quad \mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401 \quad \textcircled{\leftarrow}$$

Example Cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Solution.

$$1 \quad \mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401 \quad \leftarrow$$

$$2 \quad \mathbb{P}(-2 \leq Z \leq 2) = \Phi(2) - \Phi(-2) = .9772 - .0228 = .9544 \quad \leftarrow$$

Example Cont'd

Solution.

$$1 \quad \mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401$$

$$2 \quad \mathbb{P}(-2 \leq Z \leq 2) = \Phi(2) - \Phi(-2) = .9772 - .0228 = .9544$$

$$3 \quad \mathbb{P}(Z < .5) = \Phi(.5) = .6915$$

Sums of Normal Random Variables

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

If X_i $1 \leq i \leq n$ are independent normal random variables with mean μ_i and variance σ_i^2 , respectively.

Sums of Normal Random Variables

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

If X_i $1 \leq i \leq n$ are independent normal random variables with mean μ_i and variance σ_i^2 , respectively.

- Let $S_n = \sum_{i=1}^n X_i$ then $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$

Sums of Normal Random Variables

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

If X_i $1 \leq i \leq n$ are independent normal random variables with mean μ_i and variance σ_i^2 , respectively.

- Let $S_n = \sum_{i=1}^n X_i$ then $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$

Sums of Normal Random Variables

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables


Normal Distributions


If X_i $1 \leq i \leq n$ are independent normal random variables with mean μ_i and variance σ_i^2 , respectively.


- Let $S_n = \sum_{i=1}^n X_i$ then $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$
- This can be applied for any integer n

Example

Let X_1 , X_2 , and X_3 be mutually independent, Normal random variables. Let their means and standard deviations be $3k$ and k for $k = 1, 2$, and 3 respectively. Find the following distributions:

1 $\sum_{i=1}^3 X_i$ 

2 $X_1 + 2X_2 - 3X_3$ 

3 $X_1 + 5X_3$ 

Example Cont'd

Binomial,
Hypergeometric
R.V.s & Continuous
Random Variables



Binomial Random
Variables

Hypergeometric
Random Variable

Continuous Random
Variables

Normal Distributions

Solution.

1 $\sum_{i=1}^3 X_i \sim N(\mu = 3 + 6 + 9 = 15, \sigma^2 = 1^2 + 2^2 + 3^2 = 14)$ ◀

2 $X_1 + 2X_2 - 3X_3 \sim N(\mu = 3 + 12 - 27 = -12, \sigma^2 = 1^2 + 4 \times 2^2 + 9 \times 3^2 = 98)$ ◀

3 $X_1 + 5X_3 \sim N(\mu = 3 + 45 = 48, \sigma^2 = 1^2 + 25 \times 3^2 = 226)$ ◀