

# Lecture 9

## Normal Approximation to Binomial, Sampling Distribution, and Central Limit Theorem

Text: Chapter 4

*STAT 8010 Statistical Methods I*  
February 6, 2020

Whitney Huang  
Clemson University

# Agenda

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

## 1 Normal approximation of Binomial Distribution

## 2 Sampling Distribution

## 3 Central Limit Theorem (CLT)

# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$

# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$

# Normal approximation of Binomial Distribution

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$
- If  $X \sim \text{Bin}(n, p)$  with  $np > 5$  and  $n(1 - p) > 5$  then we can use  $X^* \sim N(\mu = np, \sigma^2 = np(1 - p))$  to approximate  $X$

# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$
- If  $X \sim \text{Bin}(n, p)$  with  $np > 5$  and  $n(1 - p) > 5$  then we can use  $X^* \sim N(\mu = np, \sigma^2 = np(1 - p))$  to approximate  $X$



# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$
- If  $X \sim \text{Bin}(n, p)$  with  $np > 5$  and  $n(1 - p) > 5$  then we can use  $X^* \sim N(\mu = np, \sigma^2 = np(1 - p))$  to approximate  $X$
- Notice that Binomial is a discrete distribution but normal is a continuous distribution so that  $\mathbb{P}(X^* = x) = 0 \forall x$

# Normal approximation of Binomial Distribution

- We can use a Normal Distribution to approximate a Binomial Distribution if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$
- If  $X \sim \text{Bin}(n, p)$  with  $np > 5$  and  $n(1 - p) > 5$  then we can use  $X^* \sim N(\mu = np, \sigma^2 = np(1 - p))$  to approximate  $X$
- Notice that Binomial is a discrete distribution but normal is a continuous distribution so that  $\mathbb{P}(X^* = x) = 0 \forall x$

# Normal approximation of Binomial Distribution

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

- We can use a **Normal Distribution** to approximate a **Binomial Distribution** if  $n$  is large
- Rule of thumb for this approximation to be valid (in this class) is  $np > 5$  and  $n(1 - p) > 5$
- If  $X \sim \text{Bin}(n, p)$  with  $np > 5$  and  $n(1 - p) > 5$  then we can use  $X^* \sim N(\mu = np, \sigma^2 = np(1 - p))$  to approximate  $X$
- Notice that Binomial is a **discrete** distribution but normal is a **continuous** distribution so that  $\mathbb{P}(X^* = x) = 0 \ \forall x$
- **Continuity correction:** we use  $\mathbb{P}(x - 0.5 \leq X^* \leq x + 0.5)$  to approximate  $\mathbb{P}(X = x)$

## Example

Suppose a class has 400 students (to begin with), that each student drops independently of any other student with a probability of .07. Let  $X$  be the number of students that finish this course

- 1 Find the probability that  $X$  is between 370 and 373 inclusive
- 2 Is an approximation appropriate for the number of students that finish the course?
- 3 If so, what is this distribution and what are the parameter(s)?
- 4 Find the probability that is between 370 and 373 inclusive by using the approximation

# Sampling Distribution

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**

Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

# Sampling Distribution

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**

Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

# Sampling Distribution

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**
- A **statistic** is a function of a **random sample**

**Example:**

- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**
- A **statistic** is a function of a **random sample**

## Example:

- Sample mean:  $\bar{X}_n = \sum_{i=1}^n X_i / n$



- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**
- A **statistic** is a function of a **random sample**

## Example:

- Sample mean:  $\bar{X}_n = \sum_{i=1}^n X_i / n$
- Sample variance:  $\sum_{i=1}^n (X_i - \bar{X}_n)^2 / (n - 1)$

- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**
- A **statistic** is a function of a **random sample**

## Example:

- Sample mean:  $\bar{X}_n = \sum_{i=1}^n X_i / n$
- Sample variance:  $\sum_{i=1}^n (X_i - \bar{X}_n)^2 / (n - 1)$
- Sample maximum:  $\max_{i=1}^n X_i$

- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**
- A **statistic** is a function of a **random sample**

## Example:

- Sample mean:  $\bar{X}_n = \sum_{i=1}^n X_i / n$
- Sample variance:  $\sum_{i=1}^n (X_i - \bar{X}_n)^2 / (n - 1)$
- Sample maximum:  $\max_{i=1}^n X_i$

- Independent random variables  $X_1, X_2, \dots, X_n$  with the same distribution are called a **random sample**
- A **statistic** is a function of a **random sample**

## Example:

- Sample mean:  $\bar{X}_n = \sum_{i=1}^n X_i / n$
  - Sample variance:  $\sum_{i=1}^n (X_i - \bar{X}_n)^2 / (n - 1)$
  - Sample maximum:  $\max_{i=1}^n X_i$
- 
- The probability distribution of a statistic is called its **sampling distribution**

## Example

Suppose  $X_1, X_2, \dots, X_n$  is a random sample from a  $N(\mu, \sigma^2)$  population, Find the sampling distribution of sample mean.

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

## Example

Suppose  $X_1, X_2, \dots, X_n$  is a random sample from a  $N(\mu, \sigma^2)$  population, Find the sampling distribution of sample mean.

$\bar{X}_n = \frac{\sum_{i=1}^n X_i}{n} = \sum_{i=1}^n \frac{1}{n} X_i$ . From last lecture we know that sum of normal r.v.s is still a normal r.v. Hence we only need to figure its mean and variance.

$\mathbb{E}[\bar{X}_n] = \sum_{i=1}^n \mu/n = \mu$ .  $\text{Var}[\bar{X}_n] = \sum_{i=1}^n \frac{1}{n^2} \sigma^2 = \frac{\sigma^2}{n}$ . Therefore, we have  $\bar{X}_n \sim N(\mu, \frac{\sigma^2}{n})$

# Central Limit Theorem (CLT)

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



## CLT

The **sampling distribution** of the **mean** will become approximately **normally distributed** as the **sample size becomes larger**, **irrespective of the shape of the population distribution!**

Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

Let  $X_1, X_2, \dots, X_n \stackrel{i.i.d.}{\sim} F$  with  $\mu = \mathbb{E}[X_i]$  and  $\sigma^2 = \text{Var}[X_i]$ .  
Then  $\bar{X}_n = \frac{\sum_{i=1}^n X_i}{n} \xrightarrow{d} N(\mu, \frac{\sigma^2}{n})$  as  $n \rightarrow \infty$ .

## CLT In Action

- 1 Generate 100 ( $n$ ) random numbers from an Exponential distribution (population distribution)
- 2 Compute the **sample mean** of these 100 random numbers
- 3 Repeat this process 120 times

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)



# CLT: Sample Size ( $n$ ) and the Normal Approximation







Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)



# Why CLT is important?

Normal  
Approximation to  
Binomial, Sampling  
Distribution, and  
Central Limit  
Theorem



Normal approximation  
of Binomial Distribution

Sampling Distribution

Central Limit Theorem  
(CLT)

- In many cases, we would like to make statistical inference about the population mean  $\mu$ 
  - The sample mean  $\bar{X}_n$  is a sensible estimator for the population mean
  - CLT tells us the **distribution** of our estimator  
 $\Rightarrow \bar{X}_n \approx N(\mu, \frac{\sigma^2}{n})$
- Applications: Hypothesis testing, confidence interval