

Lecture 1

Overview

DSA 8070 Multivariate Analysis
August 18-20, 2021

Introduction

Objectives of
Multivariate Analysis

Useful Tools for
Multivariate Analysis

Whitney Huang
Clemson University

Introduction

Objectives of
Multivariate Analysis

Useful Tools for
Multivariate Analysis

- 1 Introduction
- 2 Objectives of Multivariate Analysis
- 3 Useful Tools for Multivariate Analysis

- In many observational or experimental studies, observations are collected simultaneously on **more than one variable** on each unit

```
> head(Boston)
```

	crim	zn	indus	chas	nox	rm	age	dis	rad	tax	ptratio	black	lstat	medv
1	0.00632	18	2.31	0	0.538	6.575	65.2	4.0900	1	296	15.3	396.90	4.98	24.0
2	0.02731	0	7.07	0	0.469	6.421	78.9	4.9671	2	242	17.8	396.90	9.14	21.6
3	0.02729	0	7.07	0	0.469	7.185	61.1	4.9671	2	242	17.8	392.83	4.03	34.7
4	0.03237	0	2.18	0	0.458	6.998	45.8	6.0622	3	222	18.7	394.63	2.94	33.4
5	0.06905	0	2.18	0	0.458	7.147	54.2	6.0622	3	222	18.7	396.90	5.33	36.2
6	0.02985	0	2.18	0	0.458	6.430	58.7	6.0622	3	222	18.7	394.12	5.21	28.7


- **Multivariate analysis** is the collection of statistical methods that can be used to analyze these multiple measurements
⇒ *some are extensions of familiar methods (t-test, ANOVA, linear regression,...) while others are unique to multivariate analysis*
- Idea is to exploit potential **“correlations”** among the multiple measurements to improve inference (see an example in the next slide)

Using Multivariate Methods Could Lead to Sharper Inference

Introduction

Objectives of
Multivariate Analysis

Useful Tools for
Multivariate Analysis



Source: Fig. 1.1 of Applied Multivariate Statistics with R by Zeltermann

Introduction

Objectives of
Multivariate Analysis

Useful Tools for
Multivariate Analysis

Objectives of Multivariate Analysis

Dimensionality Reduction or Structural Simplification

- **Goal:** to reduce the “dimensionality” by considering a small number of (linear) combinations of a large number of measurements without losing important information
- **Examples:**
 - A single index of patient reaction to radiotherapy can be constructed from measurements on several response variables
 - Wildlife ecologists can construct a few indices of habitat preference from measurements of dozens of features of nesting sites selected by a certain bird species
- **Techniques:**
 - **Principal Component Analysis** (Week 9)
 - **Factor Analysis** (Week 10)
 - **Multidimensional Scaling** (Week 14)

- **Goal:** to **identify** groups of “similar” units or to **classify** units into previously defined groups
- **Examples:**
 - The US IRS uses data collected from tax returns (income, amount withheld, deductions, ...) to **classify** taxpayers into two groups: those who will be audited and those who will not
 - Using the concentration of elements (copper, silver, tin, antimony) in the lead alloy used in bullets, the FBI **identifies** ‘similar’ bullets that may be used to infer whether bullets were produced from the same batch of lead
- **Techniques:**
 - **Classification Analysis** (Week 12)
 - **Cluster Analysis** (Week 13)

Investigation of the Dependence among Variables and Prediction

- **Goal:** to estimate the relationship among variables and, if the variables are associated, to predict the value of some of them given information on the others
- **Examples:**
 - The associations between measures of risk-taking propensity and measures of socioeconomic characteristics for top-level business executives were used to assess the relation between risk-taking behavior and performance
 - The association between test scores, and several college performance variables were used to develop predictors of success in college
- **Techniques:**
 - **Multivariate Regression** (Week 7)
 - **Canonical Correlation Analysis** (Week 11)

- **Goal:** to test if differences in sets of response mean vectors for two or more groups large enough to be distinguished from sampling variation
- **Examples:**
 - A transportation company wants to know if means for gasoline mileage, repair costs, downtime due to repairs differ for different truck models
 - An insurance company wants to know if changing case management practices leads to changes in mean length of hospital stay, mean infection rates, and mean costs
- **Techniques:**
 - **Hotelling's T^2 and MAVONA** (Week 5 and Week 6)

Introduction


Objectives of
Multivariate Analysis

Useful Tools for
Multivariate Analysis

Useful Concepts/Tools for Multivariate Analysis

Statistical Distance

Multivariate methods rely on “distances” between data points: **clustering** (group units that are “close”); **classification** (allocate each unit to the “closest” group)



Question: which one ((x_0, y_0) or (x_1, y_1)) is closer the center of the observations?

The study of multivariate methods is greatly facilitated by the use of matrix algebra

- Many operations performed on multivariate data are presented using vector/matrix notation, e.g., $\mathbf{X}_{n \times p}$ (Data matrix); $\hat{\boldsymbol{\mu}}_{p \times 1}$ (estimated mean vector); $\hat{\boldsymbol{\Sigma}}_{p \times p}$ (estimated covariance matrix)
- The computation of **eigenvalues** and **eigenvectors** (i.e., the **spectral decomposition**) plays an important role in multivariate analysis
- We will use R to perform the needed matrix operations

- We will (almost always) assume the joint distribution of $\mathbf{X} = (X_1, X_2, \dots, X_p)^T$ follows a multivariate normal distribution with the probability density function:

$$f(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{\frac{d}{2}} \det(\boldsymbol{\Sigma})^{\frac{1}{2}}} \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right]$$

- The multivariate normal assumption is often appropriate:
 - Variables can sometimes be assumed to be multivariate normal (perhaps after transformation)
 - **Central limit theorem** tells us that distribution of many **multivariate sample statistics** is approximately normal, regardless of the form of the population distribution