

# Lecture 9

## Law of Total Probability & Bayes' Rule

Text: Chapter 4

STAT 8010 Statistical Methods I  
September 9, 2019

Whitney Huang  
Clemson University



Notes

---

---

---

---

---

---

---

### Agenda

- 1 Law of Total Probability
- 2 Bayes' Rule
- 3 Random Variables



Notes

---

---

---

---

---

---

---

### Law of Partitions & Multiplication Rule

#### Law of partitions

Let  $A_1, A_2, \dots, A_k$  form a partition of  $\Omega$ . Then, for all events  $B$ ,

$$\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(A_i \cap B)$$

#### Multiplication rule

- 2 events:

$$\mathbb{P}(B \cap A) = \mathbb{P}(A) \times \mathbb{P}(B|A) = \mathbb{P}(B) \times \mathbb{P}(A|B)$$

- More than 2 events:

$$\begin{aligned} \mathbb{P}(\cap_{i=1}^n A_i) &= \mathbb{P}(A_1) \times \mathbb{P}(A_2|A_1) \times \mathbb{P}(A_3|A_1 \cap A_2) \\ &\quad \times \dots \times \mathbb{P}(A_n|A_{n-1} \cap \dots \cap A_1) \end{aligned}$$



Notes

---

---

---

---

---

---

---

Law of Total Probability

Let  $A_1, A_2, \dots, A_k$  form a partition of  $\Omega$ . Then, for all events  $B$ ,

$$\begin{aligned} \mathbb{P}(B) &= \sum_{i=1}^k \underbrace{\mathbb{P}(A_i \cap B)}_{\text{Law of partitions}} \\ &= \sum_{i=1}^k \underbrace{\mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}_{\text{Multiplication rule}} \end{aligned}$$

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

94

Notes

---

---

---

---

---

---

---

Motivating example

The Monty Hall Problem

There was an old television show called Let's Make a Deal, whose original host was named Monty Hall. The set-up is as follows. You are on a game show and you are given the choice of three doors. Behind one door is a car, behind the others are goats. You pick a door, and the host, who knows what is behind the doors, opens another door (not your pick) which has a goat behind it. Then he asks you if you want to change your original pick. The question we ask you is, "Is it to your advantage to switch your choice?"

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

95

Notes

---

---

---

---

---

---

---

The Monty Hall Problem



Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

96

Notes

---

---

---

---

---

---

---

The Monty Hall Problem Solution

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.7

Notes

---

---

---

---

---

---

---

Bayes' Rule

General form

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

Let  $A_1, A_2, \dots, A_k$  form a partition of the sample space.  
Then for every event  $B$  in the sample space,

$$\mathbb{P}(A_j|B) = \frac{\mathbb{P}(B|A_j) \times \mathbb{P}(A_j)}{\sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)}, \quad j = 1, 2, \dots, k$$

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.8

Notes

---

---

---

---

---

---

---

Example

Let us assume that a specific disease is only present in 5 out of every 1,000 people. Suppose that the test for the disease is accurate 99% of the time a person has the disease and 95% of the time that a person lacks the disease. What is the probability that the person has the disease given that they tested positive?

Solution.

$$\mathbb{P}(D|+) = \frac{\mathbb{P}(D \cap +)}{\mathbb{P}(+)} = \frac{.005 \times .99}{.005 \times .99 + .995 \times .05} = \frac{.00495}{.0547} = .0905$$

The reason we get such a surprising result is because the disease is so rare that the number of false positives greatly outnumbers the people who truly have the disease.

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.9

Notes

---

---

---

---

---

---

---

Review of Probability (we learned so far)

Basic Concepts:

- Random Experiment, Sample Space, Outcome, Event
- Frequentist Interpretation of Probability and Equally Likely Framework
- Union and Intersection
- Mutually Exclusive, Exhaustive, Partition
- Venn Diagram

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.10

Notes

---

---

---

---

---

---

---

Review: Probability Rules

- $0 \leq \mathbb{P}(A) \leq 1$  for any event  $A$ ,  $\mathbb{P}(\emptyset) = 0$ ,  $\mathbb{P}(\Omega) = 1$
- Complement rule:  $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$
- General addition rule:  
 $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- Multiplication rule:  
 $\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \times \mathbb{P}(B) = \mathbb{P}(B|A) \times \mathbb{P}(A)$
- Conditional probability:  $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$
- Law of total probability:  
 $\mathbb{P}(B) = \sum_{i=1}^k \mathbb{P}(B \cap A_i) = \sum_{i=1}^k \mathbb{P}(B|A_i) \times \mathbb{P}(A_i)$
- Independence: if  $A$  and  $B$  are independent, then  $\mathbb{P}(A|B) = \mathbb{P}(A)$ ,  $\mathbb{P}(B|A) = \mathbb{P}(B)$ , and  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.11

Notes

---

---

---

---

---

---


---

Random Variables

A **random variable** is a real-valued function whose domain is the sample space of a random experiment. In other words, a random variable is a function

$$X : \Omega \mapsto \mathbb{R}$$

where  $\Omega$  is the sample space of the random experiment under consideration and  $\mathbb{R}$  represents the set of all real numbers.



Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.12

Notes

---

---

---

---

---

---

---

Example

The following is a chart describing the number of siblings each student in a particular class has.

Siblings ( $X$ )	Frequency	Relative Frequency
0	8	.200
1	17	.425
2	11	.275
3	3	.075
4	1	.025
Total	40	1

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.13

Notes

---

---

---

---

---

---

---

Example cont'd

Let's define the event  $A$  as the event that a randomly chosen student has 2 or more siblings. What is  $\mathbb{P}(X \in A)$ ?

**Solution.**

$$\begin{aligned}\mathbb{P}(X \in A) &= \mathbb{P}(X \geq 2) = \mathbb{P}(X = 2) + \mathbb{P}(X = 3) + \mathbb{P}(X = 4) \\ &= .275 + 0.075 + 0.025 = .375\end{aligned}$$

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.14

Notes

---

---

---

---

---

---

---

Types of Random Variables

There are two main types of quantitative random variables: **discrete** and **continuous**. A discrete random variable often involves a count of something. Examples may include number of cars per household, etc.

**Discrete random variable**

A random variable  $X$  is called a discrete random variable if the outcome of the random variable is limited to a countable set of real numbers (usually integers).

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.15

Notes

---

---

---

---

---

---

---

Probability Mass Function

Let  $X$  be a discrete random variable. Then the probability mass function (pmf) of  $X$  is the real-valued function defined on  $\mathbb{R}$  by

$$p_X(x) = \mathbb{P}(X = x)$$

The capital letter,  $X$ , is used to denote random variable. Lowercase letter,  $x$ , is used to denote possible values of the random variable.

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.16

Notes

---

---

---

---

---

---

---

Example

Flip a fair coin 3 times. Let  $X$  denote the number of heads tossed in the 3 flips. Create a pmf for  $X$

Solution.

$X$	0	1	2	3
$p_X(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.17

Notes

---

---

---

---

---

---

---

Properties of a PMF

- $0 \leq p_X(x) \leq 1, \forall x \in R$
- $\{x \in \mathbb{R} : p_X(x) \neq 0\}$  is countable
- $\sum_x p_X(x) = 1$

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability

Bayes' Rule

Random Variables

9.18

Notes

---

---

---

---

---

---

---

Example

Let  $X$  be a random variable with pmf defined as follows:

$$p_X(x) = \begin{cases} k(5 - x) & \text{if } x = 0, 1, 2, 3, 4 \\ 0 & \text{otherwise} \end{cases}$$

- 1 Find the value of  $k$  that makes  $p_X(x)$  a legitimate pmf.
- 2 What is the probability that  $X$  is between 1 and 3 inclusive?
- 3 If  $X$  is not 0, what is the probability that  $X$  is less than 3?

Law of Total Probability & Bayes' Rule

CLEMSON UNIVERSITY

Law of Total Probability  
Bayes' Rule  
Random Variables

9/19

Notes

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---