

# Lecture 7

## Binomial, Hypergeometric R.V.s & Continuous Random Variables

Text: Chapter 4

STAT 8010 Statistical Methods I  
January 30, 2020

Whitney Huang  
Clemson University



### Notes

---

---

---

---

---

---

---

### Agenda

- 1 Binomial Random Variables
- 2 Hypergeometric Random Variable
- 3 Continuous Random Variables
- 4 Normal Distributions



### Notes

---

---

---

---

---

---

---

### Binomial Random Variable

We define the **Binomial** r.v. as the number of successes in  $n$  Bernoulli trials, where the probability of success in one trial is  $p$ . Let  $X$  be a Binomial r.v.

- The definition of  $X$ : **# of successes in  $n$  trials of Bernoulli trials.**
- The support:  $0, 1, \dots, n$
- Its parameter(s) and definition(s):  **$p$ : the probability of success on 1 trial;  $n$  is the sample size**
- The probability mass function (pmf):

$$p_X(x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n$$

- The expected value:

$$\mathbb{E}[X] = np$$

- The variance:

$$\text{Var}(X) = np(1-p)$$



### Notes

---

---

---

---

---

---

---

### Example

To test for Extrasensory perception (ESP), we have 4 cards. They will be shuffled and one randomly selected each time, and you are to guess which card is selected. This is repeated 10 times. Suppose you do not have ESP. Let  $R$  be the number of times you guess a card correctly. What are the distribution and parameter(s) of  $R$ ? What is the expected value of  $R$ ? Furthermore, suppose that you get certified as having ESP if you score at least an 8 on the test. What is the probability that you get certified as having ESP?

**Solution.**

$$R \sim \text{Binomial}(n = 10, p = \frac{1}{4} = .25)$$

$$\mathbb{E}[R] = n \times p = 2.5$$

$$\mathbb{P}(X \geq 8) = .000416$$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

74

Notes

---

---

---

---

---

---

---

---

### Example

Suppose that 95% of consumers can recognize Coke in a blind taste test. Assume consumers are independent of one another. The company randomly selects 4 consumers for a taste test. Let  $X$  be the number of consumers who recognize Coke.

❶ What is the probability that  $X$  is at least 1?

❷ What is the probability that  $X$  is at most 3?

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

75

Notes

---

---

---

---

---

---

---

---

### Binomial and Hypergeometric r.v.s

The binomial distribution describes the probability of  $k$  successes in  $n$  trials **with replacement**.

We want a distribution to describe the probability of  $k$  successes in  $n$  trials **without replacement** from a finite population of size  $N$  containing exactly  $K$  successes.

⇒ **Hypergeometric Distribution**

Important applications are **quality control** and statistical **estimation of population proportions**. The hypergeometric r.v. is the equivalent of a Binomial r.v. except that sampling is done **without replacement**.

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

76

Notes

---

---

---

---


---

---

---

---

## An Example of Hypergeometric r.v.



Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.7

## Notes

## Hypergeometric r.v.s

Let  $X$  be a hypergeometric r.v.

- The definition of  $X$ : # of successes in  $n$  trials of a random experiment, where sampling is done without replacement (or trials are dependent)
- The support:  $k \in \{\max(0, n + K - N), \dots, \min(n, K)\}$
- Its parameter(s) and definition(s):  $N$ : the population size,  $n$ : the sample size, and  $K$ : number of success in the population
- The probability mass function (pmf):  

$$p_X(k) = \frac{\binom{K}{k} \times \binom{N-K}{n-k}}{\binom{N}{n}}$$
- The expected value:  $\mathbb{E}[X] = n \frac{K}{N}$
- The variance:  $\text{Var}(X) = n \frac{K}{N} \frac{N-K}{N} \frac{N-n}{N-1}$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.8

## Notes

## Example

There are 100 identical looking 52" TVs at Best Buy in Anderson, SC. Let 10 of them be defective. Suppose we want to buy 8 of the aforementioned TVs (at random). What is the probability that we don't get any defective TVs?

### Solution.

Let  $D$  be the number of defective TVs in the sample.

$$D \sim \text{Hyp}(N = 100, n = 8, K = 10)$$

$$\mathbb{P}(D = 0) = \frac{\binom{10}{0} \binom{90}{8}}{\binom{100}{8}} = 0.4166$$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable


Continuous  
Random Variables

Normal  
Distributions

7.9

## Notes

# From Discrete to Continuous Random Variables



Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.10

Notes

---

---

---

---


---

---

---

---

## Probability Mass Functions vs. Probability Density Functions



Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.11

Notes

---

---

---

---

---

---

---

---

### Remarks:

- pmf assigns probabilities to each possible values of a discrete random variable
- pdf describes the relative likelihood for a continuous random variable to take on a given interval

## Probability Mass Functions v.s. Probability Density Functions cont'd

Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$  for all possible values of  $x$
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^b p_X(x)$

For continuous distributions, the properties for probability density functions (Pdfs) are similar:

- $f_X(x) \geq 0$  for all possible values of  $x$
- $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- $\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.12

Notes

---

---

---

---

---

---

---

---

Cumulative Distribution Functions (cdfs) for Continuous r.v.s

- The cdf  $F_X(x)$  is defined as
$$F_X(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f_X(x) \, dx$$
  - we use cdf to calculate probabilities of a continuous random variable within an interval, i.e.
$$\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) \, dx = \int_{-\infty}^b f_X(x) \, dx - \int_{-\infty}^a f_X(x) \, dx = F_X(b) - F_X(a)$$
- Remark:**  $\mathbb{P}(X = x) = \int_x^x f_X(x) \, dx = 0$  for all possible values of  $x$

Binomial, Hypergeometric R.V.s & Continuous Random Variables

CLEMSON UNIVERSITY

Binomial Random Variables

Hypergeometric Random Variable

Continuous Random Variables

Normal Distributions

7.13

Notes

---

---

---

---

---

---

---

Expected Value and Variance

- Recall the expected value formula for the discrete random variable:  $\mathbb{E}[X] = \sum_x xp_X(x)$
- For continuous random variables, we have similar formulas:
- Let  $a$ ,  $b$ , and  $c$  are constant real numbers
- $\mathbb{E}[X] = \int_{-\infty}^{\infty} xf_X(x) \, dx$
  - $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f_X(x) \, dx$
  - $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
  - $\mathbb{E}[cX] = c\mathbb{E}[X]$
  - $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$
  - $\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \int_{-\infty}^{\infty} x^2f_X(x) \, dx - \left(\int_{-\infty}^{\infty} xf_X(x) \, dx\right)^2$
  - $\text{Var}(cX) = c^2\text{Var}(X)$
  - $\text{Var}(X - c) = \text{Var}(X)$

Binomial, Hypergeometric R.V.s & Continuous Random Variables

CLEMSON UNIVERSITY

Binomial Random Variables

Hypergeometric Random Variable

Continuous Random Variables

Normal Distributions

7.14

Notes

---

---

---

---

---

---

---

Example

- Let  $X$  represent the diameter in inches of a circular disk cut by a machine. Let  $f_X(x) = c(4x - x^2)$  for  $1 \leq x \leq 4$  and be 0 otherwise. Answer the following questions:
- Find the value of  $c$  that makes this a valid pdf
  - Find the expected value and variance of  $X$
  - What is the probability that  $X$  is within .5 inches of the expected diameter?
  - Find  $F_X(x)$

Binomial, Hypergeometric R.V.s & Continuous Random Variables

CLEMSON UNIVERSITY

Binomial Random Variables

Hypergeometric Random Variable

Continuous Random Variables

Normal Distributions

7.15

Notes

---

---

---

---

---

---

---

## Normal Distribution

Characteristics of the Normal random variable:

Let  $X$  be a Normal r.v.

- The support for  $X$ :  $(-\infty, \infty)$
- Its parameter(s) and definition(s):  $\mu$  : mean and  $\sigma^2$  : variance
- The probability density function (pdf):  $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for  $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value  $\Phi(\frac{x-\mu}{\sigma})$  for  $-\infty < x < \infty$  from standard normal table
- The expected value:  $\mathbb{E}[X] = \mu$
- The variance:  $\text{Var}(X) = \sigma^2$



## Notes

---

---

---

---


---

---


---

---

## Normal Density Curves



- The parameter  $\mu$  determines the center of the distribution
- The parameter  $\sigma^2$  determines the spread of the distribution
- Also called bell-shaped distribution



## Notes

---

---

---

---

---

---

---

---


## Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

- Normal random variable  $X$  with mean  $\mu$  and standard deviation  $\sigma$  can convert to standard normal  $Z$  by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by  $\Phi(z)$ , can be found from the standard normal table
- The probability  $\mathbb{P}(a \leq X \leq b)$  where  $X \sim N(\mu, \sigma^2)$  can be compute

$$\begin{aligned} \mathbb{P}(a \leq X \leq b) &= \mathbb{P}\left(\frac{a - \mu}{\sigma} \leq Z \leq \frac{b - \mu}{\sigma}\right) \\ &= \Phi\left(\frac{b - \mu}{\sigma}\right) - \Phi\left(\frac{a - \mu}{\sigma}\right) \end{aligned}$$



## Notes

---

---

---

---

---

---

---

---

Properties of  $\Phi$

- $\Phi(0) = .50 \Rightarrow$  Mean and Median ( $50_{th}$  percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$
- $\mathbb{P}(Z > z) = 1 - \Phi(z) = \Phi(-z)$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.19

Notes

---

---

---

---

---

---

---

Example

Let us examine  $Z$ . Find the following probabilities with respect to  $Z$ :

- 1  $Z$  is at most  $-1.75$
- 2  $Z$  is between  $-2$  and  $2$  inclusive
- 3  $Z$  is less than  $.5$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.20

Notes

---

---

---

---

---

---

---

Example Cont'd

Solution.

- 1  $\mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401$
- 2  $\mathbb{P}(-2 \leq Z \leq 2) = \Phi(2) - \Phi(-2) = .9772 - .0228 = .9544$
- 3  $\mathbb{P}(Z < .5) = \Phi(.5) = .6915$

Binomial,  
Hypergeometric  
R.V.s &  
Continuous  
Random  
Variables

CLEMSON  
UNIVERSITY

Binomial Random  
Variables

Hypergeometric  
Random Variable

Continuous  
Random Variables

Normal  
Distributions

7.21

Notes

---

---

---

---

---

---

---

Sums of Normal Random Variables

If  $X_i$ ,  $1 \leq i \leq n$  are independent normal random variables with mean  $\mu_i$  and variance  $\sigma_i^2$ , respectively.

- Let  $S_n = \sum_{i=1}^n X_i$  then  $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$
- This can be applied for any integer  $n$

Binomial, Hypergeometric R.V.s & Continuous Random Variables

CLEMSON UNIVERSITY

Binomial Random Variables

Hypergeometric Random Variable

Continuous Random Variables

Normal Distributions

7.22

Notes

---

---

---

---

---

---

---

Example

Let  $X_1$ ,  $X_2$ , and  $X_3$  be mutually independent, Normal random variables. Let their means and standard deviations be  $3k$  and  $k$  for  $k = 1, 2$ , and  $3$  respectively. Find the following distributions:

- $\sum_{i=1}^3 X_i$
- $X_1 + 2X_2 - 3X_3$
- $X_1 + 5X_3$

Binomial, Hypergeometric R.V.s & Continuous Random Variables

CLEMSON UNIVERSITY

Binomial Random Variables

Hypergeometric Random Variable

Continuous Random Variables

Normal Distributions

7.23

Notes

---

---

---

---

---

---

---

Example Cont'd

Solution.

- $\sum_{i=1}^3 X_i \sim N(\mu = 3 + 6 + 9 = 15, \sigma^2 = 1^2 + 2^2 + 3^2 = 14)$
- $X_1 + 2X_2 - 3X_3 \sim N(\mu = 3 + 12 - 27 = -12, \sigma^2 = 1^2 + 4 \times 2^2 + 9 \times 3^2 = 98)$
- $X_1 + 5X_3 \sim N(\mu = 3 + 45 = 48, \sigma^2 = 1^2 + 25 \times 3^2 = 226)$

Binomial, Hypergeometric R.V.s & Continuous Random Variables

CLEMSON UNIVERSITY

Binomial Random Variables

Hypergeometric Random Variable

Continuous Random Variables

Normal Distributions

7.24

Notes

---

---

---

---

---

---

---