

# Lecture 14

## Multidimensional Scaling

Reading: Izenman Chapter 13

The main reference for these slides is from Dr. Markus Kalisch's Lecture Notes at

<https://stat.ethz.ch/education/semesters/ss2012/ams/slides/v4.1.pdf>

DSA 8070 Multivariate Analysis

Whitney Huang  
Clemson University

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.1

Notes

---

---

---

---

---

---

---

### Agenda

1 Main Idea

2 Classical Multidimensional Scaling

3 Non-metric Multidimensional Scaling

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.2

Notes

---

---

---

---


---

---

---

### Basic Idea of Multidimensional Scaling (MDS)

- Represent high-dimensional point cloud in low (usually 2) dimensional Euclidean space while *preserving as well as possible the inter-point distances*
- Classical/Metric MDS: Use a clever projection
- Non-metric MDS: Squeeze data on table



Source: Dr. Markus Kalisch's Lecture Notes on MDS

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.3

Notes

---

---

---

---

---

---

---

Classical MDS (cMDS)

- **Goal:** Given pairwise distances among points, recover the position of the points!
- **Example:** Distance between 10 US major cities

```
> UScitiesD
```

	Atlanta	Chicago	Denver	Houston	LosAngeles	Miami	NewYork	SanFrancisco	Seattle
Chicago	587								
Denver	1212	928							
Houston	701	940	879						
LosAngeles	1936	1745	831	1374					
Miami	604	1188	1726	968	2339				
NewYork	746	713	1631	1420	2451	1092			
SanFrancisco	2139	1858	949	1645	347	2594	2571		
Seattle	2182	1737	1021	1891	959	2734	2408	678	
Washington.DC	543	597	1494	1220	2300	923	205	2442	2329

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.4

Notes

---

---

---

---


---

---

---

Classical MDS: First Try

```
loc <- cmdscale(UScitiesD)
x <- loc[, 1]; y <- loc[, 2]
plot(x, y, type = "n", xlab = "", ylab = "", asp = 1,
     axes = FALSE, main = "cmdscale(UScitiesD)")
text(x, y, rownames(loc), cex = 0.8)
'''
```



Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.5

Notes

---

---

---

---


---

---

---

Classical MDS: Flip Axes

```
# Flip Axes
x1 <- -loc[, 1]; y1 <- -loc[, 2]
plot(x1, y1, type = "n", xlab = "", ylab = "", asp = 1,
     axes = FALSE, main = "cmdscale(UScitiesD)")
text(x1, y1, rownames(loc), cex = 0.8)
'''
```



Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.6

Notes

---

---

---

---

---

---

---

Another Example: Air Pollution in US Cities

```
> summary(dat)
      SO2      temp      manu      popul
Min.   : 8.00   Min.   :43.50   Min.   : 35.0   Min.   : 71.0
1st Qu.:13.00   1st Qu.:50.60   1st Qu.:181.0   1st Qu.:299.0
Median :26.00   Median :54.60   Median :347.0   Median :515.0
Mean   :30.05   Mean   :55.76   Mean   :463.1   Mean   :608.6
3rd Qu.:35.00   3rd Qu.:59.30   3rd Qu.:462.0   3rd Qu.:717.0
Max.   :110.00   Max.   :75.50   Max.   :3344.0   Max.   :3369.0

      wind      precip      predays
Min.   :6.000   Min.   :7.05   Min.   :36.0
1st Qu.:8.700   1st Qu.:30.96   1st Qu.:103.0
Median :9.300   Median :38.74   Median :115.0
Mean   :9.444   Mean   :36.77   Mean   :113.9
3rd Qu.:10.600   3rd Qu.:43.11   3rd Qu.:128.0
Max.   :12.700   Max.   :59.80   Max.   :166.0
```

- Range of manu and popul is much bigger than range of wind
- Need to standardize to give every variable equal weight

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.7

Notes

---

---

---


---

---

---

---

Air Pollution in US Cities Example



Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.8

Notes

---

---

---

---

---

---

---

Classical MDS: Technical Details

- **Input:**  $D = \{d_{ij}\}_{i,j=1}^n$ , the Euclidean distances between  $n$  objects in  $p$  dimensions
- **Output:**  $\{X_i\}_{i=1}^n$ , the “position” of points up to rotation, reflection, shift
- Two steps:
  - Compute inner products matrix  $B = XX^T$  from distance
$$b_{ij} = -\frac{1}{2}(d_{ij}^2 - d_{i.}^2 - d_{.j}^2 + d_{..}^2)$$
  - Perform spectral decomposition to compute positions from  $B$  (see next slide)

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.9

Notes

---

---

---

---

---

---

---

### Classical MDS: Technical Details

- Since  $B = XX^T$ , we need the “square root” of  $B$

- Since  $B$  is a symmetric and positive definite  $n \times n$  matrix  $\Rightarrow B$  can be diagonalized:

$$B = V\Lambda V^T$$

$\Lambda$  is a diagonal matrix with  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  on diagonal

- Assuming the rank of  $B = p$ , so that the last  $n - p$  of its eigenvalues will be zero  $\Rightarrow B$  can be written as

$$B = V_1\Lambda_1V_1^T,$$

where  $V_1$  contains the first  $p$  eigenvectors and  $\Lambda_1$  the  $p$  non-zero eigenvalues. Take “square root”:

$$X = V_1\Lambda_1^{-\frac{1}{2}}$$



### Notes

---

---

---

---

---

---

---

---

### Classical MDS: Low-Dimensional Representation

- Keep only few (e.g. 2) largest eigenvalues and corresponding eigenvectors


- The resulting  $X$  will be the low-dimensional representation we were looking for

- “Goodness of fit” (GOF) if we reduce to  $m$  dimensions:

$$\text{GOF} = \frac{\sum_{i=1}^m \lambda_i}{\sum_{i=1}^n \lambda_i}$$

- Finds “optimal” low-dim representation: Minimizes

$$S = \sum_{i=1}^n \sum_{j=1}^n (d_{ij}^2 - (d_{ij}^m)^2)^2$$



### Notes

---

---

---

---

---

---

---

---

### Classical MDS: Pros and Cons

- + Optimal for Euclidean input data
- + Still optimal, if  $B$  has non-negative eigenvalues
- + Very fast to compute
- - There is no guarantee it will be optimal if  $B$  has negative eigenvalues



### Notes

---

---

---

---

---


---

---

---

## Non-metric MDS: Idea

- Sometimes, there is no well-defined metric on original points
- Absolute values are not that meaningful, but the ranking is important
- Non-metric MDS finds a low-dimensional representation, which **respects the ranking of distances**



### Notes

---

---

---

---

---

---

---

## Non-metric MDS: Theory


- $\delta_{ij}$  is the true dissimilarity,  $d_{ij}$  is the distance of representation

- Minimize STRESS:

$$S = \frac{\sum_{i < j} (\theta(\delta_{ij}) - d_{ij})^2}{\sum_{i < j} d_{ij}^2},$$

where  $\theta(\cdot)$  is an increasing function

- Optimize over both position of points and  $\theta$
- $\hat{d}_{ij} = \theta(\delta_{ij})$  is called "disparity"
- Solved numerically (isotonic regression); Classical MDS as starting value; very time consuming



### Notes

---

---

---

---


---

---

---

## Non-metric MDS: Pros and Cons

- +: Fulfills a clear objective (minimize STRESS) without many assumptions
- +: Results don't change with rescaling or monotonic variable transformation
- +: Works even if you only have rank information
- -: computation can be slow in "large" problems
- -: Usually only local (not global) optimum found
- -: Only gets ranks of distances right



### Notes

---

---

---

---

---

---

---

House of Representatives Voting Data

Romesburg (1984) gives a set of data that shows the number of times 15 congressmen from New Jersey voted differently in the House of Representatives on 19 environmental bills

```
> voting[1:6, 1:6]
      Hunt(R) Sandman(R) Howard(D) Thompson(D) Freylinghuysen(R) Forsythe(R)
Hunt(R)      0         8       15       15         10         9
Sandman(R)    8         0       17       12         13        13
Howard(D)    15        17        0        9         16        12
Thompson(D)  15        12        9        0         14        12
Freylinghuysen(R) 10        13       16       14         0         8
Forsythe(R)   9         9       12       12         8         0
```

**Question:** Do people in the same party vote alike?

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.16

Notes

---

---

---

---


---

---

---

---

Non-metric MDS: Voting Example



Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.17

Notes

---

---

---

---

---

---

---

---

Summary

- **Classical MDS:**
  - Finds low-dim projection that respects distances
  - Optimal for euclidean distances
  - No clear guarantees for other distances
  - Fast to compute (can use `cmdscale` in R)
- **Non-metric MDS:**
  - Squeezes data points on table
  - Respects only rankings of distances
  - (Locally) solves clear objective
  - Computation can be slow (can use `isoMDS` from the R package "MASS")

Multidimensional Scaling

CLEMSON UNIVERSITY

Main Idea

Classical Multidimensional Scaling

Non-metric Multidimensional Scaling

14.18

Notes

---

---

---

---

---

---

---

---