

# Lecture 26

## Time Series Analysis

STAT 8020 Statistical Methods II  
December 1, 2020

Whitney Huang  
Clemson University

Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data  
Objectives of Time Series Analysis  
Features of Times Series  
Means & Autocovariances  
A Case Study

26.1

Notes

---

---

---

---

---

---

---

### Agenda

- 1 Time Series Data
- 2 Objectives of Time Series Analysis
- 3 Features of Times Series
- 4 Means & Autocovariances
- 5 A Case Study

Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data  
Objectives of Time Series Analysis  
Features of Times Series  
Means & Autocovariances  
A Case Study

26.2

Notes

---

---

---

---


---

---

---

### Level of Lake Huron 1875–1972

Annual measurements of the level of Lake Huron in feet.  
[Source: [Brockwell & Davis, 1991](#)]



Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data  
Objectives of Time Series Analysis  
Features of Times Series  
Means & Autocovariances  
A Case Study

26.3

Notes

---

---

---

---


---

---

---

Mauna Loa Atmospheric CO<sub>2</sub> Concentration

Monthly atmospheric concentrations of CO<sub>2</sub> at the Mauna Loa Observatory [Source: [Keeling & Whorf, Scripps Institution of Oceanography \(SIO\)](#)]



Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data

Objectives of Time Series Analysis

Features of Times Series

Means & Autocovariances

A Case Study

26.4

Notes

---

---

---


---

---

---

---

US Unemployment Rate 1948 Jan. – 2020 Oct.



Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data

Objectives of Time Series Analysis

Features of Times Series

Means & Autocovariances

A Case Study

26.5

Notes

---

---

---

---


---

---

---

Airflow Signal

A “normal” patient's 100 Hz sleep airflow signal [Source: [Huang et al. 2020+](#)]



Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data

Objectives of Time Series Analysis

Features of Times Series

Means & Autocovariances

A Case Study

26.6

Notes

---

---

---

---

---

---

---

## Time Series Data & Models

- A **time series** is a set of observations made sequentially in "time"
- **Time series analysis** is the area of statistics which deals with the analysis of **dependency** between different observations in time series data
- A **time series model** is a probabilistic model that describes ways that the series data  $\{y_t\}$  could have been generated
- More specifically, a time series model is usually a probability model for  $\{Y_t : t \in T\}$ , a **collection of random variables indexed in time**



### Notes

---

---

---

---

---

---

---

## Some Objectives of Time Series Analysis

- Find a **statistical model** that adequately explains the **dependence** observed in a time series
- To conduct **statistical inferences**, e.g., Is there evidence of a decreasing trend in the Lake Huron depths?
- To **forecast** future values of the time series based on those we have already observed



### Notes

---

---

---

---

---

---

---

## Features of Times Series

- **Trends**
  - One can think of trend,  $\mu_t$  as continuous changes, usually in the mean, over longer time scales
  - Usually the form of the trend is unknown and needs to be estimated. When the trend is removed, we obtain a **detrended** series
- **Seasonal or periodic components**
  - A seasonal component  $s_t$  constantly repeats itself in time, i.e.,  $s_t = s_{t+kd}$
  - We need to estimate the form and/or the period  $d$  of the seasonal component, i.e. to **deseasonalize** the series
- **The "noise" process**
  - The noise process,  $\eta_t$ , is the component that is neither trend nor seasonality
  - We will focus on finding plausible (typically stationary) statistical models for this process



### Notes

---

---

---

---

---

---

---

## Combining Trend $\mu_t$ , Seasonality $s_t$ , and Noise $\eta_t$ Together

There are two commonly used approaches

- Additive model:

$$y_t = \mu_t + s_t + \eta_t$$

- Multiplicative model:

$$y_t = \mu_t s_t \eta_t$$

If all  $\{y_t\}$  are positive then we obtain the additive model by taking logarithms:

$$\log y_t = \log \mu_t + \log s_t + \log \eta_t$$



### Notes

---

---

---

---

---

---

---

## Means, Autocovariances, and Stationary Processes

- The mean function of  $\{Y_t\}$  is

$$\mu_t = E[Y_t], \quad t \in T$$

- The autocovariance function of  $\{Y_t\}$  is

$$\gamma(t, t') = \text{Cov}(Y_t, Y_{t'}) = E[(Y_t - \mu_t)(Y_{t'} - \mu_{t'})], \quad t, t' \in T$$

When  $t = t'$  we obtain

$$\gamma(t, t') = \text{Cov}(Y_t, Y_t) = \text{Var}(Y_t) = \sigma_t^2, \text{ the variance function of } Y_t$$



### Notes

---

---

---

---

---

---

---

## Autocorrelation Function

The autocorrelation function (ACF) of  $\{Y_t\}$  is

$$\rho(t, t') = \text{Corr}(Y_t, Y_{t'}) = \frac{\gamma(t, t')}{\sqrt{\gamma(t, t)\gamma(t', t')}}.$$

It measures the strength of linear association between  $Y_t$  and  $Y_{t'}$

Properties:

- 1  $-1 \leq \rho(t, t') \leq 1, \quad t, t' \in T$
- 2  $\rho(t, t') = \rho(t', t), \quad \forall t, t' \in T; \rho(t, t) = 1, \quad \forall t \in T$
- 3  $\rho(t, t')$  is a non-negative definite function



### Notes

---

---

---

---

---

---

---

Stationary Processes

We will still try to keep our models for  $\{\eta_t\}$  as simple as possible by assuming stationarity, meaning that some characteristic of  $\{\eta_t\}$  does not depend on the time points, only on the “time lag” between time points:

- $E[\eta_t] = 0, \quad \forall t \in T$
- $Cov(\eta_t, \eta_{t'}) = \gamma(t' - t) = Cov(\eta_{t+s}, \eta_{t'+s})$

⇒ autocorrelation function (ACF):

$$\rho(h) = \frac{\gamma(h)}{\gamma(0)}$$

Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data

Objectives of Time Series Analysis

Features of Times Series

Means & Autocovariances

A Case Study

26.13

Notes

---

---

---

---

---

---

---

Autoregressive Moving Average (ARMA) Models

Let  $\{Z_t\}$  be independent and identical random variables that follow  $N(0, \sigma^2)$

- Moving Average Processes (MA(q)):**  
 $\eta_t = Z_t + \theta_1 Z_{t-1} + \theta_2 Z_{t-2} \cdots + \theta_q Z_{t-q}$
- Autoregressive Processes (AR(p)):**  
 $\eta_t = \phi_1 \eta_{t-1} + \phi_2 \eta_{t-2} + \cdots + \phi_p \eta_{t-p} + Z_t$
- Autoregressive Moving Average Processes ARMA(p,q):**  $\eta_t = \phi_1 \eta_{t-1} + \phi_2 \eta_{t-2} + \cdots + \phi_p \eta_{t-p} + Z_t + \theta_1 Z_{t-1} + \theta_2 Z_{t-2} + \cdots + \theta_q Z_{t-q}$

Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data

Objectives of Time Series Analysis

Features of Times Series

Means & Autocovariances

A Case Study

26.14

Notes

---

---

---


---

---

---

---

Autocorrelation Plot



Time Series Analysis

CLEMSON UNIVERSITY

Time Series Data

Objectives of Time Series Analysis

Features of Times Series

Means & Autocovariances

A Case Study

26.15

Notes

---

---

---

---

---

---

---

Lake Huron Case Study



- Detrending
- Model selection and fitting
- Forecasting

See R lab 22 for a demo

Time Series  
Analysis

CLEMSON  
UNIVERSITY

Time Series Data  
Objectives of Time  
Series Analysis  
Features of Times  
Series  
Means &  
Autocovariances  
A Case Study

26/16

Notes

---

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

---