

Lecture 12

Spectral Analysis of Time Series II

Readings: CC08 Chapter 14; BD16 Chapter 4, Chapter 10.1;
SS17 Chapter 1.5-1.6, Chapter 4.4-Chapter 4.6, Chapter 4.8,
Chapter 5.5

MATH 8090 Time Series Analysis
Week 12

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Whitney Huang
Clemson University

- 1 **Review**
- 2 **Southern Oscillation Index Example**
- 3 **Parametric Spectral Estimation**
- 4 **Lagged Regression Models**

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Review: Nonparametric Spectral Estimation

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

- **Periodogram:** $I(\omega_j) = |d(\omega_j)|^2$, where

$$d(\omega_j) = n^{-\frac{1}{2}} \sum_{t=1}^n y_t e^{-2\pi i \omega_j t}, \quad \omega_j = \frac{j}{n}, \quad j = 0, 1, \dots, n-1$$

- $\frac{I(\omega_j)}{\frac{1}{2}f(\omega_j)} \overset{\text{approx. i.i.d.}}{\sim} \chi^2_2, \quad j = 1, \dots, m = \frac{n-1}{2} \Rightarrow \mathbb{E}[I(\omega_j)] \approx f(\omega_j)$
(unbiased)
- But $\text{Var}[I(\omega_j)] \approx f^2(\omega_j)$ (inconsistent because it does not go to 0 as $n \rightarrow \infty$)

- Smooth the periodogram

- **Averaged periodogram:**

$$\bar{f}(\omega_j) = \frac{1}{L} \sum_{k=-l}^l I(\omega_{j+k}), \quad L = 2l + 1$$

- **Smoothed periodogram:** $\bar{f}(\omega_j) = \sum_{k=-l}^l W_l(k) I(\omega_{j+k})$

- Pointwise CI for $f(\omega_j)$:

$$\frac{\nu \bar{f}(\omega_j)}{\chi^2_\nu(1 - \alpha/2)} \leq f(\omega_j) \leq \frac{\nu \bar{f}(\omega_j)}{\chi^2_\nu(\alpha/2)}, \quad \nu = \frac{2}{\sum_{k=-l}^l W_l^2(k)}$$

- For odd $n = 2m + 1$, the inverse transform can be written

$$y_t - \bar{y} = \frac{2}{\sqrt{n}} \sum_{j=1}^m [d_{\cos}(\omega_j) \cos(2\pi\omega_j t) + d_{\sin}(\omega_j) \sin(2\pi\omega_j t)].$$

- Square and sum over t ; orthogonality of sines and cosines implies that

$$\begin{aligned} \sum_{t=1}^n (y_t - \bar{y})^2 &= 2 \sum_{j=1}^m [d_{\cos}(\omega_j)^2 + d_{\sin}(\omega_j)^2] \\ &= 2 \sum_{j=1}^m I(\omega_j) \end{aligned}$$

We have partitioned $\sum_{t=1}^n (y_t - \bar{y})^2$ into $2 \times \sum_{j=1}^m I(\omega_j)$. This leads to **Spectral ANOVA \Rightarrow total variance = sum of frequency components**

Source	df	SS	MS
ω_1	2	$2I(\omega_1)$	$I(\omega_1)$
ω_2	2	$2I(\omega_2)$	$I(\omega_2)$
\vdots	\vdots	\vdots	\vdots
ω_m	2	$2I(\omega_m)$	$I(\omega_m)$
Total	$2m = n - 1$	$\sum (y_t - \bar{y})^2$	

Toy example:

```
> x <- c(1, 2, 3, 2, 1) - mean(x)
> c1 <- cos(2 * pi * (1:5) * (1 / 5)); s1 <- sin(2 * pi * (1:5) * (1 / 5))
> c2 <- cos(2 * pi * (1:5) * (2 / 5)); s2 <- sin(2 * pi * (1:5) * (2 / 5))
> omega1 <- cbind(c1, s1); omega2 <- cbind(c2, s2)
> anova(lm(x ~ omega1 + omega2))
```

Warning in anova.lm(lm(x ~ omega1 + omega2)) :

ANOVA F-tests on an essentially perfect fit are unreliable

Analysis of Variance Table

Response: x

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
omega1	2	2.74164	1.37082		
omega2	2	0.05836	0.02918		
Residuals	0	0.00000			

Review



Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Example: Southern Oscillation Index (SOI)

Southern Oscillation Index (SOI) for a period of 453 months ranging over the years 1950-1987



What are the hidden periods of SOI?


Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

SOI Example: Raw Periodogram



An approximate 95% confidence interval for $f(\omega)$:

ω	Period	Power	Lower	Upper
$\frac{1}{48}$	4 years	0.0537	0.0146	2.1222
$\frac{1}{12}$	1 year	0.9722	0.2636	38.4011


Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

SOI Example: Averaged Periodogram (Daniell with $l = 4$)



An approximate 95% confidence interval for $f(\omega)$:

ω	Period	Power	Lower	Upper
$\frac{1}{48}$	4 years	0.0495	0.0279	0.1113
$\frac{1}{12}$	1 year	0.1191	0.0670	0.2677


Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

SOI Example: Smoothed Periodogram (modified Daniell $c(3, 3)$)



Review

Southern Oscillation
Index Example


Parametric Spectral
Estimation

Lagged Regression
Models

An approximate 95% confidence interval for $f(\omega)$:

ω	Period	Power	Lower	Upper
$\frac{1}{48}$	4 years	0.0502	0.0283	0.1129
$\frac{1}{12}$	1 year	0.1675	0.0943	0.3767

SOI Example: Apply Tapering to Alleviate Spectral Leakage



The tapered spectrum does a better job in separating the yearly cycle $\omega = 1$ and the El Niño cycle $\omega = \frac{1}{4}$

Seasonally Adjusted SOI [Source: Peter Bloomfield's ST 730 Lecture Notes]

- The Southern Oscillation Index data provided by Shumway and Stoffer **is not seasonally adjusted**, which explains the substantial peaks in the periodogram at the annual frequency


Seasonally Adjusted SOI [Source: Peter Bloomfield's ST 730 Lecture Notes]

- The Southern Oscillation Index data provided by Shumway and Stoffer **is not seasonally adjusted**, which explains the substantial peaks in the periodogram at the annual frequency
- So the series is non-stationary, and has neither an autocovariance function nor a spectral density function

Seasonally Adjusted SOI [Source: Peter Bloomfield's ST 730 Lecture Notes]

- The Southern Oscillation Index data provided by Shumway and Stoffer **is not seasonally adjusted**, which explains the substantial peaks in the periodogram at the annual frequency
- So the series is non-stationary, and has neither an autocovariance function nor a spectral density function
- A more sensible analysis uses the seasonally adjusted series. (Bloomfield did this by fitting a seasonal means model using data from 1876-2010.)

SOI Example from Bloomfield: Smoothed Periodogram



Note that the peak at the annual frequency disappear due to the removal of the annual cycle

- **Parametric estimation**: estimate a model that is specified by a fixed number of parameters
- **Nonparametric estimation**: estimate a model that is specified by a number of parameters that can grow as the sample grows

The smoothed periodogram estimates we have considered are **nonparametric**: the estimates of the spectral density can be parameterized by estimated values at ω_j 's. As $n \uparrow$, the number of distinct frequency values increases

The time domain models we considered are **parametric**. For example, an $\text{ARMA}(p,q)$ process can be completely specified with $p + q + 1$ parameters

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

The typical approach is to use the maximum likelihood parameter estimates $(\hat{\phi}_1, \dots, \hat{\phi}_p, \hat{\sigma}^2)$ for the parameters of an $AR(p)$, and then compute $\hat{f}(\omega)$ for this estimated AR model:

$$\hat{f}(\omega) = \frac{\hat{\sigma}^2}{|\hat{\phi}(e^{-2\pi i \omega})|^2}$$

For large n ,

$$\text{Var}(\hat{f}(\omega)) \approx \frac{2p}{n} f^2(\omega)$$

- The **bias** decreases as $p \uparrow$, the number of parameters increase, as one can model more complex spectra

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

The typical approach is to use the maximum likelihood parameter estimates $(\hat{\phi}_1, \dots, \hat{\phi}_p, \hat{\sigma}^2)$ for the parameters of an $AR(p)$, and then compute $\hat{f}(\omega)$ for this estimated AR model:

$$\hat{f}(\omega) = \frac{\hat{\sigma}^2}{|\hat{\phi}(e^{-2\pi i \omega})|^2}$$

For large n ,

$$\text{Var}(\hat{f}(\omega)) \approx \frac{2p}{n} f^2(\omega)$$

- The **bias** decreases as $p \uparrow$, the number of parameters increase, as one can model more complex spectra
- The **variance** increase linealy with p

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

- Sometimes ARMA models are used instead
- Estimate the parameters of an ARMA(p, q) model and compute its spectral density:

$$\hat{f}(\omega) = \hat{\sigma}^2 \left| \frac{\hat{\theta}(e^{-2\pi i\omega})}{\hat{\phi}(e^{-2\pi i\omega})} \right|^2.$$

- However, it is more common to use large AR models, rather than ARMA models

- The main advantage of parametric spectral estimation over nonparametric is that it often gives better **frequency resolution** of a small number of peaks

- The main advantage of parametric spectral estimation over nonparametric is that it often gives better **frequency resolution** of a small number of peaks
- This is especially important if there is more than one peak at nearby frequencies

- The main advantage of parametric spectral estimation over nonparametric is that it often gives better **frequency resolution** of a small number of peaks
- This is especially important if there is more than one peak at nearby frequencies
- The disadvantage of parametric spectral estimation is the inflexibility due to the use of the restricted class of ARMA models.

Given data y_1, y_2, \dots, y_n ,

- 1 Estimate the AR parameters $(\phi_1, \phi_2, \dots, \phi_p, \sigma^2)$ using maximum likelihood or Yule-Walker/least squares, choose a suitable model order p using AIC or BIC

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Given data y_1, y_2, \dots, y_n ,

- 1 Estimate the AR parameters $(\phi_1, \phi_2, \dots, \phi_p, \sigma^2)$ using maximum likelihood or Yule-Walker/least squares, choose a suitable model order p using AIC or BIC
- 2 Use the estimates $(\hat{\phi}_1, \hat{\phi}_2, \dots, \hat{\phi}_p, \hat{\sigma}^2)$ to compute the estimated spectral density:

$$\hat{f}(\omega) = \frac{\hat{\sigma}^2}{|\hat{\phi}(e^{-2\pi i \omega})|^2}$$

Example: AR(1) with $\phi = 0.8$



- 1 Use AIC to select p , the order of the AR model
- 2 Use the estimates $(\hat{\phi}_1, \hat{\phi}_2, \dots, \hat{\phi}_p, \hat{\sigma}^2)$ to compute the estimated spectral

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models



Example: ARMA(1, 1) with $\phi = 0.8$ and $\theta = 0.5$



- 1 Use AIC to select p , the order of the AR model
- 2 Use the estimates $(\hat{\phi}_1, \hat{\phi}_2, \dots, \hat{\phi}_p, \hat{\sigma}^2)$ to compute the estimated spectral

Review


Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models



Example: AR(2) with $\phi_1 = 1.5$ and $\phi_2 = -0.95$



- Black: True
- Blue: Parametric Estimate
- Red: Non-parametric Estimate (Daniell $l = 3$)

Review


Southern Oscillation
Index Example

Parametric Spectral
Estimation


Lagged Regression
Models

SOI Example

Series: soi | AIC order = 15



Series: soiAdj | AIC order = 14



[Review](#)

[Southern Oscillation
Index Example](#)

[Parametric Spectral
Estimation](#)

[Lagged Regression
Models](#)

A **lagged regression model** expresses the output time series as a linear combination of current and past (and possibly future) values of an input series:

$$Y_t = \sum_{h=-\infty}^{\infty} \beta_h X_{t-h} + V_t,$$

where

- X_t : observed **input time series**
- Y_t : observed **output time series**
- V_t : **stationary noise process**

Applications:

- Identify the (best linear) dynamic relationship between two time series, X_t and Y_t
- Forecast the future of Y_t based on X_t (often setting $\beta_h = 0$ for $h < 0$ to use only past inputs)

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models


An Example of Lagged Regression Model

Review

Southern Oscillation
Index Example


Parametric Spectral
Estimation

Lagged Regression
Models



- We may wish to identify how the values of the recruitment series is related to the SOI

An Example of Lagged Regression Model



- We may wish to identify how the values of the recruitment series is related to the SOI
- We may wish to predict future values of recruitment from the SOI

- **Time domain:** model the input series, extract the white time series driving it ("prewhitening"), regress with transformed output series
 - Cross-covariance function
 - Cross-correlation function
- **Frequency domain:** Calculate the input's spectral density, and the cross-spectral density between input and output, and find the **transfer function** relating them, in the frequency domain
 - Cross spectrum
 - Coherence

Recall that the autocovariance function of a stationary process $\{Y_t\}$ is

$$\gamma_X(h) = \mathbb{E}[(X_{t+h} - \mu_X)(X_t - \mu_X)].$$

The **cross-covariance function** of two jointly stationary processes $\{Y_t\}$ and $\{X_t\}$ is

$$\gamma_{XY}(h) = \mathbb{E}[(X_{t+h} - \mu_X)(Y_t - \mu_Y)].$$

Note: Jointly stationary = constant means, autocovariances depending only on the lag h , and cross-covariance depends only on h

[Review](#)

[Southern Oscillation
Index Example](#)

[Parametric Spectral
Estimation](#)

[Lagged Regression
Models](#)

The **cross-correlation function** of jointly stationary $\{X_t\}$ and $\{Y_t\}$ is

$$\rho_{XY}(h) = \frac{\gamma_{XY}(h)}{\sqrt{\gamma_X(0)\gamma_Y(0)}}.$$

Notice that $\rho_{XY}(h) = \rho_{YX}(-h)$ but $\rho_{XY}(h)$ is not necessarily equal to $\rho_{XY}(-h)$

Example: Suppose that $Y_t = \beta X_{t-\ell} + W_t$ for $\{X_t\}$ stationary and uncorrelated with $\{W_t\}$, and $\{W_t\}$ a zero mean white noise. Then $\{X_t\}$ and $\{Y_t\}$ are jointly stationary, with $\mu_Y = \beta\mu_X$,

$$\gamma_{XY}(h) = \beta\gamma_X(h + \ell).$$

- If $\ell > 0$, we say X_t leads Y_t
- If $\ell < 0$, we say X_t lags Y_t

[Review](#)[Southern Oscillation
Index Example](#)[Parametric Spectral
Estimation](#)[Lagged Regression
Models](#)


The sample cross-covariance is

$$\hat{\gamma}_{XY}(h) = \frac{1}{n} \sum_{i=1}^{n-h} (x_{t+h} - \bar{x})(y_t - \bar{y})$$

for $h \geq 0$. Then sample CCF is

$$\hat{\rho}_{XY}(h) = \frac{\hat{\gamma}_{XY}(h)}{\sqrt{\hat{\gamma}_X(0)\hat{\gamma}_Y(0)}}$$

SOI vs Recruitment



Example: CCF of SOI and recruitment has a peak at $h = -6$. Thus, SOI leads recruitment by 6 months

[Review](#)[Southern Oscillation
Index Example](#)[Parametric Spectral
Estimation](#)[Lagged Regression
Models](#)

Suppose we wish to fit a lagged regression model of the form

$$Y_t = \beta(B)X_t + V_t = \sum_{j=0}^{\infty} \beta_j X_{t-j} + V_t,$$

where X_t is an observed input series, Y_t is the observed output series, and V_t is a stationary noise process, uncorrelated with X_t

Suppose we wish to fit a lagged regression model of the form

$$Y_t = \beta(B)X_t + V_t = \sum_{j=0}^{\infty} \beta_j X_{t-j} + V_t,$$

where X_t is an observed input series, Y_t is the observed output series, and V_t is a stationary noise process, uncorrelated with X_t

One approach (pioneered by [Box and Jenkins](#)) is to fit ARMA models for X_t and V_t , and then find a simple rational representation for $\beta(B)$. This is the [transfer function models](#)

$$Y_t = \beta(B)X_t + V_t = \sum_{j=0}^{\infty} \beta_j X_{t-j} + V_t,$$

For example:

$$X_t = \frac{\theta_X(B)}{\phi_X(B)} W_t,$$

$$V_t = \frac{\theta_V(B)}{\phi_V(B)} Z_t,$$

$$\beta(B) = \frac{\delta(B)}{\omega(B)} B^d$$

Notice the delay B^d , indicating that Y_t lags X_t by d steps

How do we choose all of these parameters?

- Fit $\theta_X(B)$, $\phi_X(B)$ to model the input series $\{X_t\}$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

How do we choose all of these parameters?

- Fit $\theta_X(B)$, $\phi_X(B)$ to model the input series $\{X_t\}$
- **Prewhiten** the input series by applying the inverse operator $\phi_X(B)/\theta_X(B)$:

$$\tilde{Y}_t = \frac{\phi_X(B)}{\theta_X(B)} Y_t = \beta(B) W_t + \frac{\phi_X(B)}{\theta_X(B)} V_t$$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

How do we choose all of these parameters?

- Fit $\theta_X(B)$, $\phi_X(B)$ to model the input series $\{X_t\}$
- **Prewhiten** the input series by applying the inverse operator $\phi_X(B)/\theta_X(B)$:

$$\tilde{Y}_t = \frac{\phi_X(B)}{\theta_X(B)} Y_t = \beta(B) W_t + \frac{\phi_X(B)}{\theta_X(B)} V_t$$

- Calculate the cross-correlation of \tilde{Y}_t with W_t ,

$$\gamma_{\tilde{Y}, W}(h) = \mathbb{E}[\tilde{Y}_{t+h} W_t] = \mathbb{E}\left[\sum_{j=0}^{\infty} \beta_j W_{t+h-j} W_t\right] = \sigma_W^2 \beta_h$$

to give an indication of the behavior of $\beta(B)$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

How do we choose all of these parameters?

- Fit $\theta_X(B)$, $\phi_X(B)$ to model the input series $\{X_t\}$
- **Prewhiten** the input series by applying the inverse operator $\phi_X(B)/\theta_X(B)$:

$$\tilde{Y}_t = \frac{\phi_X(B)}{\theta_X(B)} Y_t = \beta(B) W_t + \frac{\phi_X(B)}{\theta_X(B)} V_t$$

- Calculate the cross-correlation of \tilde{Y}_t with W_t ,

$$\gamma_{\tilde{Y}, W}(h) = \mathbb{E}[\tilde{Y}_{t+h} W_t] = \mathbb{E}\left[\sum_{j=0}^{\infty} \beta_j W_{t+h-j} W_t\right] = \sigma_W^2 \beta_h$$

to give an indication of the behavior of $\beta(B)$

- Estimate the coefficients of $\beta(B)$ and hence fit an ARMA model for the noise series V_t

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

The prewhitening step inverts the linear filter $X_t = \frac{\theta_X(B)}{\phi_X(B)} W_t$.

Then the lagged regression is between the transformed Y_t and a white series W_t . This makes it easy to determine a suitable lag

Example: In the SOI/recruitment series, we treat SOI as an input, estimate an AR(1) model, prewhiten it, and consider the cross-correlation between the transformed recruitment series and the prewhitened SOI. This shows a large peak at lag -5 (corresponding to the SOI series leading the recruitment series)

This sequential estimation procedure ϕ_X, θ_X , then β , then ϕ_V, θ_V is rather ad hoc. [State space methods](#) (ARMAX model) offer an alternative, and they are also convenient for vector-valued input and output series

Lagged Regression in the Frequency Domain: Coherence

To analyze lagged regression in the frequency domain, we'll need the notion of **coherence**, the analog of cross-correlation in the frequency domain

Define the **cross-spectrum** as the Fourier transform of the cross-correlation,

$$f_{XY}(\omega) = \sum_{h=-\infty}^{\infty} \gamma_{XY}(h) e^{-2\pi i \omega h},$$
$$\gamma_{XY}(h) = \int_{-\frac{1}{2}}^{\frac{1}{2}} f_{XY}(\omega) e^{2\pi i \omega h} d\omega,$$

provided that $\sum_{h=-\infty}^{\infty} |\gamma_{XY}(h)| < \infty$

Notice that $f_{XY}(\omega)$ is complex: $f_{XY}(\omega) = c_{XY}(\omega) - iq_{XY}(\omega)$.
Also, $\gamma_{YX}(h) = \gamma_{XY}(-h)$ implies $f_{YX}(\omega) = \overline{f_{XY}(\omega)}$

$$\Rightarrow c_{YX}(\omega) = c_{XY}(\omega) \quad \text{and} \quad q_{YX}(\omega) = -q_{XY}(\omega)$$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

- The **squared coherence** function is

$$\rho_{Y,X}^2(\omega) = \frac{|f_{YX}(\omega)|^2}{f_X(\omega)f_Y(\omega)}.$$

measures the strength of the relationship between X_t and Y_t at frequency ω

- $\rho_{Y,X}^2(\omega)$ is an analog of R^2 , it measures the fraction of variance in Y_t at frequency ω , $f_Y(\omega)$, explained by X_t
- $\rho_{Y,X}^2(\omega) = |\rho_{Y,X}(\omega)|^2$, where

$$\rho_{Y,X}(\omega) = \frac{f_{YX}(\omega)}{\sqrt{f_X(\omega)f_Y(\omega)}}$$

Estimating Squared Coherence

Recall that we estimated the spectral density using the smoothed squared modulus of the DFT of the series,

$$\begin{aligned}\bar{f}_X(\omega_j) &= \frac{1}{L} \sum_{k=-(L-1)/2}^{(L-1)/2} |d_X(\omega_j)|^2 \\ &= \frac{1}{L} \sum_{k=-(L-1)/2}^{(L-1)/2} d_X(\omega_{j+k}) \overline{d_X(\omega_{j+k})}.\end{aligned}$$

We can estimate the cross spectral density using the same sample estimate,

$$\bar{f}_{XY}(\omega_j) = \frac{1}{L} \sum_{k=-(L-1)/2}^{(L-1)/2} d_X(\omega_{j+k}) \overline{d_Y(\omega_{j+k})}$$

Also, we can estimate the squared coherence using these estimates,

$$\bar{\rho}_{Y,X}^2(\omega) = \frac{|\bar{f}_{YX}(\omega)|^2}{\bar{f}_X(\omega) \bar{f}_Y(\omega)}.$$


Estimating Squared Coherence: SOI/Recruitment Example

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models



$$Y_t = \sum_{j=-\infty}^{\infty} \beta_j X_{t-j} + V_t$$

The projection theorem tells us that the coefficients that minimize the mean squared error,

$$\mathbb{E} \left[\left(Y_t - \sum_{j=-\infty}^{\infty} \beta_j X_{t-j} \right)^2 \right]$$

satisfy the orthogonality conditions

$$\mathbb{E} \left[\left(Y_t - \sum_{j=-\infty}^{\infty} \beta_j X_{t-j} \right) X_{t-k} \right] = 0, \quad k = 0, \pm 1, \pm 2, \dots$$

Taking the expectations inside leads to the normal equations

$$\sum_{j=-\infty}^{\infty} \beta_j \gamma_X(k-j) = \gamma_Y(k), \quad k = 0, \pm 1, \pm 2, \dots$$

We could solve these equations for the β_j using the sample autocovariance and sample cross-covariance. But it is more convenient to use estimates of the spectra and cross-spectrum because **convolution with $\{\beta_j\}$ in the time domain is equivalent to multiplication by the Fourier transform of $\{\beta_j\}$ in the frequency domain**

We replace the autocovariance and cross-covariance with the inverse Fourier transforms of the spectral density and cross-spectral density in the orthogonality conditions, i.e., replace

$$\sum_{j=-\infty}^{\infty} \beta_j \gamma_X(k-j) \quad k = 0, \pm 1, \pm 2, \dots$$

by

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \sum_{j=-\infty}^{\infty} \beta_j e^{2\pi i \omega(k-j)} f_X(\omega) d\omega$$

Lagged Regression Models in the Frequency Domain

This gives, for $k = 0, \pm 1, \pm 2, \dots$,

$$\begin{aligned}\int_{-\frac{1}{2}}^{\frac{1}{2}} \sum_{j=-\infty}^{\infty} \beta_j e^{2\pi i \omega(k-j)} f_X(\omega) d\omega &= \int_{-\frac{1}{2}}^{\frac{1}{2}} e^{2\pi i \omega k} f_{YX}(\omega) d\omega, \\ \Rightarrow \int_{-\frac{1}{2}}^{\frac{1}{2}} e^{2\pi i \omega k} B(\omega) f_X(\omega) d\omega &= \int_{-\frac{1}{2}}^{\frac{1}{2}} e^{2\pi i \omega k} f_{YX}(\omega) d\omega,\end{aligned}$$

where $B(\omega) = \sum_{j=-\infty}^{\infty} e^{-2\pi i \omega j} \beta_j$ is the Fourier transform of the coefficient sequence β_j . Since the Fourier transform is unique, the orthogonality conditions are equivalent to

$$B(\omega) f_X(\omega) = f_{YX}(\omega).$$

Then we may take

$$\hat{B}(\omega_k) = \frac{\hat{f}_{YX}(\omega_k)}{\hat{f}_X(\omega_k)}$$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Lagged Regression Models in the Frequency Domain

We can write the mean squared error at the solution as follows

$$\begin{aligned}\mathbb{E}\left[\left(Y_t - \sum_{j=-\infty}^{\infty} \beta_j X_{t-j}\right) Y_t\right] &= \gamma_Y(0) - \sum_{j=-\infty}^{\infty} \beta_j \gamma_{XY}(-j) \\&= \int_{-\frac{1}{2}}^{\frac{1}{2}} (f_Y(\omega) - B(\omega) f_{XY}(\omega)) d\omega \\&= \int_{-\frac{1}{2}}^{\frac{1}{2}} f_Y(\omega) \left(1 - \frac{f_{YX}(\omega) f_{XY}(\omega)}{f_X(\omega) f_Y(\omega)}\right) d\omega \\&= \int_{-\frac{1}{2}}^{\frac{1}{2}} f_Y(\omega) \left(1 - \frac{|f_{YX}(\omega)|^2}{f_X(\omega) f_Y(\omega)}\right) d\omega \\&= \int_{-\frac{1}{2}}^{\frac{1}{2}} f_Y(\omega) (1 - \rho_{Y,X}^2(\omega)) d\omega.\end{aligned}$$

$$\Rightarrow \text{MSE} = \int_{-\frac{1}{2}}^{\frac{1}{2}} f_Y(\omega) (1 - \rho_{Y,X}^2(\omega)) d\omega$$

$$\Rightarrow f_V(\omega) = (1 - \rho_{Y,X}^2(\omega)) f_Y(\omega)$$

Lagged Regression Models in the Frequency Domain

$$\text{Recall MSE} = \int_{-\frac{1}{2}}^{\frac{1}{2}} f_Y(\omega)(1 - \rho_{Y,X}^2(\omega)) d\omega.$$

Thus, $\rho_{Y,X}^2(\omega)$ indicates how the variance of $\{Y_t\}$ at a frequency ω is accounted for by $\{X_t\}$. Compare with the corresponding decomposition for random variables:

$$\mathbb{E}(Y - \beta X) = \sigma_Y^2(1 - \rho_{Y,X}^2)$$

We can estimate the β_j in the frequency domain:

$$\hat{B}(\omega_k) = \frac{\hat{f}_{YX}(\omega_k)}{\hat{f}_X(\omega_k)}.$$

We can approximate the inverse Fourier transform of $\hat{B}(\omega)$,

$$\hat{\beta}_j = \int_{-\frac{1}{2}}^{\frac{1}{2}} e^{2\pi i \omega j} \hat{B}(\omega) d\omega$$

via the sum,

$$\hat{\beta}_j = \frac{1}{M} \sum_{k=0}^{M-1} \hat{B}(\omega_k) e^{-2\pi i \omega_k j}.$$

Here is the procedure:

- 1 Estimate the spectral density $f_X(\omega)$ and cross-spectral density $f_{YX}(\omega)$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Here is the procedure:

- 1 Estimate the spectral density $f_X(\omega)$ and cross-spectral density $f_{YX}(\omega)$
- 2 Compute the transfer function $\hat{B}(\omega)$:

$$\hat{B}(\omega_k) = \frac{\hat{f}_{YX}(\omega_k)}{\hat{f}_X(\omega_k)}.$$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models

Here is the procedure:

- 1 Estimate the spectral density $f_X(\omega)$ and cross-spectral density $f_{YX}(\omega)$
- 2 Compute the transfer function $\hat{B}(\omega)$:

$$\hat{B}(\omega_k) = \frac{\hat{f}_{YX}(\omega_k)}{\hat{f}_X(\omega_k)}.$$

- 3 Take the inverse Fourier transform to obtain the impulse response function β_j :

$$\hat{\beta}_j = \frac{1}{M} \sum_{k=0}^{M-1} \hat{B}(\omega_k) e^{-2\pi i \omega_k j}.$$

Review

Southern Oscillation
Index Example

Parametric Spectral
Estimation

Lagged Regression
Models