
A
Bangladesh (R=1.17)
Poland (R=2.87)
Viet Nam (R=1.07)
Singapore (R=2.34)
Australia (R=0.34)
SDI–year reference (95% CI)
Reference-uncertain
Resolved, central
Low tail
High tail
LRI incidence per 100 000 children <5
10 000
3 000
1 000
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
0.9
Mean SDI, 2011–2023
B
Diarrhoeal mortality
LRI mortality
Meningitis mortality
Diarrhoeal incidence
LRI incidence
Meningitis incidence
0.69
0.79
0.79
0.69
0.84
0.79
0.5
0.6
0.7
0.8
0.9
1.0
Spearman ρ, 1990–2000 vs 2011–2023
C
Position in 2011–2023 (%)
100
80
60
40
20
0
11
29
59
70
9
21
High tail
Low tail
Position in 1990–2000 (n=231 each)
High tail
Reference-uncertain
Resolved, central
Low tail
D
17 locations had both
persistent high-tail and
persistent low-tail outcomes
Locations
40
30
20
10
0
38
36
22
23
8
13
4
4
0
4
1
0
1
2
3
4
5
6
Outcomes persistently in the tail (of 6)
Persistent high tail (73 locations)
Persistent low tail (80 locations)
E
Incidence–mortality Spearman ρ
1.0
0.8
0.6
0.4
0.2
0.0
Meningitis
12% opposite
(7–18)
LRI
24% opposite
(18–32)
Diarrhoeal
35% opposite
(27–44)
1990–
2000
2001–
2010
2011–
2023