
A
Rank stability, early vs late
Main analysis
Fixed curve shape
Period-specific fits
No time terms
Spline, 3 df
Spline, 5 df
Population-weighted fit
Small locations excluded
Leave-location-out reference
Tertile tails
Quintile tails
GBD uncertainty, 200 draws
0.4 0.5 0.6 0.7 0.8 0.9 1.0
Spearman ρ (range over six outcomes)
B
Early tails still in the same tail
High tail
Low tail
40 50 60 70 80
% of early-tail records
C
Correlation between diseases
Mortality
Incidence
0.0 0.2 0.4 0.6
Mean Spearman ρ over pairs and periods
D
Incidence vs mortality, same disease
Diarrhoeal
LRI
Meningitis
Main analysis
Fixed curve shape
Period-specific fits
No time terms
Spline, 3 df
Spline, 5 df
Population-weighted fit
Small locations excluded
Leave-location-out reference
Tertile tails
Quintile tails
GBD uncertainty, 200 draws
0.0 0.2 0.4 0.6 0.8 1.0
Spearman ρ, 2011–2023
E
Water and sanitation, high minus low tail
Low & low-middle SDI
High SDI
0 20 40 60
Percentage points
F
Catch-up in drinking water
Gap narrowed
Gap not narrowed
20 30 40 50 60 70 80
% of locations with falling mortality residual