
A
Death rates, fixed SDI groups
Deaths per 100 000 children <5
100
10
1
0.5
LRI
Diarrhoeal
Meningitis
Low SDI
High SDI
1990
2000
2010
2020
B
Relative gap by grouping design
Death rate ratio, low / high SDI
200
100
50
20
Diarrhoeal
LRI
Meningitis
1990
2000
2010
2020
C
Why grouping changes the LRI gap
Δ log rate ratio, 2023 (reassigned minus fixed)
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
−1.00
Low-SDI group change
High-SDI group change (−)
Net change (coloured by disease)
0.92×
0.52×
1.26×
Diarrhoeal
LRI
Meningitis
Fixed, 1990–2000 membership
Fixed, 2011–2023 membership
Reassigned each period
D
Example: position relative to the SDI–year reference (LRI incidence, 2011–2023)
LRI incidence per 100 000 children <5
10 000
3 000
1 000
Bangladesh (R=1.17)
Poland (R=2.87)
Viet Nam (R=1.07)
Singapore (R=2.34)
Australia (R=0.34)
SDI–year reference (95% CI)
Reference-uncertain
Resolved, central
Low tail
High tail
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
0.9
Mean SDI, 2011–2023
E
Late high-SDI locations
Diarrhoea death
6/32
8/44
LRI death
6/32
17/44
Meningitis death
5/32
8/44
LRI incidence
7/32
19/44
0
5
10
15
20
25
Locations in high tail, 2011–2023