
A
Rank stability of residuals
Diarrhoea, death
LRI, death
Meningitis, death
Diarrhoea, incidence
LRI, incidence
Meningitis, incidence
0.69
0.79
0.79
0.69
0.84
0.79
0.6
0.8
1.0
Spearman ρ, 1990–2000 vs 2011–2023
B
Where early tails ended up
Position in 2011–2023 (%)
100
80
60
40
20
0
11
29
59
70
9
21
High tail
Low tail
Position in 1990–2000 (n=231 records each)
High tail
Reference-uncertain
Resolved, central
Low tail
C
Correlation between outcomes
Diar death
LRI death
Men death
Diar inc.
LRI inc.
Men inc.
0.49 0.53 0.29 0.29 0.39
0.49 0.48 0.08 0.55 0.51
0.53 0.48 0.22 0.20 0.73
0.29 0.08 0.22 0.28 0.20
0.29 0.55 0.20 0.28 0.25
0.39 0.51 0.73 0.20 0.25
1.0
0.5
0.0
−0.5
−1.0
D
Pairs of diseases
Spearman ρ between diseases
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0.0
Mortality
Incidence
1990–2000
2001–2010
2011–2023
E
Incidence vs death, same disease
Incidence–death Spearman ρ
1.0
0.8
0.6
0.4
0.2
0.0
Meningitis
12% opposite
LRI
24% opposite
Diarrhoeal
35% opposite
1990–2000
2001–2010
2011–2023
F
Diarrhoeal disease, 2011–2023 (ρ=0.29)
Mortality residual (log)
2
0
−2
Greenland
Australia
Other locations
High tail, both
Low tail, both
−2
0
2
Incidence residual (log)