
A
Position relative to development: LRI incidence, 2011–2023
Bangladesh (R=1.17)
Poland (R=2.87)
Viet Nam (R=1.07)
Singapore (R=2.34)
Australia (R=0.34)
SDI–year reference (95% CI)
Reference-uncertain
Resolved, central
Low tail
High tail
LRI incidence per 100 000 children <5
10 000
3 000
1 000
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
0.9
Mean SDI, 2011–2023
B
Rank stability
Diarrhoeal mortality
LRI mortality
Meningitis mortality
Diarrhoeal incidence
LRI incidence
Meningitis incidence
0.69
0.79
0.79
0.69
0.84
0.79
0.5
0.6
0.7
0.8
0.9
1.0
Spearman ρ, 1990–2000 vs 2011–2023
C
Where early tails ended up
Position in 2011–2023 (%)
100
80
60
40
20
0
11
29
59
70
9
21
High tail
Low tail
Position in 1990–2000 (n=231 each)
High tail
Reference-uncertain
Resolved, central
Low tail
D
Persistent tails by number of outcomes
Locations
40
30
20
10
0
38
36
22
23
8
13
4
4
0
4
1
0
17 locations had both persistent high-tail and persistent low-tail outcomes
1
2
3
4
5
6
Outcomes persistently in the tail (of 6)
Persistent high tail (73 locations)
Persistent low tail (80 locations)
E
Incidence vs mortality
Incidence–mortality Spearman ρ
1.0
0.8
0.6
0.4
0.2
0.0
Meningitis
12% opposite
(7–18)
LRI
24% opposite
(18–32)
Diarrhoeal
35% opposite
(27–44)
1990–
2000
2001–
2010
2011–
2023