
A
Low and low-middle SDI
Middle and high-middle SDI
High SDI
Basic services
Immunisation
Nutrition
Health system
Air pollution
Drinking water
Sanitation
Clean cooking fuels
DTP3
Hib3
PCV3
Stunting
Low birthweight
Undernourishment
UHC index
Health expenditure
Hospital beds
Nurses and midwives
Ambient PM2.5
Second-hand smoke
n high/low
13/17 17/17 7/11 10/13 17/17 20/20 9/8 8/6 6/8 7/10 6/9 7/7 17/14 14/16 26/20 23/17 17/14 13/13
Diar. inc.
Diar. mort.
LRI inc.
LRI mort.
Men. inc.
Men. mort.
100
50
0
−50
−100
High tail minus low tail, % with adverse departure
B
Profiles by domain, 2011–2023
Basic services
Immunisation
Nutrition
Health system
Air pollution
−40 −20 0 20 40 60 80
High tail minus low tail, % adverse (pp)
Low and low-middle
Middle and high-middle
High
Mortality
Incidence
C
Profile change with tail membership
Basic services
Immunisation
Nutrition
Health system
Air pollution
−30 −20 −10 0 10 20 30
Difference in change of % adverse, 2001–2010 to 2011–2023 (pp)
Left vs stayed in high tail
Entered high tail vs stayed in low tail