
Identification and characterization of retinoblastoma tumor progressionrelated somatic copy number alteration (SCNA) driver genes.

Kevin Stachelek^{1,2}, Jinlun Bai¹ James Hicks^{5,6} and David Cobrinik^{1,3,4,5}

¹The Vision Center and Saban Research Institute, Children's Hospital Los Angeles, Los Angeles, CA; ²Cancer Biology and Genomics Program, Keck School of Medicine, University of Southern California, Los Angeles, CA; ³Department of Ophthalmology and Roski Eye Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA; ³Department of Ophthalmology and Roski Eye Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA; ⁵Norris Cancer Center, Cancer Cen CA; ⁶Department of Biological Sciences, USC Dornsife College of Letters, Arts, and Sciences, University of Southern California, Los Angeles





THE SABAN RESEARCH INSTITUTE






- Identify retinoblastoma tumors with subclonal heterogeneity for SCNAs
- Capture scRNAseg information
- Infer SCNA subclones and identify SCNA-related gene
- Define effects of candidate 16q- driver genes in a retinal organoid model of retinoblastoma progression.
 - Identify candidate driver genes with downregulation in 16q- retinoblastoma tumors
- Create heterozygous knockout of candidate 16q- drivers by CRISPR inactivation of GFP-marked RB-/- organoid cone precursors.
- Define effects of hemizygous knockout by measuring immunofluorescence staining





Numbat analysis of existing scRNAseg reveals SCNA subclones -AMP -DEL 4 5 6 7 8 9 10 11 12 14 16 18 20 2p+GT-2 16q-14-GT-3 8p-11p-SCNA subclone proportions vary according to cell states

SCNA inference reveals clonal dynamics in retinoblastoma




SCNA subclone proportions vary according cell cycle and proliferation state




Aim 2 Approach

Define effects of candidate 16g- driver genes by CRISPR-mediated heterozygous knockout in a retinal organoid model




A novel chimeric stem cell-derived retinal organoid model with RB-/-. cone-specific GNAT2-GFP, and Cerulean-Ki67 expression




Aim 2 Preliminary Studies

Identification of 16q genes downregulated in Subtype 1 (early)



Differential expression (16q- vs. diploid)



Potential tumor suppressive features of SETD6, DDX19A, and RBL2

- SETD6: protein lysine methyltransferase that methylates transcriptional coactivator BRD4 to negatively regulate genes involved in translation and inhibit total mRNA translation13.
- DDX19A: DEAD-box ATPase that remodels mRNPs at the nuclear pore complex to terminate mRNP export.
- RBL2: pRB-related; combined RB1 and RBL2 KD leads to more rapid retinoblastoma tumors
- These activities are highly relevant to retinoblastoma where activation of translation related genes by MYCN is implicated in tumor progression.

Conclusions

- scGTO was used to concurrently define single cell SCNAs and
- scGTO-derived transcriptomes formed the same transcriptomic groups as standard Smart-seq based scRNA-seq.
- Retinoblastoma cells with recurrent SCNAs contributed to the same transcriptomic groups as cells with no SCNAs, implying that differential expression should be performed between cells with and without SCNAs within each transcriptomic group.
- Application of Numbat to a public retinoblastoma scRNA-seq dataset revealed patterns of SCNA subclone distribution according to gene expression states

- Identified candidate 16q- driver genes in a publicly available
- CRISPR-based indel production of SETD6, DDX19A, and RBL2 is in progress



Acknowledgements Thesis Committee David Craig David Cobrinik

James Hicks

Suhn Rhie

Research to Prevent Blindness
NIH K08CA232344 (JLB)
He Larry and Celia Moh Foundat
The Naconatal Blindness Research

the St. Baldrick's Foundati NIH R01CA137124 (DC).