

# Irodov: Problems in General Physics

Will Huang

April 25, 2023

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Notation and Conventions . . . . .	2
<b>2</b>	<b>Physical Fundamentals of Mechanics</b>	<b>4</b>
2.1	Kinematics . . . . .	4
2.2	Laws of Conservation . . . . .	8

# 1 Introduction

These are my solutions to every problem in Igor Irodov's *Problem in General Physics* as part of my efforts to prepare for graduate studies in physics. They are organized according to the sections given in the book. Irodov provides some hints (more guidance) at the beginning of the book which I will rehash here as I will make every effort to follow this paradigm for each problem:

- Being each problem by recognizing its meaning and the data needed to arrive at an answer. Missing data may be found in the reference tables he provides in the appendices. Start each problem by drawing a diagram, if possible, "elucidating the essence of the problem."
- Solve each problem in the general form, that is, in letter notation. Verify that this solution has the correct dimensions as expected for the problem and, if possible, investigate its behavior in certain extreme cases (he provides the example that gravitational force between two bodies must turn into the well known law of gravitational attraction as length approaches infinity).
- When starting the final numerical calculation, keep in mind that physical quantities are always approximate. Adhere to the proper rules when operating with approximations and pay attention to numerical accuracy. When a final numerical answer is achieved, evaluate its plausibility (i.e. velocity should not surpass the speed of light).

## 1.1 Notation and Conventions

The notation in this book is a mix of Irodov's and my own. In particular there are a few conventions that he adopts that I am not familiar with (i.e.  $w$  for acceleration), so I will use the notation that I am familiar with in place of those. The beginning of each subsection will include formulas defined the way I use them to make any differences in notation/convention clear. Additional conventions are listed in this subsection.

All vectors will be written using boldface, e.g.  $\mathbf{x}, \mathbf{F}$ , and their magnitudes written in normal italics, e.g.  $r = |\mathbf{r}|$ . The unit vectors are:

- $\mathbf{i}, \mathbf{j}, \mathbf{k}$  in Cartesian coordinates
- $\rho, \theta, \phi$  in spherical coordinates ( $\phi$  is azimuthal)

The normal and tangential components of a vector will be denote  $\mathbf{x}_n, \mathbf{x}_t$  respectively.

Differences are denoted in one of three ways:

- $\Delta$  denotes a finite difference, e.g.  $\Delta x = x_2 - x_1$
- $d$  denotes an infinitesimal increment, e.g.  $dr$
- $\delta$  denotes an elementary value, e.g.  $\delta W$  the elementary work<sup>1</sup>

---

<sup>1</sup>I have not seen the term elementary value used before. From looking up the work example, I believe this is close enough to the more familiar concept of variation for which  $\delta$  usually denotes.

The time derivative is denoted, in typical physics fashion, with a dot over the variable, e.g.  $\mathbf{v} \equiv \dot{\mathbf{x}}$ . The three vector operations are denoted using a nabla:

- $\nabla\phi$  the gradient of a potential function  $\phi$
- $\nabla \cdot \mathbf{E}$  the divergence of a vector field  $\mathbf{E}$
- $\nabla \times \mathbf{E}$  the curl of a vector field  $\mathbf{E}$

Integrals of any multiplicity will all use a single sign  $\int$  and will be distinguished only by the integrating element:  $dx$  or  $d\mathbf{x}$ ,  $d\mathbf{S}$ ,  $dV$  for line, surface, and volume integrals respectively. An integral over a closed surface or loop will be denoted with the sign  $\oint$ .

## 2 Physical Fundamentals of Mechanics

### 2.1 Kinematics

For a given point undergoing motion, the average and instantaneous velocities are

$$\langle \mathbf{v} \rangle = \frac{\Delta \mathbf{x}}{\Delta t} \quad \mathbf{v} = \frac{d\mathbf{x}}{dt}$$

where  $\mathbf{x}$  is the displacement vector. Similarly the average and instantaneous accelerations are

$$\langle \mathbf{a} \rangle = \frac{\Delta \mathbf{v}}{\Delta t} \quad \mathbf{a} = \frac{d\mathbf{v}}{dt}$$

The distance covered by a point is given by the integral

$$s = \int_{t_0}^{t_1} v dt$$

The acceleration of a point may be broken down into components tangential and normal to the a trajectory, let  $R$  be the radius of curvature at a given point, then the components are:

$$a_t = \frac{dv_t}{dt} \quad a_n = \frac{v^2}{R}$$

The angular velocity and accelerations are given by

$$\omega = \frac{d\theta}{dt} \quad \alpha = \frac{d\omega}{dt}$$

To convert between angular and linear quantities for a rotating body

$$\mathbf{v} = (\omega \mathbf{r}) \quad \omega_n = \omega^2 R \quad |\omega_t| = \alpha R$$

where  $R$  is the distance from the rotation action and  $\mathbf{r}$  is the radius vector.

**Exercise 1.**

**Exercise 2.**

**Exercise 3.**

**Exercise 4.**

**Exercise 5.**

**Exercise 6.**

Exercise 7.

Exercise 8.

Exercise 9.

Exercise 10.

Exercise 11.

Exercise 12.

Exercise 13.

Exercise 14.

Exercise 15.

Exercise 16.

Exercise 17.

Exercise 18.

Exercise 19.

Exercise 20.

Exercise 21.

Exercise 22.

Exercise 23.

Exercise 24.

Exercise 25.

Exercise 26.

Exercise 27.

Exercise 28.

Exercise 29.

Exercise 30.

Exercise 31.

Exercise 32.

Exercise 33.

Exercise 34.

Exercise 35.

Exercise 36.

Exercise 37.

Exercise 38.

Exercise 39.

Exercise 40.

Exercise 41.

Exercise 42.

Exercise 43.

Exercise 44.

Exercise 45.

Exercise 46.

Exercise 47.

Exercise 48.

Exercise 49.

Exercise 50.

Exercise 51.

Exercise 52.

Exercise 53.

Exercise 54.

Exercise 55.

Exercise 56.

Exercise 57.

Exercise 58.