

⑤ Let $h(x) = \frac{a^T x + \alpha}{b^T x + \beta}$ and $K = \{x \in \mathbb{R}^n : b^T x + \beta > 0\}$.

Obviously $\nabla h(x) = \frac{(b^T x + \beta)a - (a^T x + \alpha)b}{(b^T x + \beta)^2}$

thus

$$\nabla h(x)^T (y-x) \geq 0 \Leftrightarrow (b^T x + \beta) a^T (y-x) \geq (a^T x + \alpha) b^T (y-x)$$

If $x \in K$,

$$a^T (y-x) \geq \frac{a^T x + \alpha}{b^T x + \beta} b^T (y-x), \text{ or equivalently}$$

$$a^T y + \alpha - a^T x - \alpha \geq \frac{a^T x + \alpha}{b^T x + \beta} (b^T y + \beta - b^T x - \beta), \text{ or}$$

$$a^T y + \alpha \geq \frac{a^T x + \alpha}{b^T x + \beta} (b^T y + \beta - \cancel{b^T x - \beta} + \cancel{b^T x + \beta}),$$

$$h(y) = \frac{a^T y + \alpha}{b^T y + \beta} \geq \frac{a^T x + \alpha}{b^T x + \beta} = h(x). \text{ Hence } h \text{ is pseudoconvex.}$$

This implies also that $\frac{(-a)^T x + (-\alpha)}{b^T x + \beta}$ is pseudoconvex,

which asserts that $-h$ is pseudoconvex.

⑥ Evidently $\nabla f(x_1, x_2) = 2(x_1 - \frac{3}{2}, x_2 - 5)$ and so
 $\nabla f(1, 3) = (-1, -4)$. This yields

$$\nabla f(1, 3)^T (x_1, x_2) - (1, 3) = -x_1 + 4x_2 + 13 \dots (1)$$

On the other hand,

$$(x_1, x_2) \in K \Rightarrow -x_1 + x_2 \leq 2 \text{ and } 2x_1 + 3x_2 \leq 11$$

$$\Rightarrow (\text{summing up}) \quad x_1 + 4x_2 \leq 13 \dots (2)$$

(1) and (2) give

$$\nabla f(\bar{x}) (x - \bar{x}) \geq 0 \quad \forall x \in K,$$

which means that \bar{x} is minimum of f on K
 in view of the convexity of f .

② (a) ✓ (b) ✓

(c) Clearly $C_1 \cup C_2$ is a cone.

$$\left. \begin{array}{l} C_1 \subseteq C_1 + C_2 \\ C_2 \subseteq C_1 + C_2 \end{array} \right\} \Rightarrow C_1 \cup C_2 \subseteq C_1 + C_2 \Rightarrow \text{co}(C_1 \cup C_2) \subseteq C_1 + C_2$$

since $C_1 + C_2$ is convex (by (b)).

Let $x_1 \in C_1, x_2 \in C_2$. Then

$$x_1 + x_2 = \frac{1}{2}(2x_1) + \frac{1}{2}(2x_2) \in \frac{1}{2}(C_1 \cup C_2) + \frac{1}{2}(C_1 \cup C_2)$$

$$\Rightarrow x_1 + x_2 \in \text{co}(C_1 \cup C_2) \therefore C_1 + C_2 \subseteq \text{co}(C_1 \cup C_2)$$

$$\begin{aligned}
 \textcircled{7} \quad (a) \quad f(ty_1 + (1-t)y_2) &= \sup_{x \in K} (ty_1 + (1-t)y_2)^T x \quad (0 < t < 1) \\
 &= \sup_{x \in K} (ty_1^T x + (1-t)y_2^T x) \leq \sup_{x \in K} ty_1^T x + \sup_{x \in K} (1-t)y_2^T x \\
 &= t \sup_{x \in K} y_1^T x + (1-t) \sup_{x \in K} y_2^T x = t f(y_1) + (1-t) f(y_2).
 \end{aligned}$$

(b) We need to prove

$$f(z) \geq f(y) + \bar{x}^T(z-y) \quad \forall z \in \mathbb{R}^n.$$

$$\text{Clearly, for } z \in \mathbb{R}^n, \quad f(z) = \sup_{x \in K} z^T x \geq z^T x \quad \forall x \in K.$$

In particular,

$$f(z) \geq z^T \bar{x} = z^T \bar{x} + y^T \bar{x} - y^T \bar{x} = y^T \bar{x} + \bar{x}^T(z-y)$$

$$\Rightarrow f(z) \geq f(y) + \bar{x}^T(z-y), \text{ i.e., } \bar{x} \in \partial f(y).$$

$$\textcircled{3} \quad n=1, \quad A = \mathbb{Q} \text{ (los números racionales)}. \quad L = \mathbb{R} \text{ y } (a \in A)$$

(b) se cumple claramente, y

(a) no!

$\textcircled{4}$ Since M is a convex cone and every $x \in M, x \neq 0$, verifies $x = \frac{1}{2}(0) + \frac{1}{2}(2x)$, x cannot be extremal. Thus the only possible extremal point of M should be the origin. For example, $n=2, A = \begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix}$.