

$$h(x, y) = \arctan \left[\frac{x+y}{x-y} \right]$$

• Curvas de nivel

$$z = h(x, y) \Leftrightarrow z = \arctan \left[\frac{x+y}{x-y} \right]$$


$$\Leftrightarrow \tan(z) = \frac{x+y}{x-y}$$

$$\Leftrightarrow z = \frac{x+y}{x-y}$$

$$z=0 \Rightarrow x=-y$$

$$z=1 \Rightarrow x-y = x+y \Rightarrow y=0$$

$$z=-1 \Rightarrow y-x = x+y \Rightarrow x=0$$



$$z=2 \Rightarrow 2x-2y = x+y \Rightarrow x=3y \Rightarrow y = \frac{1}{3}x$$

$$z=-2 \Rightarrow -2x+2y = x+y \Rightarrow y=3x$$

veamos como se comporta $h(x, y)$ en las siguientes trayectorias

$$y = -x$$

$$h(x, -x) = \arctan \left(\frac{0}{2x} \right) = \pi$$

$$y = 0$$

$$h(x, 0) = \arctan(1)$$

$$x=0$$

$$h(0, y) = \arctan(-1)$$

$$y = \frac{1}{3}x$$

$$h(x, \frac{1}{3}x) = \arctan\left(\frac{x + \frac{1}{3}x}{x - \frac{1}{3}x}\right) = \arctan\left(\frac{4x}{2x}\right) = \arctan(2)$$

$$y = 3x$$

$$h(x, 3x) = \arctan\left(\frac{x + 3x}{x - 3x}\right) = \arctan\left(\frac{4x}{-2x}\right) = \arctan(-2)$$

$$\lim_{(x,y) \rightarrow (-1,1)} h(x,y) = \lim_{(x,y) \rightarrow (-1,1)} \arctan\left(\frac{x+y}{x-y}\right)$$

$$\text{Si } x = -1$$

$$g(u, v) = f(u+v, u-v)$$

$$\frac{\partial g}{\partial u} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial u} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}$$

$$\frac{\partial g}{\partial v} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial v} = \frac{\partial f}{\partial x} - \frac{\partial f}{\partial y}$$

$$\begin{aligned} \bullet \frac{\partial^2 g}{\partial u^2} &= \frac{\partial}{\partial u} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial x^2} \cdot \frac{\partial x}{\partial u} + \frac{\partial^2 f}{\partial x \partial y} \frac{\partial y}{\partial u} + \frac{\partial^2 f}{\partial y \partial x} \frac{\partial x}{\partial u} \\ &\quad + \frac{\partial^2 f}{\partial y^2} \frac{\partial y}{\partial u} \\ &= \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial x \partial y} + \frac{\partial^2 f}{\partial y \partial x} + \frac{\partial^2 f}{\partial y^2} \\ &= \frac{\partial^2 f}{\partial x^2} + 2 \frac{\partial^2 f}{\partial x \partial y} + \frac{\partial^2 f}{\partial y^2} \end{aligned}$$

$$\begin{aligned} \bullet \frac{\partial^2 g}{\partial v^2} &= \frac{\partial^2 f}{\partial x^2} \cdot \frac{\partial x}{\partial v} + \frac{\partial^2 f}{\partial x \partial y} \cdot \frac{\partial y}{\partial v} - \frac{\partial^2 f}{\partial y \partial x} \frac{\partial x}{\partial v} - \frac{\partial^2 f}{\partial y^2} \cdot \frac{\partial y}{\partial v} \\ &= \frac{\partial^2 f}{\partial x^2} - 2 \frac{\partial^2 f}{\partial x \partial y} + \frac{\partial^2 f}{\partial y^2} \end{aligned}$$

$$g(u, v) = f(x(u, v), y(u, v))$$

$$x = u + v$$

$$y = u - v$$

$$\Rightarrow u = \frac{x+y}{2}, v = \frac{x-y}{2}$$

$$f(x, y) = g\left(\frac{x+y}{2}, \frac{x-y}{2}\right)$$

$$\frac{\partial f}{\partial x} = \frac{\partial g}{\partial u} \cdot \frac{\partial u}{\partial x} + \frac{\partial g}{\partial v} \cdot \frac{\partial v}{\partial x} = \frac{\partial g}{\partial u} \cdot \frac{1}{2} + \frac{1}{2} \frac{\partial g}{\partial v}$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 g}{\partial u^2} \cdot \frac{1}{4} + \frac{\partial^2 g}{\partial u \partial v} \cdot \frac{1}{4} + \frac{1}{4} \frac{\partial^2 g}{\partial v \partial u} + \frac{1}{4} \frac{\partial^2 g}{\partial v^2}$$

$$= \frac{1}{4} \frac{\partial^2 g}{\partial u^2} + \frac{1}{2} \frac{\partial^2 g}{\partial u \partial v} + \frac{1}{4} \frac{\partial^2 g}{\partial v^2}$$

$$\frac{\partial f}{\partial y} = \frac{1}{2} \frac{\partial g}{\partial u} - \frac{1}{2} \frac{\partial g}{\partial v}$$

$$\frac{\partial^2 f}{\partial y^2} = \frac{1}{4} \frac{\partial^2 g}{\partial u^2} - \frac{1}{4} \frac{\partial^2 g}{\partial u \partial v} - \frac{1}{4} \frac{\partial^2 g}{\partial v \partial u} + \frac{1}{4} \frac{\partial^2 g}{\partial v^2}$$

längs $\frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial y^2}$

$$\Rightarrow \cancel{\frac{1}{4} \frac{\partial^2 g}{\partial u^2}} + \frac{1}{2} \frac{\partial^2 g}{\partial u \partial v} + \cancel{\frac{1}{4} \frac{\partial^2 g}{\partial v^2}} = \cancel{\frac{1}{4} \frac{\partial^2 g}{\partial u^2}} - \frac{1}{4} \frac{\partial^2 g}{\partial u \partial v} - \frac{1}{4} \frac{\partial^2 g}{\partial v \partial u} + \cancel{\frac{1}{4} \frac{\partial^2 g}{\partial v^2}}$$

$$\Rightarrow \frac{1}{2} \frac{\partial^2 g}{\partial u \partial v} = -\frac{1}{2} \frac{\partial^2 g}{\partial u \partial v}$$

$$\Rightarrow \frac{\partial^2 g}{\partial u \partial v} = 0 \quad / \int \Rightarrow \int \frac{\partial^2 g}{\partial u \partial v} \cdot dv = 0$$

$$\Rightarrow \frac{\partial g}{\partial u} = F(u) / \int$$

$$\Rightarrow g(u, v) = F(u) + G(v)$$

$$\Rightarrow g(u(x,y), v(x,y)) = F\left(\frac{x+y}{2}\right) + G\left(\frac{x-y}{2}\right)$$

$$\Rightarrow f\left(\frac{x+y}{2} + \frac{x-y}{2}, \frac{x+y}{2} - \frac{x-y}{2}\right) = f(x, y)$$

$$\Rightarrow f(x, y) = F\left(\frac{x+y}{2}\right) + G\left(\frac{x-y}{2}\right)$$

$$L = \lim_{(x,y) \rightarrow (0,0)} \frac{4x^2 + 2xy + 5y^2}{3x^2 + 5y^2}$$

considera la trayectoria $T = \{(x,y) \in \mathbb{R}^2 : y = mx, m > 0\}$

$$\lim_{x \rightarrow 0} \frac{4x^2 + 2mx^2 + 5m^2x^2}{3x^2 + 5m^2x^2} = \lim_{x \rightarrow 0} \frac{(4 + 2m + 5m^2)\cancel{x^2}}{(3 + m^2)\cancel{x^2}}$$

$$= \frac{4 + 2m + 5m^2}{3 + m^2}$$

luego, el límite anterior depende del valor de m , por lo que L depende de la trayectoria $\Rightarrow L$ no existe.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y}{x^2 + y^2}$$

veamos que sucede con la trayectoria $T = \{(x,y) \in \mathbb{R}^2 : x = 0\}$

$$\lim_{\substack{(x,y) \rightarrow (0,0) \\ x=0}} \frac{-y}{y^2} = \lim_{y \rightarrow 0} \frac{-1}{y}, \text{ este límite no existe}$$

$\Rightarrow L$ no existe.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^5}{x^2 + y^{10}} = \lim_{(x,y) \rightarrow (0,0)} \frac{x(y^5)}{x^2 + (y^5)^2} = \lim_{(x,u) \rightarrow (0,0)} \frac{xu}{x^2 + u^2} = 0$$

$$\text{Sea } u = y^5 \text{ si } y \rightarrow 0 \Rightarrow u \rightarrow 0$$

$$\left| \frac{xu}{x^2 + u^2} \right| = \frac{|xu|}{|x^2 + u^2|} \leq \frac{|xu| \cdot |x|}{x^2 + u^2} \leq \frac{\cancel{x^2}|u|}{\cancel{x^2}} = |u| \rightarrow 0$$

$$\lim_{(x,y) \rightarrow (0,0)} x \operatorname{Sen}\left(\frac{1}{y}\right) = 0$$

$$\left| x \operatorname{Sen}\left(\frac{1}{y}\right) \right| \leq |x| \cdot 1 = |x| \longrightarrow 0$$

$$\bullet \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} \quad \begin{array}{l} x = r \cos \theta \\ y = r \operatorname{Sen} \theta \end{array}$$

$$\left| \frac{x^2 y}{x^2 + y^2} \right| = \left| \frac{x \cdot xy}{x^2 + y^2} \right| = \frac{|x| |xy|}{x^2 + y^2} \cdot \frac{2}{2} = \frac{|x| \cdot 2 |xy|}{(x^2 + y^2) \cdot 2} \leq \frac{|x| (\cancel{x^2 + y^2})}{2 (\cancel{x^2 + y^2})} \rightarrow 0$$

$$\frac{x^2 y}{x^2 + y^2} = \frac{r^2 \cos^2 \theta \cdot r \operatorname{Sen} \theta}{r^2 \cos^2 \theta + r^2 \operatorname{Sen}^2 \theta} = \frac{r^3 \cos^2 \theta \operatorname{Sen} \theta}{r^2} = r \cos^2 \theta \operatorname{Sen} \theta$$

luego $|\cos^2 \theta| \leq 1$ y $|\operatorname{Sen} \theta| \leq 1$

$$\Rightarrow 0 \leq r \cos^2 \theta \operatorname{Sen} \theta \leq r$$

$$\Rightarrow \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} = \lim_{r \rightarrow 0} r \cos^2 \theta \operatorname{Sen} \theta = 0$$

$$\bullet \lim_{(x,y) \rightarrow (0,0)} \frac{x^2}{x^2 + y^2}$$

$$\begin{array}{l} x = r \cos \theta \\ y = r \operatorname{Sen} \theta \end{array} \quad \frac{r^2 \cos^2 \theta}{r^2 \cos^2 \theta + r^2 \operatorname{Sen}^2 \theta} = \frac{r^2 \cos^2 \theta}{r^2} = \cos^2 \theta$$

Como $|\cos^2 \theta| \leq 1 \Rightarrow \cos^2 \theta \leq r \cos^2 \theta \leq r$
 $\Rightarrow \cos^2 \theta \leq r$

así, $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2}{x^2 + y^2} = \lim_{r \rightarrow 0} \cos^2 \theta = \cos^2 \theta \Rightarrow$ el límite no existe, pues el resultado depende de θ

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + xy + y^2}{x^2 + y^2}$$

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\frac{r^2 \cos^2 \theta + r^2 \cos \theta \sin \theta + r^2 \sin^2 \theta}{r^2}$$

$$= 1 + \cos \theta \sin \theta$$

lungo $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + xy + y^2}{x^2 + y^2} = \lim_{r \rightarrow 0} (1 + \cos \theta \sin \theta) = 1 + \cos \theta \sin \theta$

lungo il limite dipende da $\theta \Rightarrow$ no esiste

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^5 + y^4 - 3x^3y + 2x^2 + 2y^2}{x^2 + y^2} = 2$$

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\frac{r^5 \cos^5 \theta + r^4 \sin^4 \theta - 3r^4 \cos^3 \theta \sin \theta + 2r^2}{r^2}$$

$$= r^3 \cos^5 \theta + r^2 \sin^4 \theta - 3r^2 \cos^3 \theta \sin \theta + 2$$

$$|\cos^n(\theta)| \leq 1, \forall n \in \mathbb{N}$$

$$|\sin^n(\theta)| \leq 1, \forall n \in \mathbb{N}$$

$$-r^3 - 3r^2 + 2 \leq r^3 \cos^5 \theta + r^2 \sin^4 \theta - 3r^2 \cos^3 \theta \sin \theta + 2 \leq r^3 + r^2 + 3r^2 + 2$$

$$\Leftrightarrow -r^3 - 3r^2 + 2 \leq r^3 \cos^5 \theta + r^2 \sin^4 \theta - 3r^2 \cos^3 \theta \sin \theta + 2 \leq r^3 + 4r^2 + 2$$

lungo $\lim_{r \rightarrow 0} (-r^3 - 3r^2 + 2) = \lim_{r \rightarrow 0} (r^3 + 4r^2 + 2) = 2$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2} \quad / \quad \text{no limits}$$

$$x=0 \Rightarrow \lim_{\substack{(x,y) \rightarrow (0,0) \\ x=0}} \frac{0 - y^2}{0 + y^2} = -1$$

$$y=0 \Rightarrow \lim_{\substack{(x,y) \rightarrow (0,0) \\ y=0}} \frac{x^2 - 0}{x^2 + 0} = 1$$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3 + x^2 + xy^2 + y^2}{x^2 + y^2}$$

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\frac{r^3 \cos^3 \theta + r^2 \cos^2 \theta + r^3 \cos \theta \sin^2 \theta + r^2 \sin^2 \theta}{r^2}$$

$$= \frac{r^3 (\cos^3 \theta + \cos \theta \sin^2 \theta) + r^2}{r^2}$$

$$= r (\cos^3 \theta + \cos \theta \sin^2 \theta) + 1$$

$$-r - r + 1 \leq r \cos^3 \theta + r \cos \theta \sin^2 \theta + 1 \leq r + r + 1$$

$$\lim_{r \rightarrow 0} -2r + 1 = 1 = \lim_{r \rightarrow 0} 2r + 1$$

$$\Rightarrow \lim_{(x,y) \rightarrow (0,0)} \frac{x^3 + x^2 + xy^2 + y^2}{x^2 + y^2} = 1$$

$$\frac{x^3 + x^2 + xy^2 + y^2}{x^2 + y^2} = \frac{x^2(x+1) + (x+1)y^2}{(x^2 + y^2)}$$

$$= \frac{(x+1)(\cancel{x^2 + y^2})}{(\cancel{x^2 + y^2})} = (x+1)$$

luego

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3 + x^2 + xy^2 + y^2}{x^2 + y^2} = \lim_{(x,y) \rightarrow (0,0)} (x+1) = 1$$

determinar c , tal que

$$g(x,y) = \begin{cases} \frac{x^3 + xy^2 + 2x^2 + 2y^2}{x^2 + y^2} & (x,y) \neq (0,0) \\ c & (x,y) = (0,0) \end{cases}$$

Sea continuo

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3 + xy^2 + 2x^2 + 2y^2}{x^2 + y^2} = \lim_{(x,y) \rightarrow (0,0)} \frac{x^2(x+2) + y^2(x+2)}{x^2 + y^2}$$

$$= \lim_{(x,y) \rightarrow (0,0)} \frac{(x+2)(\cancel{x^2 + y^2})}{(\cancel{x^2 + y^2})}$$

$$= \lim_{(x,y) \rightarrow (0,0)} (x+2) = 2$$

luego $c=2$

