AB CLOUD QUANTUM RESEARCH · MONOGRAPH SERIES

VOLUME Th5

Th5. Block reduction

Th1–Th6: hermiticity, self-adjointness, certificates, the spectral Li formula, block reduction, the Guinand–Weil trace formula

Ishak Khamzatovich Isaev

Independent researcher, Nalchik, Kabardino-Balkarian Republic

Ishak Khamzatovich Isaev · version 1.2.0 · October 10, 2026

Th5. Block reduction

Th1–Th6: hermiticity, self-adjointness, certificates, the spectral Li formula, block reduction, the Guinand–Weil trace formula

Ishak Khamzatovich Isaev · Independent researcher, Nalchik, Kabardino-Balkarian Republic

AB Research Program · monograph series V1–V19 · том Th5 · версия 1.2.0 · October 10, 2026

Contents

Th5. Block reduction by symmetry **2**

Th5.1 Certificate and boundaries **2**

Th5. Block reduction by symmetry

Theorem Th5 (V9).

If [H, A] = [H, B] = 0 for the generators of the abelian group G of magnetic translations, then H is block-diagonal in the character decomposition of G and spec H = ⊔_χ spec(H|Π_χ ℂ^N).

Proof. *Commuting unitaries A, B are simultaneously diagonalizable; their joint eigenvalues are the characters χ: G → U(1); the spectral projectors Π_χ (polynomials in A, B) commute with H: [H, Π_χ] = 0, hence each Π_χ ℂ^N is invariant and H decomposes into the direct sum of restrictions. Corollary: eigenvalues are found in smaller blocks with guaranteed coverage. Numerically: the union of 6 blocks reproduces the full spectrum to 7.1e-15; vortices break the symmetry (commutators ε_g = O(1)) — the reduction applies only to the clean lattice and to symmetric vortex configurations.*

Th5.1 Certificate and boundaries

Certificate: test_th5_block_reduction.py compares the union of the block spectra with the full spectrum; threshold 1e-9, fact 7.1e-15. The value is twofold: (a) computational — the reduction speeds up scans (each block ~$L^2/6$ instead of L^2); (b) diagnostic — the magnitude ‖[H, A]‖ measures how far a vortex configuration is from symmetric and serves as a passport feature of a gate in V4.

Boundaries: for protocol (asymmetric) vortices the reduction does not apply — this is not a defect but the boundary of its domain; partially symmetric configurations give approximate invariant subspaces, and the quantification of this idea is not included in the package.




Fig. 1. Theorem Th5: four measurement dimensions — the metric, its behaviour across gates/instances, the control and the threshold (2×2 tile, 600 dpi, campaign verdict data).