

Syncopation affects free body-movement in musical groove

Maria A. G. Witek¹  · Tudor Popescu² · Eric F. Clarke³ · Mads Hansen⁴ ·
Ivana Konvalinka⁵ · Morten L. Kringelbach^{1,6} · Peter Vuust¹

Received: 6 May 2016 / Accepted: 17 November 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract One of the most immediate and overt ways in which people respond to music is by moving their bodies to the beat. However, the extent to which the rhythmic complexity of groove—specifically its syncopation—contributes to how people spontaneously move to music is largely unexplored. Here, we measured free movements in hand and torso while participants listened to drum-breaks with various degrees of syncopation. We found that drum-breaks with medium degrees of syncopation were associated with the same amount of acceleration and synchronisation as low degrees of syncopation. Participants who enjoyed dancing made more complex movements

than those who did not enjoy dancing. While for all participants hand movements accelerated more and were more complex, torso movements were more synchronised to the beat. Overall, movements were mostly synchronised to the main beat and half-beat level, depending on the body-part. We demonstrate that while people do not move or synchronise much to rhythms with high syncopation when dancing spontaneously to music, the relationship between rhythmic complexity and synchronisation is less linear than in simple finger-tapping studies.

Keywords Music · Movement · Rhythm · Syncopation · Groove · Synchronisation

Electronic supplementary material The online version of this article (doi:10.1007/s00221-016-4855-6) contains supplementary material, which is available to authorized users.

✉ Maria A. G. Witek
maria.witek@clin.au.dk

- ¹ Center for Music in the Brain, Department of Clinical Medicine, Aarhus University & The Royal Academy of Music Aarhus/Aalborg, Noerrebrogade 44, 8000 Aarhus, Denmark
- ² Dresden Music Cognition Lab, Dresden University of Technology, August-Bebel Str 20, 01219 Dresden, Germany
- ³ Faculty of Music, University of Oxford, St Aldate's, Oxford OX1 1DB, UK
- ⁴ Department of Psychology and Behavioural Sciences, Aarhus University, Bartholins Alle 9, 8000 Aarhus, Denmark
- ⁵ Section for Cognitive Systems, Department of Applied Mathematics and Computer Science, Technical University of Denmark, Richard Petersens Plads, 2800 Lyngby, Denmark
- ⁶ Department of Psychiatry, Warneford Hospital, University of Oxford, Warneford Lane, Oxford OX3 7JX, UK

Syncopation affects free body-movement in musical groove

The body is posited as the central site for cognitive processes, according to the embodied approach to perception (Clark 2008; Leman 2007; Varela et al. 1991). For music, it is hard to imagine a context more obvious than dance as an illustration of that claim: through body-movement, music is actively perceived and physically embodied in ways that correspond to the structure of the music. An important concept in music and dance is groove. It is defined as a musical quality associated with a pleasurable desire for body-movement (Janata et al. 2012; Madison 2006; Madison et al. 2011; Stupacher et al. 2013) and music that is associated with groove is thus a particularly suitable venue for the investigation of music-directed dance. In groove research, syncopation—a form of rhythmic complexity—has been shown to be an important predictor. While some studies have shown that there is an inverted U-shaped relationship between syncopation and ratings of pleasurable wanting to

move (Sioros et al. 2014; Witek et al. 2014b), other studies show negative linear relationships between syncopation and synchronised finger-tapping (Fitch and Rosenfeld 2007; Ladinig et al. 2009; Song et al. 2013; Witek et al. 2014a). But what is the relationship between syncopation in groove and body-movement in dance? To what extent does syncopation affect how people move spontaneously to music? Here, we report on a study in which we used motion-capture to record free body-movement in response to rhythmic patterns with varying degrees of syncopation, measuring the acceleration, synchronisation, complexity, and periodicity of body-movements. Our main aim was to investigate the effects of (a) syncopation in groove and (b) musical background on these movement properties.

The progress in embodied music perception (Leman 2007) has been facilitated by the development of sophisticated motion-capture systems for measuring body-movements in space and time. Motion-capture research in music demonstrates that dancers map their body-movements to the structural (Leman and Naveda 2010), acoustic (Burger et al. 2013b), and emotional characteristics (Burger et al. 2013a; Saarikallio et al. 2013) of music. Some studies have specifically focused on rhythm and metre (Toiviainen et al. 2010), rhythm being broadly defined as a pattern of discrete durations organised into groups (Clarke 1999; Fraisse 1963, 1982), while metre is understood as the temporal framework in relation to which rhythm is perceived (Jones 2009; Lerdahl and Jackendoff 1983; London 2012). Formally, metre is organised hierarchically (Lerdahl and Jackendoff 1983; Longuet-Higgins and Lee 1984; Temperley 2010), with different periodicities corresponding to differing levels of metric salience. At the most fundamental level, metre perception requires beat perception, i.e. the temporal perception of events relative to a regular interval, or 'beat' (Honing 2012). Evidence from motion-capture confirms that the most prominent metric periodicity, or beat, expressed in dance is the tactus (the main pulse, or main beat), and that different body-parts can embody different metrical beat levels, simultaneously (Toiviainen et al. 2010). Furthermore, a number of individual factors can affect the propensity towards and nature of music-directed dance, such as biomechanical properties (Todd et al. 2007), personality (Luck et al. 2010), genre preference (Van Dyck et al. 2010), and dance experience (Witek et al. 2014b).

Dance involves sensorimotor synchronisation, which is the rhythmic coordination of perception and action allowing us to synchronise our movements to a beat. It relies fundamentally on mechanisms of prediction and adaptation (Konvalinka et al. 2010; Repp and Su 2013): in order to align one's movements with a periodic external referent, one must be able to anticipate the referent's temporal progression and continuously correct for movement errors.

The most significant musical property to affect sensorimotor synchronisation is rhythmic complexity (Chen et al. 2008; Repp and Su 2013; Witek et al. 2014a). One of the most studied forms of rhythmic complexity in music is syncopation, which is defined as a rhythmic event that violates listeners' metric expectations (Fitch and Rosenfeld 2007; Ladinig et al. 2009; Longuet-Higgins and Lee 1984; Margulis and Beatty 2008; Song et al. 2013; Temperley 2010; Witek et al. 2014a). Using a neural model of oscillations based on dynamic attending theory, Large et al. (2015) confirmed that it is possible to perceive a pulse frequency that is not acoustically present but only metrically implied in a rhythm (i.e. a syncopated rhythm). Furthermore, they found that the more spectral energy at the beat frequency, the less variable was the finger-tapping to the beat. Similar to this, Snyder and Krumhansl (Snyder and Krumhansl 2001) found that tapping to ragtime (i.e. a jazz piano style defined largely by its syncopatedness) was significantly affected by removing the left-hand part, which usually features a predictable alternating bass pattern. Without the regular beat of the left-hand part, tapping to the syncopated patterns of the right-hand part increased the tapping variability and asynchrony and caused more off-beat tapping. Consistent with these studies, a number of experiments have shown a negative linear relationship between syncopation and synchronised finger-tapping (Fitch and Rosenfeld 2007; Ladinig et al. 2009; Song et al. 2013; Witek et al. 2014a): more syncopated rhythms challenge listeners' metric predictions and thus reduce the capacity to synchronise. This effect is modulated by musical training, with musicians performing better than non-musicians (Palmer and Krumhansl 1990).

Rhythmic complexity has been shown to be an important structural aspect of groove. In a previous experiment (Witek et al. 2014b), we have shown that medium degrees of syncopation elicit the most desire to move and the most pleasure while listening to drum-breaks, in particular for participants who enjoy dancing. This suggests that listeners prefer a balance between predictability and complexity in music and that there is an inverted U-shaped relationship between syncopation and the desire for body-movement (Sioros et al. 2014). This inverted U-shape—also called Wundt curve (Wundt 1874)—between complexity and preference has previously been suggested to reflect aesthetic appreciation of art more broadly (Berlyne 1971). Finding this function in response to syncopation in groove, however, seems to contradict evidence regarding the relationship between syncopation and synchronised body-movement: tapping studies show a negative linear relationship between syncopation and synchronisation (Fitch and Rosenfeld 2007; Ladinig et al. 2009; Song et al. 2013), while studies involving groove suggest an inverted U-shaped relationship between syncopation and desire for

synchronised body-movement (Sioros et al. 2014; Witek et al. 2014b). One study showed that the more a piece of music is perceived as having groove the more participants synchronise their finger-tapping to the beat (Janata et al. 2012). However, no study has until now specifically measured the synchronisation of free body-movements to syncopated music. While finger-tapping is an embodied activity, moving the core body (i.e. torso) is more embodied in the sense that it involves moving the whole body and affords a different affective experience.

The study reported here used motion-capture to investigate the relationship between three degrees of syncopation and free body-movements in the hands and torsos of participants with varying levels of musical training and who enjoyed dancing and listening to groove to differing extents. In the light of earlier findings of an inverted U-shaped relationship between syncopation and wanting to move (Witek et al. 2014b), we hypothesised that participants would move more to drum-breaks with Medium syncopation, compared to Low and High. Since dancing to rhythmic music is generally synchronised, we tested the extent to which people synchronised to rhythms with different levels of syncopation and hypothesised that Medium syncopation would increase movement synchronisation. However, since there are no explicit instructions to synchronise when dancing spontaneously to music (this is only implied by the music's rhythmicity), we did not ask people to synchronise. As such, this is not primarily a synchronisation study, but a study of free body-movement which measures synchronisation alongside a number of other movement properties. One of these other properties was the spatial dimensionality of movements, which we measured to investigate whether syncopation affects the complexity of people's movements and whether there is a relationship between temporal synchronisation and spatial complexity in moving to groove. The possible effect of syncopation on the most prominent periodicities in movements was also investigated, on the basis that people might change the metrical level to which they synchronise according to the degree of syncopation in the drum-break. Since musicians demonstrate improved synchronisation with syncopated rhythms (Witek et al. 2014a), we hypothesised that participants with more musical training would be more synchronised than participants with less musical training. Finally, since our previous study (Witek et al. 2014b) found that dance experience influenced ratings of pleasure and wanting to move to groove, and since personality and genre preferences have been found to affect body-movements in dance (Luck et al. 2010; Van Dyck et al. 2010), we measured the effects of dancing experience and groove familiarity.

Method

Participants

Twenty-six right-handed Danish-speaking participants (7 females, 19 males) aged between 21 and 40 (Mean = 25.13, SD = 5.34) were recruited in Aarhus, Denmark, through adverts and opportunity sampling. Informed consent was obtained from all individual participants included in the study. Participants received a small payment for their participation. Before taking part in the motion-capture experiment, they participated in an unpublished fMRI study. One participant (a musician who enjoyed neither groove nor dancing) was excluded from the analysis, since his data showed that he barely moved during the experiment and he reported that he found the study highly unnatural and uncomfortable. Thus, the final number of participants was 25.

Participants completed a demographics questionnaire about musical training, dance experience, and groove familiarity, which showed that most participants enjoyed dancing and frequently danced to music, and liked and frequently listened to groove. All 25 participants' data were included in the analysis, regardless of their score on these measures. As expected, there were strong correlations between liking and frequency of listening to groove ($r = .642$, $p < .001$), and between liking and frequency of dancing ($r = .625$, $p < .001$). Thus, a principal component analysis (PCA) was run on each of these two group measures, and the resulting first principal components were used as measures of groove familiarity and dance experience. Participants were then defined as belonging to groups based on their position on the spectrum of groove familiarity and dance experience, i.e. whether they were below or above the mean. This yielded the following numbers in each category: groove-enjoyer $n = 12$; non-groove-enjoyer $n = 13$; dance-enjoyer $n = 12$; non-dance-enjoyer $n = 13$. It should be noted that the distribution of responses below and above the mean was not clearly discrete, i.e. there were some responses close to the mean. Therefore, these should not be thought of as strict categories but rather groups of participants with more or less experience with groove and dance. Importantly, groove familiarity did not correlate with dance experience ($r = .243$, $p = .231$), suggesting that these categories represented different attributes. The classification of musical training was categorical: those with six or more years of musical training were categorised as musicians ($n = 12$), and those with five or fewer years of training were categorised as non-musicians ($n = 13$).

Stimuli

Participants heard 15 synthesised drum-breaks with varying degrees of syncopation programmed using a synthesised drum-kit in GarageBand 5.1 (Apple, Inc.). Degree of syncopation was coded according to a modified version of Longuet-Higgins and Lee's (1984) syncopation index, which not only considered the notes' metric positions but also the polyphonic context and instrumental configuration (see the corrected supporting information in (Witek et al. 2014b, 2015) for a detailed description of the index). We categorised the drum-breaks into three syncopation levels: Low, Medium, and High (5 examples in each category). The drum-breaks were chosen from the larger pool of 50 funk drum-breaks used in a previous online survey (Witek et al. 2014b). They consisted of 16-s repeated 2-bar drum-kit patterns of bass drum, snare drum and hihat, at 120 bpm. The hihat was sounded on every semi-quaver. Degree of syncopation correlated marginally significantly with total number of onsets ($r = -.498$, $p < .059$). The supplementary material to this paper shows notational transcripts of the 15 drum-breaks used in the study, with Low, Medium, and High syncopation (and individual syncopation values).

Apparatus

Participants' body-movements were measured using the accelerometer data generated by the Wii remote controllers of the wireless motion-sensor videogame console Wii (Nintendo, Inc). One 'Wiimote' was strapped to the lower back (torso) and another held in the right hand. Previous research has found that movements in these body-parts are the most consistently entrained to the main pulse during spontaneous dance to music (Toiviainen et al. 2010). Acceleration in all three Euclidian dimensions (two horizontal, one vertical) was recorded simultaneously for both Wiimotes via WiiDataCapture 2.1 (Burger and Toiviainen 2013) on a MacBook laptop running OSX 10.4. Data were recorded with a sampling frequency of 100 samples per second. The drum-breaks were presented to participants over loudspeakers, and the order of presentation was counterbalanced and logged using Presentation (Neurobehavioral Systems, Inc.) on a Sony Vaio laptop.

Procedure

The room in which the motion-capture took place was large and mostly empty, apart from a table by one wall on which the laptops controlling the experiment and the speakers were placed, facing the centre of the room. A strip of tape was stuck to the floor 130 cm from the speakers, and participants were told not to move beyond the tape during the

experiment, in order to keep the loudness level as stable as possible across participants. The loudness of the drum-breaks as projected from the speakers was held constant for all participants and was measured at the tape location, using a DAWE D-1422C digital sound-level metre, with a 30–135 dB sound range. The mean loudness for the drum-breaks was 75 dB (SD = .97 dB).

The motion-capture recording was triggered by a one-second-long loud beep, and 16 s later the first drum-break was presented. In this way, the motion-capture recording was time-locked to the stimuli. During the first 16 s of silence, the experimenter left the room, thus leaving the participant to perform the motion-capture task completely alone. Participants were asked to *move freely* to the drum-breaks. Note that there were no explicit instructions to synchronise to the music. Each drum-break was heard twice during the course of the experiment, in randomised order. All drum-breaks were heard once before the repetitions were presented. The drum-breaks followed on from one another continuously, with no gap between them. This prevented any disruption to the regularity of the metre, which was continuous across drum-breaks, avoiding a loss of flow in body-movements. After the last drum-break, the experimenter entered the room and stopped the recording. Finally, subjects completed the demographics questionnaire.

Analysis

The first 4 s of the motion-capture data for each drum-break (i.e. corresponding to the first two bars of the drum-break) were excluded, to allow participants to adjust their movements to the new rhythmic pattern. Four properties of movement were extracted from the data: acceleration, synchronisation index, movement complexity, and periodicity. Data were analysed using the MoCap Toolbox 1.4 (Burger and Toiviainen 2013) for MATLAB (MathWorks, Inc.) and PASW 19.0 (IBM, Inc.).

To address how much people moved to drum-breaks with different degrees of syncopation, we measured movement acceleration, defined by the mean acceleration in each trial, averaged over the three Euclidean dimensions (as normalised using the 'mcnorm' function in the MoCap toolbox) and the two repetitions of each drum-break.

To investigate whether participants synchronised to the drum-breaks and whether the synchronisation depended on the degree of syncopation, we calculated the synchronisation index (SI) between the movements and the main pulse of the drum-break, representing the variance of the relative phase (Konvalinka et al. 2010). The three dimensions of motion-capture data were first reduced using principal component analysis (PCA). Subsequent processing and analyses were performed on the first principal component, since it involved the majority of movements' variance. In

order to measure synchronisation, we needed to focus on a single periodicity to compare between movements and music. Following Phillips-Silver et al. (2010), we filtered the data using fast Fourier transform (FFT) with a Gaussian kernel distribution centred on the frequency of the main pulse (i.e. main beat) of the stimuli (2 Hz, given the tempo of 120 bpm and a sampling frequency of 100 samples per second), including 10% of variability on either side of the centre frequency. The filtered data were then Hilbert-transformed, and instantaneous phase data were estimated. A variable representing the tactus in the stimulus (a single value at each sample representing the 2-Hz frequency, i.e. the main pulse) was also processed with the Hilbert transform. *SI*s were then calculated from the relative phase between the movement data and the tactus. The *SI* is based on the relative phase of the signals, and is output as a unitless value between 0 and 1, representing the absence of synchrony and perfect synchrony, respectively (Skewes et al. 2015; Tognoli et al. 2007). The formula in Eq. (1) was used:

$$SI = \frac{1}{N} \left| \sum_{n=1}^N e^{i(\theta_1(t_n) - \theta_2(t_n))} \right| \quad (1)$$

where N is the number of time samples in each trial, and θ_1 and θ_2 are the respective phases of the movement signal and the tactus. All *SI*s were averaged across the two stimulus repetitions.

To investigate the relationship between complexity of movements and syncopation, we computed the dimensionality of the motion-capture data, following Saarikallio et al. (2013). We used PCA on the mocap data to determine the cumulative variance of the principal components. According to Bennett (1969), there is an inverse relationship between the variance in interpoint distances within a hypersphere and the dimensionality of a hypersphere. Thus, in our analyses, a high proportion of cumulative variance represents low dimensionality, i.e. low complexity. Since the first two components of the hand and torso data contained on average about 80 and 86% of the cumulative variance, respectively, we used the cumulative variance of the second component (CumVarPC2) as our measure of dimensionality (i.e. the sum of the variances of the first and second components). Values were averaged across stimulus repetitions.

For acceleration, synchronisation, and dimensionality, we performed separate linear and quadratic regressions, using the individual syncopation values for each drum-break as the predictor and average acceleration, synchronisation and dimensionality across subjects as the output variables. For these measures, we also performed separate mixed-model $3 \times 2 \times 2 \times 2 \times 2$ ANOVAs, with

syncopation (Low, Medium, and High) and body-part (hand, torso) as the within-subjects variables and musical training, groove familiarity, and dance experience as between-subjects variables. Correlations between the measures were also calculated. Due to the large number of variables, we only report main effects and two-way interactions.

For the purpose of obtaining participants' movement periodicities, the maximum amplitude (i.e. the most prominent periodicity during dancing to each drum-break) was extracted, using the 'mcperiod' function (autocorrelation) in the MoCap toolbox. In order to do this, the Euclidean dimensions of the data were reduced with PCA. Since it did not make sense to average across the two repetitions (periodicities are not linear), we chose to use data from the first repetition only, unless no period was detected, in which case we used data from the second repetition. With this procedure, there were only two trials with no period across all participants (these were both in the hand). We then categorised the periods of each trial according to their nearest metric periodicity, with a 20% tolerance. If the period of a trial fell outside of this range, the trial was left empty, indicating that the most prominent periodicity for this trial was not metric. Then, metric periodicity trials were grouped into the three levels of syncopation (Low, Medium, and High). It was found that movements were periodic at three metric levels, the eighth-note level, the fourth-note (beat) level, and the half-note level. Using SPSS, we thus performed a $2 \times 3 \times 3$ repeated measures ANOVA, with body-part (hand and torso), syncopation level (Low, Medium, and High), and metric levels (eighth, fourth, and half-note) as independent variables. To reduce the number of variables, we did not include any between-subjects variables in this analysis.

Results

For amount of movement, the regression showed a significant negative linear relationship between acceleration in the hand and syncopation $R^2 = .585$ (adjusted $R^2 = .552$), $F(1,14) = 18.31$, $p < .001$. For the hip, there was a significant negative quadratic relationship between acceleration and syncopation $R^2 = .660$ (adjusted $R^2 = .604$), $F(1,14) = 11.67$, $p = .002$. Table 1 reports the coefficients. In the mixed ANOVA, there were significant main effects of syncopation $F(2,34) = 22.60$, $p < .001$, $\eta_p^2 = .571$, and body-part $F(1,17) = 123.37$, $p < .001$, $\eta_p^2 = .879$, and an interaction between syncopation and body-part $F(2,34) = 8.23$, $p < .001$, $\eta_p^2 = .326$. There were no effects of musical background. Bonferroni-corrected tests for simple effects (Table 2) showed that movements in the hand accelerated more than in the torso, and that for both body-parts,

Table 1 Regression coefficients. Syncopation = linear predictor. Syncopation² = quadratic predictor

Predictor	Hand		Torso	
	<i>B</i> (β)	<i>SEB</i>	<i>B</i> (β)	<i>SEB</i>
<i>Acceleration</i>				
Constant	28.87**	.741	8.31**	.403
Syncopation	-.070 (-.765)**	.016	–	–
Syncopation ²	–	–	-.001 (-1.574)**	<.001
<i>Synchronisation</i>				
Constant	.790**	.021	.781**	.028
Syncopation	-.003 (-.885)**	<.001	–	–
Syncopation ²	–	–	<.001 (-2.066)**	<.001
<i>Complexity</i>				
Constant	–	–	.879**	.002
Syncopation	–	–	–	–
Syncopation ²	–	–	<.001 (-2.328)**	<.001

** $p < .001$ **Table 2** Test of simple effects for interaction between syncopation (Low, Medium, and High) and body-part (hand and torso) on movement acceleration

Contrasts	<i>p</i>
<i>Hand</i>	
Low versus medium	>.999
Low versus high	<.001**
Medium versus high	<.001**
<i>Torso</i>	
Low versus medium	>.999
Low versus high	.002*
Medium versus high	.001*
<i>Low</i>	
Hand versus torso	<.001**
<i>Medium</i>	
Hand versus torso	<.001**
<i>High</i>	
Hand versus torso	<.001**

Corrected for multiple comparisons using the Bonferroni method

* $p < .005$, ** $p < .001$

movements accelerated more during Low and Medium syncopation, compared to High. There was no significant difference between Low and Medium. Figure 1 shows (a) the means for the three levels in both body-parts, as well as the regressed acceleration responses for each participant and the average across participants in (b) hand and (c) torso.


**Fig. 1** Effect of **a** syncopation (Low, Medium, and High) on movement acceleration in hand and torso and regression curves for **b** hand and **c** torso. Coloured lines represent curves for individual participants, bold black curve for averaged responses across participants. Error bars standard error. ** $p < .001$

Table 3 Test of simple effects for interaction of syncopation (Low, Medium, and High) with musical training (musicians, non-musicians) on movement synchronisation

Contrasts	<i>p</i>
<i>Musician</i>	
Low versus medium	>.999
Low versus high	.002*
Medium versus high	.003*
<i>Non-musician</i>	
Low versus medium	.195
Low versus high	<.001**
Medium versus high	<.001**
<i>Low</i>	
Musician versus non-musician	.384
<i>Medium</i>	
Musician versus non-musician	.877
<i>High</i>	
Musician versus non-musician	.054^

Corrected for multiple comparisons with the Bonferroni method

* $p < .005$, ** $p < .001$, ^ marginally significant

The relationship between syncopation and synchronisation to the tactus, i.e. main beat of the drum-breaks, was according to our regression analysis negatively linear for the hand $R^2 = .784$ (adjusted $R^2 = .767$), $F(1,14) = 47.08$, $p < .001$, and negatively quadratic for the torso $R^2 = .824$ (adjusted $R^2 = .795$), $F(1,14) = 28.11$, $p < .001$. Coefficients are reported in Table 1. The mixed ANOVA showed main effects of body-part $F(1,17) = 10.96$, $p = .004$, $\eta_p^2 = .392$, and syncopation $F(2,34) = 52.77$, $p < .001$, $\eta_p^2 = .756$, and a significant interaction between syncopation and musical training $F(2,34) = 4.91$, $p = .013$, $\eta_p^2 = .224$. Bonferroni-corrected tests of simple effects showed that the torso had a higher *SI* than the hand ($p = .004$). Table 3 and Fig. 2a show that syncopation affected the synchronisation of musicians and non-musicians similarly, with significant differences between Low and High, and between Medium and High, but not between Low and Medium. Musicians were also marginally significantly more synchronised during High syncopation compared to non-musicians. Figures 2b and 2c show the regression curves for individuals as well as averaged over all participants, for hand and torso, respectively. Furthermore, there was a significant between-subjects effect of dance experience $F(1,17) = 5.40$, $p = .033$, $\eta_p^2 = .241$, suggesting that non-dance-enjoyers (Mean = .742, SE = .026) were more synchronised than dance-enjoyers (Mean = .657, SE = .026). There were significant negative correlations between synchronisation and acceleration for hand $r = -.542$, $p = .006$, and torso $r = -.442$, $p = .027$.




Fig. 2 Effect of **a** syncopation (Low, Medium, and High) and musical training on synchronisation indices and regression curves for **b** hand and **c** torso. Coloured lines represent curves for individual participants, bold black curve for averaged responses across participants. Error bars standard error. * $p < .05$, ^marginally significant

For movement complexity, we found no significant regression for the hand. However, for the torso, there was a significant negative quadratic effect of syncopation $R^2 = .730$ (adjusted $R^2 = .685$), $F(1,14) = 16.20$, $p < .001$. See Table 1 for coefficients. There was no effect of syncopation on movement dimensionality in the mixed ANOVA, but a marginally significant effect of body-part




Fig. 3 Proportion of **a** trials with metric periodicities in hand and torso and **b** proportion of trials with metric periodicities in Low, Medium, and High syncopation conditions. Error bars standard error. * $p < .05$, ** $p < .001$

$F(1,18) = 3.907$, $p = .065$, $\eta_p^2 = .187$, suggesting a trend for hand movements to be more complex than torso movements. Furthermore, dance-enjoyers made more complex movements than non-dance-enjoyers, indicated by dance-enjoyers' lower CumVarPC2 (mean = .856, SE = .006) compared to non-dance-enjoyers (mean = .877, SE = .006) $F(1,17) = 6.86$, $p = .018$, $\eta_p^2 = .287$. There were also strong positive correlations between *SI* and dimensionality (averaged across body-parts) for both musicians ($r = .991$, $p < .001$) and non-musicians ($r = .974$, $p < .001$), and for dance-enjoyers ($r = .987$, $p < .001$) and non-dance-enjoyers ($r = .982$, $p < .001$).

For periodicities, we found significant main effects for metric level $F(1,48) = 27.52$, $p < .001$, $\eta_p^2 = .534$, and syncopation level $F(1,48) = 6.56$, $p = .003$, $\eta_p^2 = .215$, and also significant interactions between body-part and metric level $F(2,48) = 18.49$, $p < .001$, $\eta_p^2 = .435$ and between syncopation and metric level $F(4,96) = 3.26$, $p = .015$, $\eta_p^2 = .120$. Bonferroni-corrected simple comparisons

showed that there were significant differences between all metric levels in the hand (all $p < .05$), but only between the eighth and fourth-note beat levels and the fourth and half-note levels in the torso ($p < .01$). Figure 3a depicts the means and standard errors, which show that for the hand, the half-note level was the most prominent periodicity, followed by the fourth-note beat level and the eighth-note level, respectively. For the torso, the beat level was the most prominent periodicity compared to the other periodicities, which were not significantly different. There were also significant differences between the hand and torso at every metric level (eighth-note $p = .029$, fourth-note $p < .001$, half-note $p < .001$). The interaction between syncopation and metric levels (Fig. 3b) amounted mostly to differences within the syncopation levels, although there was also a significant difference between syncopation levels at the beat frequency, specifically participants' movements were more periodic during Medium syncopation compared to High syncopation ($p = .022$). Within each syncopation level, there were significant differences between the eighth and fourth-note beat levels, and between the eighth and half-note levels (all $p < .001$), but not between the fourth and half-note levels.

Discussion

In this study, we found that participants move the least to drum patterns with High levels of syncopation. For the hand, there was a negative linear relationship between syncopation and acceleration, while for the torso, this negative relationship was U-shaped. However, when we compared Low and Medium syncopation for both body-parts, there were no significant differences. Medium syncopation, i.e. a balance between predictability and complexity, has previously been associated with the most wanting to move and the most pleasure in groove (Witek et al. 2014b). Thus, our data suggest that there is some correspondence between ratings of desire to move and actual body-movement in groove (people neither want to nor actually move much to High syncopation), but that intermediate complexity drum-breaks elicit as much movement as rhythmically simple drum-breaks.

A similar pattern was found for synchronisation. In the hand, the regression analyses showed a negative linear relationship, suggesting that the more syncopated the rhythm, the less synchronised the movements. In the torso, this negative relationship was U-shaped. As in movement acceleration, we found no differences between Low and Medium synchronisation for either body-part. Previous research has shown that when asked to synchronise to a rhythm, participants perform worse the more complex the rhythm is

(Fitch and Rosenfeld 2007; Ladinig et al. 2009; Song et al. 2013). Here, we did not ask participants to synchronise but to move freely, and it seems that the relationship between syncopation and spontaneous synchronisation, while not quite inversely U-shaped, is less clearly negatively linear. Thus, it may be that the desire to move and pleasure specific to groove interacts with the more general detrimental effect of syncopation on sensorimotor synchronisation. In other words, because Medium syncopation motivates body-movement in groove, it may be improving synchronisation, not enough to produce a full-blown inverted U-shaped effect, but by an amount that makes it indistinguishable from Low syncopation. This interpretation would be in accordance with Janata et al.'s (2012) study, in which it was found that the more a piece of music was perceived as having groove, the more participants synchronised their tapping to the beat.

But why should intermediately syncopated rhythms invite synchronised behaviour? While the link between medium complexity and maximum pleasure in art has been addressed for several decades (Berlyne 1971) and rhythmic entrainment is increasingly being studied as a musical emotion induction mechanism (Trost and Vuilleumier 2013), it is less clear why intermediate levels of complexity would afford maximal synchronisation. When addressing the link between synchronisation and pleasure, prediction and dopamine are often mentioned, because of their role in both motor and reward functioning (Keitz et al. 2003). The idea is that the embodiment of successful temporal predictions afforded by synchronising to a beat stimulates the reward network in the brain and dopamine encodes these predictions. As for the link between intermediate complexity and sensorimotor synchronisation, it has been argued that syncopation, particularly at medium degrees, invites synchronised body-movements in groove because it opens up 'gaps' in the rhythmic structure that the body feels compelled to 'fill in' by moving to the beat (Witek 2016). In other words, syncopation affords a situation in groove where dancers can use their own body-movements to emphasise the beat. This may explain why the participants in our study synchronised more to drum-breaks with medium syncopation than previous finger-tapping studies would predict. Rather than making it more difficult to synchronise, the particular complexity of syncopation in groove makes it more *motivating* to synchronise. In other words, the desire to move elicited by the syncopations counterbalances the difficulty of synchronising to syncopated rhythms.

There were few differences between musicians and non-musicians in our study. Both groups synchronised more to Low and Medium syncopation, compared to High syncopation. At High syncopation, musicians were marginally more synchronised than non-musicians, in accordance with previous research (Chen et al. 2008; Fitch and Rosenfeld

2007; Repp and Su 2013; Witek et al. 2014a). It is possible that with stricter criteria for our groups, this improvement for musicians would be more robust and that there would be significant differences in the other syncopation conditions as well. There were also strong positive correlations between synchronisation and movement dimensionality for both musical training groups, suggesting that the more spatially complex are their movements, the less are they temporally synchronised. This was also the case for dance-enjoyers and non-dance-enjoyers. Furthermore, dance-enjoyers made significantly more complex movements and were marginally less synchronised compared to non-dance-enjoyers. While professional dancers may be better at synchronising when specifically asked to do so (Miura et al. 2013), our study suggests that when moving spontaneously to music, people who enjoy dancing are more concerned with making spatially complex movements, and this is at the expense of their overall temporal synchronisation. We found no effects of groove familiarity, suggesting that liking and frequently listening to groove does not affect the way people move to music.

There were a number of differences between the two body-parts whose movements we measured. We found that the hand accelerated more, i.e. it moved more than the torso. This can be explained by the greater spatial and temporal degrees of freedom for movement in the hand, due to its smaller size and weight, and greater mobility at the end of a limb. We also found that the torso was more synchronised than the hand. As noted by Toiviainen et al. (2010), the torso has a higher moment of inertia than the arm, due to its greater mass. We thus suggest that the torso is better suited to synchronising to the beat than the hand, while the hand is better suited to making complex non-beat-related movements. This was suggested by the finding that hand movements were marginally more complex than torso movements.

The hand and torso also differed in how strongly they entrained to different metrical levels in the drum-breaks, although overall movements were most strongly entrained to the beat, half-note, and eighth-note levels, as expected (Toiviainen et al. 2010), and only occasionally to non-metric periodicities. For the hand, the half-note was the most prominent periodicity, followed by the beat level and the eighth-note level in that order. In the torso, the beat level was the most prominent periodicity, followed by the half-note and the eighth-note, which were similarly prominent. There were also significant differences between the two body-parts at each metric level; the eighth-note and beat levels were more prominent in the torso, while the half-note level was most prominent in the hand. Toiviainen et al. (2010) used biomechanical properties to explain different periodicity patterns found in the hand compared to the torso, proposing that body-parts with greater inertia

necessarily have longer periodicities in movement. This is opposite to our finding that the torso, with much greater inertia, produced much more prevalent eighth-note periodicities than the hand. Therefore, it may be that inertia cannot fully explain the relationship between body-parts and metric periodicities. We suggest that it could be the movements in the legs that determined the metric levels embodied in the torsos of our participants. While we did not measure leg movements, they may have affected the spatio-temporal patterns in the torso. Specifically, we suspect that participants moved to the eighth-note level in the legs by bending the knees up and down to each eighth note, perhaps in alternating fashion between the right and left leg. This movement would be mirrored in the torso and may explain why we see relatively frequent eighth-note periodicities in torso movements. Furthermore, our drum-breaks were instrumentally sparser than the stimuli used in Toiviainen et al.'s (2010) study, who used a full-band 12-bar blues progression, and the difference in our findings may be due to the difference in stimuli.

Between the different levels of syncopation, there were few differences in movement periodicities, except that Medium syncopation was associated with more beat-level periodicities than High syncopation. This is consistent with the finding that participants were more synchronised to Medium than to High syncopation.

To conclude, this study has shown that during free body-movement to music, syncopation relates to the amount of movement, degree of synchronisation, complexity, and the prevalence of different periodicities in hand and torso movements in ways that interact with musical training and dance experience. Contrary to previous finger-tapping research, we found that during free body-movement to music, the relationship between rhythmic complexity—here in the form of syncopation—and sensorimotor synchronisation is not clearly negatively linear, with no difference in synchronisation between Low and Medium degrees of syncopation. Our findings further contribute to an understanding of what it is about music that motivates spontaneous motor behaviour, and how musical structure shapes our body-movements. Our study emphasises that when moving freely to music, dancers do not just embody the music by synchronising to the beat but also respond by changing their movements' spatial complexity. As an instance of embodied cognition (Clark 2008; Leman 2007; Varela et al. 1991), we demonstrate how musical properties are spontaneously expressed in body-movement, and how musical dance involves the coordination of perceptual, cognitive, and sensorimotor capacities.

Acknowledgements Center for Music in the Brain is funded by the Danish National Research Foundation (DNRF117). IK is funded by the Danish Council for Independent Research—Technology and

Production Sciences. Financial support to TP was in part provided by the Zukunftskonzept at TU Dresden funded by the Exzellenzinitiative of the Deutsche Forschungsgemeinschaft. MLK is supported by the ERC Consolidator Grant: CAREGIVING (n. 615539). We thank Kristian Nymoen and Birgitta Burger for help with the analysis.

Compliance with ethical standards

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the Institutional and/or National Research Committee and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

References

- Bennett R (1969) The intrinsic dimensionality of signal collections. *IEEE Trans Inf Theory* 15:517–525
- Berlyne DE (1971) *Aesthetics and psychobiology*. Appleton-Century-Crofts, East Norwalk
- Burger B, Toiviainen P (2013) MoCap Toolbox: a matlab toolbox for computational analysis of movement data. In: *Proceedings of 10th sound and music computing conference (SMC)*, Stockholm, Sweden Berlin: Logos Verlag
- Burger B, Saarikallio S, Luck G, Thompson MR, Toiviainen P (2013a) Relationships between perceived emotions in music and music-induced movement. *Music Percept* 30:517–533
- Burger B, Thompson MR, Luck G, Saarikallio S, Toiviainen P (2013b) Influences of rhythm- and timbre-related musical features on characteristics of music-induced movement. *Front Psychol* 4:183
- Chen JL, Zatorre RJ, Penhune VB (2008) Moving on time: brain network for auditory-motor synchronization is modulated by rhythm complexity and musical training. *J Cogn Neurosci* 20:226–239
- Clark A (2008) *Supersizing the mind: embodiment, action, and cognitive extension*. Oxford University Press, New York
- Clarke EF (1999) Rhythm and timing in music. In: Deutsch D (ed) *The psychology of music*, 2nd edn. Academic Press, New York
- Fitch WT, Rosenfeld AJ (2007) Perception and production of syncopated rhythms. *Music Percept* 25:43–58
- Fraisse P (1963) *The psychology of time*. Harper & Row, New York
- Fraisse P (1982) Rhythm and tempo. In: Deutsch D (ed) *The psychology of music*, 1st edn. Academic Press, New York
- Honing H (2012) Without it no music: beat induction as a fundamental musical trait. *Ann N Y Acad Sci* 1252:85–91. doi:10.1111/j.1749-6632.2011.06402.x
- Janata P, Tomic ST, Haberman JM (2012) Sensorimotor coupling in music and the psychology of the groove. *J Exp Psychol Gen* 141:54–75. doi:10.1037/a0024208
- Jones MR (2009) Musical time. In: Hallam S, Cross I, Thaut M (eds) *The Oxford handbook of music psychology*. Oxford University Press, New York, pp 81–92
- Keitz M, Martin-Soelch C, Leenders K (2003) Reward processing in the brain: a prerequisite for movement preparation? *Neural Plas* 10:121–128
- Konvalinka I, Vuust P, Roepstorff A, Frith CD (2010) Follow you, follow me: continuous mutual prediction and adaptation in joint tapping. *Q J Exp Psychol* 63:2220–2230
- Ladinig O, Honing H, Haden G, Winkler I (2009) Probing attentive and preattentive emergent meter in adult listeners without extensive musical training. *Music Percept* 26:377–386
- Large EW, Herrera JA, Velasco MJ (2015) Neural networks for beat perception in musical rhythm. *Front Syst Neurosci* 9:159

- Leman M (2007) Embodied music cognition and mediation technology. MIT Press, Cambridge
- Leman M, Naveda L (2010) Basic gestures as spatiotemporal reference frames for repetitive dance/music patterns in Samba and Charleston. *Music Percept* 28:71–91
- Lerdahl F, Jackendoff R (1983) A generative theory of tonal music. MIT Press, Cambridge
- London J (2012) Hearing in time. Oxford University Press, New York
- Longuet-Higgins HC, Lee C (1984) The rhythmic interpretation of monophonic music. *Music Percept* 1:424–440
- Luck G, Saarikallio S, Burger B, Thompson MR, Toiviainen P (2010) Effects of the Big Five and musical genre on music-induced movement. *J Res Pers* 44:714–720. doi:[10.1016/j.jrp.2010.10.001](https://doi.org/10.1016/j.jrp.2010.10.001)
- Madison G (2006) Experiencing groove induced by music: consistency and phenomenology. *Music Percept* 24:201–208
- Madison G, Gouyon F, Ullén F, Hörnström K (2011) Modeling the tendency for music to induce movement in humans: first correlations with low-level audio descriptors across music genres. *J Exp Psychol Hum Percept Perform* 37:1578–1594. doi:[10.1037/a0024323](https://doi.org/10.1037/a0024323)
- Margulis EH, Beatty AP (2008) Musical style, psychoaesthetics, and prospects for entropy as an analytical tool. *Comput Music J* 32:64–78
- Miura A, Kudo K, Ohtsuki T, Kanehisa H, Nakazawa K (2013) Relationship between muscle cocontraction and proficiency in whole-body sensorimotor synchronization: a comparison study of street dancers and nondancers. *Mot Control* 17:18–33
- Palmer C, Krumhansl CL (1990) Mental representation for musical meter. *J Exp Psychol Hum Percept Perform* 16:728–741
- Phillips-Silver J, Aktipis AC, Bryant G (2010) The ecology of entrainment: foundations of coordinated rhythmic movement. *Music Percept* 28:3–14
- Repp BH, Su Y-H (2013) Sensorimotor synchronization: a review of recent research (2006–2012). *Psychon Bull Rev* 20:403–452
- Saarikallio S, Luck G, Burger B, Thompson M, Toiviainen P (2013) Dance moves reflect current affective state illustrative of approach–avoidance motivation. *Psychol Aesthet Creat Arts* 7:296
- Sioros G, Miron M, Davies M, Gouyon F, Madison G (2014) Syncopation creates the sensation of groove in synthesized music examples. *Front Psychol* 5:1036
- Skewes JC, Skewes L, Michael J, Konvalinka I (2015) Synchronised and complementary coordination mechanisms in an asymmetric joint aiming task. *Exp Brain Res* 233:551–565
- Snyder JS, Krumhansl CL (2001) Tapping to ragtime: cues to pulse finding. *Music Percept* 18:455–489
- Song C, Simpson AJ, Harte CA, Pearce MT, Sandler MB (2013) Syncopation and the score. *PLoS ONE* 8:e74692
- Stupacher J, Hove MJ, Novembre G, Schütz-Bosbach S, Keller PE (2013) Musical groove modulates motor cortex excitability: a TMS investigation. *Brain Cogn* 82:127–136
- Temperley D (2010) Modeling common-practice rhythm. *Music Percept* 27:355–376
- Todd NPMA, Cousins R, Lee CS (2007) The contribution of anthropometric factors to individual differences in the perception of rhythm. *Empir Musicol Rev* 2:13
- Tognoli E, Lagarde J, DeGuzman GC, Kelso JS (2007) The phi complex as a neuromarker of human social coordination. *Proc Natl Acad Sci* 104:8190–8195
- Toiviainen P, Luck G, Thompson MR (2010) Embodied meter: hierarchical eigenmodes in music-induced movement. *Music Percept* 28:59–70
- Trost W, Vuilleumier P (2013) Rhythmic entrainment as a mechanism for emotion induction by music: a neurophysiological perspective. In: Cochrane T, Fantini B, Scherer KR (eds) The emotional power of music: multidisciplinary perspectives on musical arousal, expression, and social control. Oxford University Press, New York, pp 213–225
- Van Dyck E, Moelants D, Demey M, Coussement P, Deweppe A, Leman M (2010) The impact of the basis drum on body movement in spontaneous dance. In: Proceedings of the 11th international conference on music perception and cognition, Seattle, Washington, USA, 23–27.8.2010
- Varela FJ, Thompson E, Rosch E (1991) The embodied mind. Cognitive science and human experience. MIT Press, Cambridge
- Witek MAG (2016) Filling in: syncopation, pleasure and distributed embodiment in groove. *Music Anal*. doi:[10.1111/musa.12082](https://doi.org/10.1111/musa.12082)
- Witek MAG, Clarke EF, Kringelbach ML, Vuust P (2014a) Effects of polyphonic context, instrumentation and metric location on syncopation in music. *Music Percept* 32:201–217
- Witek MAG, Clarke EF, Wallentin M, Kringelbach ML, Vuust P (2014b) Syncopation, body-movement and pleasure in groove music. *PLoS ONE* 9:e94446. doi:[10.1371/journal.pone.0094446](https://doi.org/10.1371/journal.pone.0094446)
- Witek MA, Clarke EF, Wallentin M, Kringelbach ML, Vuust P (2015) Correction: syncopation, body-movement and pleasure in groove music. *PLoS ONE* 10
- Wundt W (1874) Grundzüge der physiologischen psychologie. Englemann, Leipzig