

# Nucleation and Crystallization of the Metastable Hard Sphere Fluid - Master Colloquium

Wilkin Wöhler,  
Albert-Ludwigs-Universität Freiburg

May 13, 2021

# Content


- Introduction
  - Metastable hard sphere fluid
  - Memory effects in non-stationary processes
  - Nucleation rate discrepancy
- Simulation scheme and testing
- Cluster growth rate
- Nucleation rate
  - Definition
  - Estimation
  - Comparison
- Memory kernel shape analysis

# The Hard Sphere Fluid

- Simplest model of a fluid

$$V(r_{ij}) = \begin{cases} \infty & r_{ij} \leq \sigma \\ 0 & r_{ij} > \sigma \end{cases}$$

- Experimentally and theoretically studied
- Liquid-Solid phase transition
  - $\eta_{RCP} \approx 64\%$ ,  $\eta_{HCP} \approx 74\%$
  - Entropy increase




**Fig.:** Carnahan-Starling[1], Almarza[2] and equilibrium equation of state

# Nucleation and Crystallization - Phase Transitions

- Technological and scientific importance
- Commonly described by classical nucleation theory
- Other Approaches
  - Modifying the free energy landscape
  - Markovian reaction coordinates
  - Memory effects


# Classical Nucleation Theory (CNT)

- Spherical clusters
- Coarse grained observable  $R$
- $\Delta F = 4\pi R \gamma_{fs} - \frac{4\pi}{3} R^3 \rho \Delta |\mu|$
- Markovian fluctuations
- Nucleation rate  $\kappa \propto \exp\left(-\frac{\Delta F}{k_B T}\right)$



# Classical Nucleation Theory (CNT)

- Spherical clusters
- Coarse grained observable  $R$
- $\Delta F = 4\pi R \gamma_{fs} - \frac{4\pi}{3} R^3 \rho \Delta |\mu|$
- Markovian fluctuations
- Nucleation rate  $\kappa \propto \exp\left(-\frac{\Delta F}{k_B T}\right)$



# Nucleation and Crystallization - Phase Transitions

- Technological and scientific importance
- Commonly described by classical nucleation theory
  - qualitatively correct
  - quantitatively not accurate
- Other Approaches
  - Modifying the free energy landscape
  - Markovian reaction coordinates
  - Including memory effects

# Memory Effects Nucleation Processes

Earlier work in the group:

- Memory effects observed in Lennard-Jones nucleation, Anja[3]
- Non-stationary generalized Langevin equation, Hugues[4]

$$\frac{dA_t}{dt} = \omega(t)A_t + \int_0^t K(\tau, t)A_\tau d\tau + \eta_{0,t}$$


- Non-Markovian kernel in the Lennard-Jones nucleation, Philipp P.[5]

This work:

- Memory kernel of hard sphere nucleation process

# Nucleation Rate Discrepancy

- Steeper decrease in simulations
- Possible differences:
  - 1 Hydrodynamics
  - 2 Sedimentation
  - 3 Polydispersity
  - 4 Measurement geometry
  - 5 ...



**Fig.: Experimental and theoretical nucleation rates [6–13]**

# Simulation Scheme and Testing


## Event driven molecular dynamics (EDMD)

- Discontinuous potential
- Resolve dynamics
- Based on analytical-
  - ballistic trajectory
  - collision time
  - collision outcome
- Validated dynamics by diffusion
- Cell system
  - ⇒  $\mathcal{O}(N)$  calculation effort

# Simulation Scheme and Testing

## Event driven molecular dynamics (EDMD)

- Discontinuous potential
- Resolve dynamics
- Based on analytical-
  - ballistic trajectory
  - collision time
  - collision outcome
- Validated dynamics by diffusion
- Cell system  
 $\Rightarrow \mathcal{O}(N)$  calculation effort



**Fig.: Logarithmic plot of long time diffusion constant [14, 15]**

# Simulation Scheme and Testing

## Event driven molecular dynamics (EDMD)

- Discontinuous potential
- Resolve dynamics
- Based on analytical-
  - ballistic trajectory
  - collision time
  - collision outcome
- Validated dynamics by diffusion
- Cell system  
 $\Rightarrow \mathcal{O}(N)$  calculation effort




Fig.: CPU time per simulation step

# Choice of Data Production Parameters

- $\mathcal{O}(N)$  calculation effort

$$\Rightarrow \frac{\delta T_{\text{CPU}}}{\delta t_{\text{Sim.}}} \propto N \propto V$$

- Without initial induction time

$$\Rightarrow \langle \tau_{\text{Nucleation}} \rangle \propto \frac{1}{V}$$


- Time to nucleation

$$\Rightarrow \langle T_{\text{CPU}} \rangle \propto \frac{\delta T_{\text{CPU}}}{\delta t_{\text{Sim.}}} \cdot \langle \tau_{\text{Nucleation}} \rangle \propto \text{const.}$$


$\Rightarrow$  Large system sizes possible

# Cluster Growth - Constant Attachment Rate

- Constant attachment rate  $\Rightarrow R \propto t$
- Spherical growth  $\Rightarrow V \propto R^3 \propto t^3 \Rightarrow N(t) = c^3(t - t_0)^3$



**Fig.:** Third root of the number of particles in the largest cluster



**Fig.:** Comparison of growth rates in the constant attachment regime.

# Nucleation Time Distribution

Assume simulation geometry and  $\kappa(t) = \text{const.}$

- $N$  subvolumes of size  $V_{\text{box}}$
- $n(t)$  nucleated boxes

$$\langle \Delta n \rangle = (N - n(t)) V_{\text{box}} \kappa \Delta t$$

- In the continuous limit of  $N \rightarrow \infty$  with  $V_{\text{box}} \kappa = k$

$$\begin{aligned} &\Leftrightarrow x_s = \frac{n(t)}{N} = 1 - \exp(-t k) \\ &\Leftrightarrow \dot{x}_s = k \exp(-t k) \end{aligned}$$

⇒ Exponentially distributed nucleation times

# Nucleation Rate Estimator I - Maximum Log-Likelihood Estimator

- Probability Density Function of censored exponential distribution

$$p(t) = \begin{cases} k \exp(-t k) & t < T \\ \exp(-T k) & t \geq T \end{cases}$$

- Likelihood of measurement


$$\mathcal{L}(k) = \binom{N}{m} k^n \exp\left(-k \sum_{i=1}^n t_i\right) \exp(-T k)^m$$

- Necessary condition of maximum likelihood

$$\hat{k}^{-1} = \frac{1}{n} \left( \sum_{i=1}^n t_i + mT \right)$$


# Nucleation Rate Estimator II - Monte Carlo Uncertainty

- 1 Exponentially distributed random samples with  $k = \hat{k}$
- 2 Censoring at T
- 3 Calculate  $\hat{k}_{MC}$
- 4  $\sigma_{\hat{k}}$  estimated by  $\sigma_{\hat{k}_{MC}}$




# Nucleation Rate Estimator II - Monte Carlo Uncertainty

- 1 Exponentially distributed random samples with  $k = \hat{k}$
- 2 Censoring at T
- 3 Calculate  $\hat{k}_{MC}$
- 4  $\sigma_{\hat{k}}$  estimated by  $\sigma_{\hat{k}_{MC}}$



# Nucleation Rate Comparison



Simulations and experiments:

- Different rates
- Different slopes


**Fig.:** Experimental and theoretical nucleation rates from the literature[6–13]

# Comparison to Real World Experiments

Simulation		Experiment
Microscopic disjunct subvolumes		Macroscopic volume
$\tau_{\text{cryst.}} \ll \tau_{\text{nuc.}}$		$\tau_{\text{cryst.}} \not\ll \tau_{\text{nuc.}}$
$x_s \approx \frac{n(t)}{m} = 1 - \exp(-\kappa t)$		$x_s(t) = t^4 \frac{\kappa c^3}{4\rho_{\text{melt}}}, x_s \ll 1$

- $\kappa$  and  $c$  are measured
- Assume  $x_s(t^*) \stackrel{!}{=} 1/8 \Rightarrow t^* = \sqrt[4]{\frac{\rho_{\text{melt}}}{2\kappa c^3}}$

# Modified Nucleation Rate Comparison




Simulations and experiments:

- Different rates
- Similar slopes

**Fig.:** Experimental nucleation rates[6–10] alongside with  $(t^*)^{-1}$  calculated from simulation studies[11–13]


# Memory Kernel I - Largest Cluster Trajectories

Last but not least,



- $N = 16384$  particles
- $\eta = 54.0\%$
- $\tau_{ind.} \approx 400\delta t$
- $\tau_{cry.} \approx 150\delta t$

# Memory Kernel II -Shape Analysis





Shape analysis:

- Exclude Markovian contribution
- FWHM of crest as first estimate
- Best Gaussian fit as second estimate

# Memory Kernel II -Shape Analysis

- Constant width  $\sim \tau_{cry}$ .
- Variable amplitude





⇒ Possibly only memory depending on the arbitrary transition width is observed.

# Conclusion - Summary

- Constant attachment rate
- Estimate of nucleation rate density + statistical uncertainty  
    ⇒ Confirming the difference
- System geometries in simulation and experiment  
    ⇒ Possible accordance
- Memory kernel shape analysis  
    ⇒ Future research in other observables

# Thank you for your attention



# Bibliography I

-  N. F. Carnahan and K. E. Starling, "Equation of state for nonattracting rigid spheres", *The Journal of Chemical Physics* **51**, 635–636 (1969).
-  N. G. Almarza, "A cluster algorithm for Monte Carlo simulation at constant pressure", *Journal of Chemical Physics* **130**, 184106 (2009).
-  A. Kuhnhold et al., "Derivation of an exact, nonequilibrium framework for nucleation: Nucleation is a priori neither diffusive nor Markovian", *Physical Review E* **100**, 52140 (2019).
-  H. Meyer, "Generalized Langevin Equations and memory effects in non-equilibrium statistical physics", *Dissertation (Université du Luxembourg, Albert-Ludwigs-Universität Freiburg, 2020)*.
-  P. Pelagejcev, "Simulation study of non-Markovian effects in crystal nucleation", *Master Thesis (Albert-Ludwigs-Universität Freiburg, 2019)*.
-  J. L. Harland and W. van Megen, "Crystallization kinetics of suspensions of hard colloidal spheres", *Physical Review E* **55**, 3054–3067 (1997).

## Bibliography II

-  Y. He et al., "Dynamics of crystallization in hard-sphere suspensions", Physical Review E **54**, 5286–5297 (1996).
-  K. Schätzel and B. J. Ackerson, "Density fluctuations during crystallization of colloids", Physical Review E **48**, 3766–3777 (1993).
-  C. Sinn, A. Heymann, A. Stipp, and T. Palberg, "Solidification kinetics of hard-sphere colloidal suspensions", Trends in Colloid and Interface Science XV. Progress in Colloid and Polymer Science **118**, 266–275 (2001).
-  S. Auer and D. Frenkel, "Prediction of absolute crystal-nucleation rate in hard-sphere colloids", Nature **409**, 1020–1023 (2001).
-  L. Filion, M. Hermes, R. Ni, and M. Dijkstra, "Crystal nucleation of hard spheres using molecular dynamics, umbrella sampling, and forward flux sampling: A comparison of simulation techniques", Journal of Chemical Physics **133**, 244115 (2010).

## Bibliography III

-  G. Fiorucci, G. M. Coli, J. T. Padding, and M. Dijkstra, "The effect of hydrodynamics on the crystal nucleation of nearly hard spheres", *Journal of Chemical Physics* **152**, 064903 (2020).
-  T. Schilling, S. Dorosz, H. J. Schöpe, and G. Opletal, "Crystallization in suspensions of hard spheres: A Monte Carlo and molecular dynamics simulation study", *Journal of Physics Condensed Matter* **23**, 194120 (2011).
-  S. Pieprzyk et al., "Thermodynamic and dynamical properties of the hard sphere system revisited by molecular dynamics simulation", *Physical Chemistry Chemical Physics* **21**, 6886–6899 (2019).
-  D. M. Heyes, M. J. Cass, J. G. Powles, and W. A. Evans, "Self-diffusion coefficient of the hard-sphere fluid: System size dependence and empirical correlations", *Journal of Physical Chemistry B* **111**, 1455–1464 (2007).