

Microbial Ecology and Biogeography

— OF THE —

Southern Ocean

David Wilkins

Submitted in fulfillment of the requirements for the Degree of Doctor of Philosophy.

SCHOOL OF BIOTECHNOLOGY AND BIOMOLECULAR SCIENCES
UNIVERSITY OF NEW SOUTH WALES, SYDNEY

April 2013

Contents

List of Figures	ix
List of Tables	xi
List of Acronyms	xiv
1 Introduction	1
Physical oceanography of the Southern Ocean	1
Fronts and zones	1
Water masses and circulation	2
Effect of climate change	3
Microbial ecology of the Southern Ocean	3
Bacteria	4
Alphaproteobacteria	4
Roseobacter clade	4
SAR11	5
SAR116	6
Betaproteobacteria	6
Gammaproteobacteria	6
SAR86	6
OMG group	7
Ant4D3	7
GSO-EOSA-1	7
Deltaproteobacteria	8
CFB	8
Cyanobacteria	9
Verrucomicrobia	9
Other bacteria	9
Archaea	10
Viriplankton	11
Project aims	11
Aim 1	11
Aim 2	11
Aim 3	11
Aim 4	11
2 MINSPEC	13
Abstract	13
Introduction	13
Metagenomic analysis of microbial assemblages	13
The maximum parsimony approach	15
Methods	15
Implementation of MINSPEC	15
Validation of MINSPEC	16
Results	17

Discussion	17
Conclusions	17
3 The Polar Front	21
Abstract	21
Introduction	21
Methods	22
Sampling and metagenomic sequencing	22
Phylogenetic analysis of metagenomic data	25
BLAST comparison to RefSeq database	25
OTU abundances and variance between zones	25
Fragment recruitment to verify Operational Taxonomic Unit (OTU) identification	25
Additional samples to test “polynya hypothesis”	25
Functional analysis of metagenomic data	26
BLAST comparison to KEGG database	26
Analysis of functional potential	26
Taxonomic decomposition	26
Results	27
Metagenomic sequencing	27
Phylogenetic analysis of metagenomic data	27
Fragment recruitment to verify OTU identification	31
Additional samples to test alternative “polynya hypothesis”	31
Functional analysis of metagenomic data	31
Discussion	31
Taxonomic groups differentiating the zones	31
GSO-EOSA-1	31
Ammonia-oxidizing Crenarchaeota	35
Cyanobacteria	35
SAR11 and SAR116 clades	36
Bacteroidetes	36
Rhodobacterales	37
Alteromonadales	37
Verrucomicrobia	37
Functional capacities differentiating the zones	37
Conclusions: Biogeographic role of the Polar Front	40
4 Deep water formation	41
Abstract	41
Methods	41
Results and Discussion	43
5 The advection effect	45
Abstract	45
Introduction	45
Methods	46
Sampling	46
DNA extraction	47
Sequencing	48
Taxonomic assignment	48
Physicochemical and spatial distances	49
Generation of advection distance matrix	49
Ordination of distance matrices and comparison to water masses	50
Testing of advection effect	50
Results	50
Sequencing and taxonomic assignment	50
Environment and distance effects	54

Testing the advection effect	56
Testing advection effect mechanisms	58
Discussion	59
Future work	60
References	61

List of Figures

1.1	Major fronts and water masses of the Southern Ocean	2
2.1	Results of MINSPEC validation	18
3.1	Map showing sites of seawater samples used in the Polar Front study	24
3.2	Rank-abundance curves for OTUs in each zone and size fraction	27
3.3	Contribution of OTUs to variance between the North and South zones	29
3.4	Tree of GSO-EOSA-1 related 16S rRNA genes	34
3.5	Read recruitment to reference genomes	34
3.6	Taxonomic decomposition of KEGG modules	38
4.1	Map showing sites of preliminary Antarctic Bottom Water (AABW) samples	42
4.2	Non-Metric Multidimensional Scaling (nMDS) of AABW, North Zone (NZ) and South Zone (SZ) samples	44
5.1	Map showing sites of samples used in the advection study	47
5.2	OTU assignments in the advection study.	54
5.3	nMDS of advective distances between samples.	55
5.4	nMDS of advective distances between samples.	55
5.5	dbRDA ordination of relationship between environment and community.	57
5.6	nMDS of advective distances between samples.	58
5.7	Advection effect at different taxonomic resolutions	59

List of Tables

2.1 Examples of spurious OTU identifications	14
3.1 Details of samples used in Polar Front study	23
3.2 Additional samples used to test polynya hypothesis	26
3.3 Twenty most abundant OTUs	28
3.4 Highest-contributing OTUs to the difference between the North and South zones	30
3.5 Contributions of KEGG modules to variance between the North and South zones	32
3.6 Contributions of KEGG ortholog groups to variance between the North and South zones	33
4.1 AABW samples used in the preliminary analysis	41
4.2 Twenty most abundant OTUs in preliminary AABW samples	43
5.1 Full sample data for advection study	52
5.1 (cont.) Full sample data for advection study.	53
5.2 Correlations between dbRDA axes and physicochemical variables	56

List of Acronyms

GAAS Genome relative Abundance and Average Size.

AABW Antarctic Bottom Water.

AAD Australian Antarctic Division.

AAP Aerobic Anoxygenic Phototrophic.

ACC Antarctic Circumpolar Current.

ANOSIM Analysis of Similarities.

AOA Ammonia-Oxidizing Archaea.

AZ Antarctic Zone.

CASO Climate of Antarctica and the Southern Ocean.

CEAMARC Collaborative East Antarctic Marine Census.

DFAA Dissolved Free Amino Acids.

DMSP dimethylsulfoniopropionate.

DOC Dissolved Organic Carbon.

DOM Dissolved Organic Matter.

FISH Fluorescence *In Situ* Hybridization.

GOS Global Ocean Sampling expedition.

GSO-EOSA-1 Gammaproteobacterial Sulfur Oxidizer-EOSA-1.

HMW High Molecular Weight.

HNLC High Nutrient, Low Chlorophyll.

KEGG Kyoto Encyclopedia of Genes and Genomes.

MGI Marine Group I Crenarchaeota.

MMPA methylmercaptopropionate.

nMDS Non-Metric Multidimensional Scaling.

NZ North Zone.

OTU Operational Taxonomic Unit.

PF Polar Front.

PFZ Polar Frontal Zone.

POM Particulate Organic Matter.

SAF Subantarctic Front.

SAZ Subantarctic Zone.

SIMPER Similarity Percentages.

SO Southern Ocean.

STF Subtropical Front.

SZ South Zone.

Acknowledgements

Chapter 3

The Polar Front as a major biogeographic boundary in the Southern Ocean

Sections of this chapter have been previously published in Wilkins D., Lauro F. M., Williams T. J., De-Maere M. Z., Brown M. V., Hoffman J. M., Andrews-Pfannkoch C., McQuaid J. B., Riddle M. J., Rintoul S. R., and Cavicchioli R. (2012). Biogeographic partitioning of Southern Ocean microorganisms revealed by metagenomics. *Environmental Microbiology*.

Abstract

The Polar Front (PF) is a major oceanographic feature of the Southern Ocean (SO), dividing surface water masses with distinct physicochemical characteristics. To confirm and characterise the PF's role as a biogeographic barrier in the SO, sixteen metagenomic samples were collected on a latitudinal transect from waters near Hobart, Australia (44° S) to near the Mertz Glacier, Antarctica (67° S). The samples were shotgun sequenced, and the community composition and functional potential of the captured microorganisms were analysed. The SAR11 and SAR116 clades and the cyanobacterial genera *Prochlorococcus* and *Synechococcus* were strongly overrepresented north of the PF. Conversely, the Gammaproteobacterial Sulfur Oxidizer-EOSA-1 (GSO-EOSA-1) complex, the phyla Bacteroidetes and Verrucomicrobia and order Rhodobacterales were overrepresented in waters south of the PF. Functions enriched south of the PF included a range of transporters, sulphur reduction and histidine degradation to glutamate, while branched-chain amino acid transport, nucleic acid biosynthesis and methionine salvage were overrepresented north of the PF. The taxonomic and functional characteristics suggested a shift of primary production from cyanobacteria in the north to eukaryotic phytoplankton in the south, and reflected the different trophic states of the two regions. Overall, this study supported the PF as a defining biogeographic feature of the SO.

Introduction

The Southern Ocean (SO) consists of zones separated by circumpolar fronts, the locations of which vary with time and longitude (Whitworth, 1980; Orsi *et al.*, 1995; Sokolov and Rintoul, 2002). From north to south, the major fronts are the Subtropical Front (STF), the Subantarctic Front (SAF) and the Polar Front (PF) (Fig. 1.1). The SAF and PF are associated with the Antarctic Circumpolar Current (ACC), the world's largest ocean current and a defining oceanographic feature of the SO. Anthropogenic climate change may be driving the warming and freshening of the ACC (Böning *et al.*, 2008). It may also be shifting the ACC and associated fronts poleward; the mean path of the current has moved \sim 50 km south since the 1950s (Gille, 2002). If this migration continues, by 2100 it will have displaced a volume of water south of the ACC approximately equivalent to the Arctic Ocean (Fyfe and Saenko, 2005).

Climate change may also be affecting the zones separated by these fronts. The major zones of the SO are the Subantarctic Zone (SAZ), between the STF and SAF; the Polar Frontal Zone (PFZ), between the SAF and the PF; and the Antarctic Zone (AZ), between the PF and the Antarctic continent (Fig. 1.1). These zones have different physicochemical properties, such as density, salinity, temperature and nutrient concentrations (Sokolov and Rintoul, 2002), and the fronts represent stepwise transitions in these properties (Whitworth III and Nowlin Jr., 1987). However, as a consequence of climate change, waters on the poleward side of the ACC have become warmer and more saline, while those to the north cooler and fresher (Böning *et al.*, 2008).

The PF has been suggested to be a major biogeographic boundary in the distribution and abundance of both zooplankton (Chiba *et al.*, 2001; Hunt *et al.*, 2001; Esper and Zonneveld, 2002; Ward *et al.*, 2003) and bacterioplankton (Selje *et al.*, 2004; Abell and Bowman, 2005b; Giebel *et al.*, 2009; Weber and Deutsch, 2010). The effects of climate change on the physical oceanography of the SO may therefore have global ecological significance, particularly as the SO is a major site for sequestration of anthropogenic CO₂ (Sabine *et al.*, 2004; Mikaloff Fletcher *et al.*, 2006) through both physicochemical processes and the microbe-driven “biological pump” of CO₂ fixation (Thomalla *et al.*, 2011), potentially forming a feedback loop (Cox *et al.*, 2000). However, the microbial assemblages that inhabit the SO are generally poorly understood, and their diversity and functional capacity poorly characterised (reviewed in Murray and Grzymski, 2007). A community-level understanding is needed to predict the effects of a shifting ACC on the distribution, abundance and ecosystem functions of microorganisms.

Large scale metagenome surveys have not previously been performed in the SO. Compared to traditional environmental microbiological methods such as culture-based or DGGE surveys, metagenomic studies are able to offer both deeper (capture of sequence from rare taxa) and broader (larger sample sizes with greater statistical significance) data on the taxonomic makeup and functional capacities of marine microbial communities (e.g. Rusch *et al.*, 2007; Angly *et al.*, 2006; Dinsdale *et al.*, 2008). Metagenome sampling is also better suited to large-scale biogeographic studies, as it allows large numbers of samples to be taken across broad spatial and temporal scales with uniform sampling, storage and processing, compared to e.g. culture-based methods where technical replicability is more difficult.

This project is part of a metagenomic survey of the SO begun in the austral summer of 2006, based on the sampling design of the Global Ocean Sampling expedition (GOS) (Rusch *et al.*, 2007). The GOS sampling strategy involves size fractionation of plankton assemblages by passing seawater through a 20 µm prefilter then capturing planktonic biomass on sequential 3.0 µm, 0.8 µm and 0.1 µm filter membranes. As well as allowing deeper sequencing of metagenomes and thus better representation of low-abundance taxa (Rusch *et al.*, 2007), this approach allows comparison to other samples collected using the GOS methods (e.g. Brown *et al.*, 2012). This study will thus focus on the 0.1–20 µm fraction of the sampled microbial assemblages.

Methods

Sampling and metagenomic sequencing

Sampling¹ was conducted on board the RSV *Aurora Australis* during Australian Antarctic Division (AAD) cruise V3 Collaborative East Antarctic Marine Census (CEAMARC) / Climate of Antarctica and the Southern Ocean (CASO) from 13 December 2007–25 January 2008. This cruise occupied the SR3 latitudinal transect from Hobart, Australia (44° S) to the Mertz Glacier, Antarctica (67° S) within a longitudinal range of 140–150° E. Sixteen samples were obtained along almost the entire latitudinal range (Fig. 3.1).

A range of physicochemical data were recorded by integrated instruments on the RSV *Aurora Australis* including location, water column depth, water temperature, salinity, fluorescence and meteorological conditions (Table 3.1). These data were used to locate the PFZ based on a surface temperature gradient of ~1.35 °C across a distance of 45–65 km, placing the PF at approximately 59.70° S, consistent with previous descriptions (Moore *et al.*, 1999; Sokolov and Rintoul, 2002). Samples were accordingly grouped into “North” and “South” zones (Table 3.1). The North Zone (NZ) represents waters from the Subtropical, Subantarctic and PFZ regions, while the South Zone (SZ) represents the AZ.

¹Sampling was performed by Jeffrey M. Hoffman and Jeffrey B. McQuaid.

Table 3.1: Sampling time, location and physicochemical properties of samples used in this study. All data were retrieved from underway instruments aboard the RSV *Aurora Australis*.

Sample	Zone	Date	Latitude	Longitude	Water Column Depth (m)	Sample Depth (m)	Temperature (°C)	Salinity (PSU)	Fluorescence ($\mu\text{g L}^{-1}$)	Volume filtered (L)
346	North	20/12/2007	-59.31	142.59	4294	2	2.9	33.75	0.3	500
347	South	23/12/2007	-66.02	142.74	450	2	0.6	34.20	4.0	250
349	South	27/12/2007	-66.57	142.32	370	1.5	-1.3	34.40	2.3	250
351	South	28/12/2007	-66.56	143.43	823	1.5	-0.6	34.30	1.3	500
352	South	29/12/2007	-66.77	143.32	164	2.5	-0.8	34.30	3.1	500
353	South	30/12/2007	-67.05	144.68	180	2	-1.8	34.40	0.3	500
357	South	05/01/2008	-66.17	143.02	580	2	-0.4	34.15	2.5	500
358	South	09/01/2008	-64.30	150.03	3550	2	0	33.55	0.5	500
359	South	12/01/2008	-66.19	143.53	540	2	-0.2	34.21	2.5	500
360	South	13/01/2008	-66.58	141.02	316	2	-0.7	34.04	6.2	500
362	South	19/01/2008	-65.54	140.83	1064	2	0.7	32.20	0.5	500
363	North	22/01/2008	-60.00	141.31	4473	2	3.3	33.77	0.1	500
364	North	23/01/2008	-56.70	141.88	3693	2	4	33.70	0.5	500
366	North	24/01/2008	-52.02	144.14	3180	2	7.6	33.84	0.3	500
367	North	25/01/2008	-48.25	145.90	3490	2	11	34.43	0.2	500
368	North	26/01/2008	-44.72	145.78	3201	2	14.8	34.96	1.3	560




Figure 3.1: Sites of seawater samples used in this study. Squares indicate surface samples from the North Zone; crosses samples from the South Zone. The dashed line represents the Polar Front.

At each station, ~250–560 L of seawater was pumped from ~1.5–2.5 m below the sea surface into drums stored at ambient temperature on deck. Seawater samples were prefiltered through a 20 µm plankton net, then filtrand was captured on sequential 3.0 µm 0.8 µm and 0.1 µm 293 mm polyethersulfone membrane filters (Pall, Port Washington, USA), and immediately stored at –20 °C (Rusch *et al.*, 2007; Ng *et al.*, 2010).

DNA extraction² was performed at the J. Craig Venter Institute (Rockville, USA) as described in Rusch *et al.* (2007). Pyrosequencing was performed on a GS20 FLX Titanium instrument (Roche, Branford, USA) also at the J. Craig Venter Institute as described in Lauro *et al.* (2011). Duplicate reads and reads with many pyrosequencing errors were removed as described in Lauro *et al.* (2011).

²DNA extraction was performed by Cynthia Andrews-Pfannkoch and others at the J. Craig Venter Institute.

Phylogenetic analysis of metagenomic data

BLAST comparison to RefSeq database

A subset of the RefSeq microbial (bacterial and archaeal) genome database (release 41, retrieved 31 May 2010 from <ftp://ftp.ncbi.nih.gov/refseq/release/>) was prepared by excluding sequences with the words “shotgun”, “contig”, “partial”, “end” or “part” in their headers (following Angly *et al.*, 2009). Because this database was not expected to contain representative genomes for every species present, Operational Taxonomic Units (OTUs) in this study are defined by the best species match to this database, and may for example represent congeners.

The metagenomic reads from each sample were compared against this database using TBLASTX, with default parameters except for: E-value threshold 1.0×10^{-3} , cost to open gap 11, cost to extend gap 1, masking of query sequence by SEG masking with lookup table only. The outputs of all TBLASTX searches against RefSeq were processed by MINSPEC, and hits not belonging to the minimal OTU sets were removed.

OTU abundances and variance between zones

The relative OTU abundances for each sample were determined using the PERL script Genome relative Abundance and Average Size (GAAS) (Angly *et al.*, 2009). Briefly, GAAS estimates the relative abundance of OTUs from the number and quality of BLAST matches (“hits”) to each OTU, taking into account differences in genome size. GAAS was run with the default settings. To normalise for reads which did not yield acceptable hits, the relative abundances for each sample were scaled by that sample’s effective BLAST hit rate. An OTU profile was generated for each sample by encoding the scaled relative abundance of each OTU from each size fraction as a separate variable.

To test the hypothesis that the oceanic zones harbour significantly different communities, Analysis of Similarities (ANOSIM) with 999 permutations was performed on a standardised, log-transformed Bray-Curtis resemblance matrix of OTU profiles. Similarity Percentages (SIMPER) analysis was performed to identify the contribution of individual OTUs to differences between the zones. All statistical procedures were performed in PRIMER 6 as described by Clarke and Warwick (2001).

Fragment recruitment to verify OTU identification

To confirm that identified OTUs were truly present, and not resulting from hits to small regions of genomic identity, plots of read recruitment and coverage to reference genomes were generated using R and PERL. Six samples, three from the SZ and three from the NZ, were randomly selected and their 0.1 μm fractions used as sources of recruited reads. As shotgun sequencing of a metagenome should be essentially random, it was predicted that these plots would show fairly even coverage of the reference genomes. Genomes where all recruited reads were concentrated in a few small regions would be likely candidates for OTUs which were not present, but shared a small genomic region with an OTU which was present.

Additional samples to test “polynya hypothesis”

Because many samples from the SZ were taken in the region of the Mertz Glacier polynya, an alternative hypothesis for any difference observed between the SZ and NZ would be that the difference represented the effect of the polynya, rather than the PF. A set of samples collected³ from waters south of the PF on different voyages and in a different longitudinal range were used to test this hypothesis. Sampling, DNA extraction and sequencing were performed as described above, although only the 0.1–0.8 μm size fraction was processed. The resulting metagenomic reads were compared to RefSeq and weighted relative abundances calculated with GAAS as described above.

Two ANOSIM tests were performed on the standardised and log-transformed Bray-Curtis resemblance matrices of the samples’ OTU abundance profiles and similarly prepared matrices representing only the 0.1 μm fractions of the SZ and NZ samples. In the first ANOSIM test (“polynya hypothesis”), all samples from the Mertz Glacier polynya region (i.e. all SZ samples apart from 358) were grouped

³Collection was performed by Ricardo Cavicchioli, Federico M. Lauro and Mark V. Brown. TODO who else was on these voyages?

Table 3.2: Dates and locations of additional samples used to falsify the hypothesis that the Mertz Glacier polynya was responsible for the observed biogeographic partitioning.

Sample	Date	AAD Voyage Number	Latitude	Longitude
388	20/10/2008	V1 2008–09	−63.8090	115.1613
398	22/10/2008	V1 2008–09	−64.8018	112.3730
390	30/10/2008	V1 2008–09	−64.8152	80.7204
392	13/12/2008	V2 2008–09	−64.1826	76.4537
393	15/12/2008	V2 2008–09	−55.2573	74.2531

together, while the samples from the open ocean were placed in a separate group. In the second test (“PF hypothesis”), the additional samples were grouped with all the SZ samples.

Functional analysis of metagenomic data

BLAST comparison to KEGG database

In order to identify functional differences between the zones, the set of metagenomic reads from each sample was compared against the Kyoto Encyclopedia of Genes and Genomes (KEGG) GENES database (retrieved 2 July 2010 from <ftp://ftp.genome.jp/pub/kegg/genes/fasta/genes.pep>) with BLASTX, with default parameters except for: maximum number of database sequence alignments 10; E-value threshold 1.0×10^{-3} ; gap opening penalty 11; gap extension penalty 1; masking of query sequence by SEG masking for lookup table only.

Analysis of functional potential

Genes identified by BLASTX were aggregated to KEGG ortholog groups according to the KEGG Orthology schema (<ftp://ftp.genome.jp/pub/kegg/genes/ko>, retrieved Mar 29 2011), and ortholog group abundances calculated for each sample. Following Coleman and Chisholm (2010), a read was considered a hit to a given ortholog group if the top three hits for that read (or all hits if fewer than three total hits) were to genes from the same ortholog group, and had bit scores > 40 . If the bit score difference between any two top hits was greater than 30, only the hits above this difference were considered.

Ortholog group counts were then used to calculate the abundance of KEGG modules. Because many ortholog groups are members of more than one module, the abundance a_m of each module m was calculated as

$$a_m = \sum_{K=1}^n \frac{C_K}{M_K}$$

where n is the number of ortholog groups K belonging to module m , C_K is the number of hits to ortholog group K , and M_K is the total number of modules to which K belongs. To account for differences in sequencing depth between samples, module abundances were scaled to 500,000 reads per sample. To test the hypothesis that the NZ and SZ harbour significantly different functional potential, one-way ANOSIM with 999 permutations was performed as above on a standardised, log-transformed Bray-Curtis distance resemblance matrix of the module and ortholog group profiles. A functional profile was generated for each sample by summing the scaled abundances of each module from all size fractions, and SIMPER performed as above to identify modules which contributed highly to the variation in functional potential between the two zones.

Taxonomic decomposition

To link modules with a high contribution to variance or otherwise of interest to specific OTUs (“taxonomic decomposition”), a script was written which decomposed each module to its constituent ortholog groups, then each ortholog group to each representative gene sequence in the KEGG database. Because each KEGG sequence is associated with a particular organism, this allowed the abundance




Figure 3.2: Rank-abundance curves for OTUs identified in each zone and size fraction. The dominance of a given OTU is its relative abundance as a percentage of all identified OTUs. The x-axis is scaled logarithmically. Generated using PRIMER 6.

of each module to be decomposed into the relative contribution from each of these organisms. This allowed functional contributions to be putatively assigned to genera which were not identified in our taxonomic analysis, as the database included gene sequences for organisms for which a full genome was not available. To aid interpretation of these results, the taxonomic decompositions for some modules were aggregated to higher taxonomic ranks, such as class or family.

Results

Metagenomic sequencing

6.6 Gbp of 454 sequence data representing microorganisms in the size range 0.1–3.0 μm was obtained from 16 samples. After removal of low-quality reads, 454 sequencing yielded 157,507–597,689 reads per sample (mean 354,399) of lengths ranging from 100 to 606 bp (mean 378).


Phylogenetic analysis of metagenomic data

The proportion of reads in each sample which yielded hits to RefSeq ranged from 25–85% (mean 62%). The most abundant OTUs in each sample are given in Table 3.3 and a full list of OTU abundances in the supplementary material (PF-all-OTUs.csv). All samples and size fractions exhibited very low OTU evenness (Fig. 3.2).

Table 3.3: Relative abundances (as percentages) of the twenty most abundant OTUs identified in this study, in each zone and size fraction.

OTU	North			South		
	0.1 µm	0.8 µm	3.0 µm	0.1 µm	0.8 µm	3.0 µm
"Candidatus Pelagibacter ubique" HTCC1062	61.76	25.00	23.87	58.85	22.40	17.61
<i>Nitrosopumilus maritimus</i> SCM1	0.01996	0.01438	0.009508	1.076	1.309	1.210
"Candidatus Ruthia magnifica" str. Cm (<i>Calyptogena magnifica</i>)	0.6699	0.6458	0.5484	2.987	2.616	1.025
<i>Roseobacter</i> sp. OCh 114	0.3125	2.932	1.588	0.4477	3.994	2.657
<i>Synechococcus</i> sp. CC9902	0.1081	9.837	4.973	0.0007484	0.004156	0.09733
<i>Silicibacter pomeroyi</i> DSS-3	0.2578	2.286	1.154	0.3070	2.505	1.576
<i>Gramella forsetii</i> strain KT0803	0.2412	1.210	1.755	0.4993	2.347	1.890
"Candidatus Vesicomysoscius okutanii" strain HA	0.4634	0.4642	0.2078	1.970	1.807	0.2174
<i>Robiginitalea biformalata</i> strain HTCC2501	0.2751	1.099	1.297	0.4722	1.878	1.405
<i>Flavobacterium psychrophilum</i> strain JIP02/86	0.1718	0.8409	1.224	0.4316	1.960	1.598
<i>Synechococcus</i> sp. CC9311	0.03014	4.624	4.409	0.0007221	0.002778	0.02764
"Candidatus Puniceispirillum marinum" IMCC1322	0.6444	2.077	1.267	0.3586	1.377	0.7109
<i>Silicibacter</i> sp. TM1040	0.2274	1.652	0.8738	0.2709	1.803	1.233
<i>Jannaschia</i> sp. DFL-12	0.1776	1.378	0.7350	0.2443	1.692	0.8009
<i>Zunongwangia profunda</i> strain SM-A87	0.1522	0.7487	1.059	0.2968	1.410	1.204
<i>Colwellia</i> sp. 34H	0.02345	0.3636	2.736	0.05207	0.5140	1.041
<i>Coraliomargarita akajimensis</i> strain DSM 45221	0.03698	0.07573	0.1197	0.1154	1.543	1.680
<i>Jannaschia</i> sp. CCS1	0.1173	0.9344	0.4784	0.1711	1.230	0.8239
<i>Pseudoalteromonas atlantica</i> strain T6c	0.01251	0.4772	1.993	0.02270	0.4089	1.132
<i>Saccharophagus degradans</i> strain 2-40	0.06532	0.4325	0.5429	0.1289	1.072	0.8663
<i>Flavobacterium johnsoniae</i> strain UW101	0.08822	0.4220	0.6141	0.2034	0.9389	0.8578
<i>Capnocytophaga ochracea</i> strain DSM 7271	0.1143	0.4830	0.5399	0.2314	0.8815	0.6814
<i>Marinomonas</i> sp. MWYL1	0.03777	0.2529	0.3026	0.1514	1.300	0.7006
<i>Cellvibrio japonicus</i> strain Ueda107	0.05884	0.3080	0.3231	0.1155	0.9917	0.4713
<i>Marinobacter hydrocarbonoclasticus</i> VT8	0.04093	0.2889	0.3883	0.08418	0.7195	0.3848
<i>Pseudoalteromonas haloplanktis</i> strain TAC125	0.01389	0.2505	0.8896	0.03427	0.3561	0.6530
<i>Terediniibacter turnerae</i> strain T7901	0.05665	0.3051	0.3081	0.1138	0.9174	0.5127
<i>Acinetobacter baumannii</i> strain SDF	0.004886	0.007187	0.4073	0.006260	0.04218	1.459

ANOSIM analysis showed that the zones harbor significantly different microbial communities ($R = 0.451$, $p = 0.004$). SIMPER was performed in order to identify the contribution of individual OTUs to the difference between the NZ and SZ. The results for the highest contributors are provided in Table 3.4, and are graphically summarised for all OTUs in Fig. 3.3.



The SIMPER analysis found that no single OTU contributed more than 2.9% of variance and 74% of variance was contributed by OTUs with a contribution less than 1%. There was also a large difference in the contribution to variance of the three size fractions, with approximately 52% of all variance

Table 3.4: The thirty OTUs with the highest contributions to the difference between the NZ and SZ. Abundances are zonal averages and have been standardised and log-transformed. As each OTU on each size fraction was encoded as a separate variable in the SIMPER analysis, the size fraction is given after each OTU name.

OTU	Abundance South	Abundance North	Contribution to variance (%)
Synechococcus sp. CC9311 0.8 µm	0.00	1.08	2.88
Synechococcus sp. CC9902 0.8 µm	0.00	1.04	2.81
Synechococcus sp. CC9311 3.0 µm	0.01	0.98	2.59
Synechococcus sp. CC9902 3.0 µm	0.04	0.76	2.03
"Candidatus Pelagibacter ubique" HTCC1062 3.0 µm	1.97	2.40	1.97
"Candidatus Ruthia magnifica" str. Cm (<i>Calyptogena magnifica</i>) 0.1 µm	0.82	0.25	1.57
"Candidatus Ruthia magnifica" str. Cm (<i>Calyptogena magnifica</i>) 0.8 µm	0.34	0.66	1.32
"Candidatus Pelagibacter ubique" HTCC1062 0.8 µm	0.74	0.25	1.32
"Candidatus Vesicomyosocius okutanii" strain HA 0.1 µm	2.32	2.48	1.32
Coraliomargarita akajimensis strain DSM 45221 0.8 µm	0.62	0.18	1.20
Coraliomargarita akajimensis strain DSM 45221 3.0 µm	0.48	0.04	1.13
Roseobacter sp. OCh 114 0.8 µm	0.49	0.06	1.10
Pseudoalteromonas atlantica strain T6c 3.0 µm	1.01	0.81	1.08
"Candidatus Vesicomyosocius okutanii" strain HA 0.8 µm	0.38	0.54	1.08
Acinetobacter baumannii strain SDF 3.0 µm	0.57	0.19	1.04
Acinetobacter baumannii strain SDF 3.0 µm	0.45	0.18	0.95
Gramella forsetii strain KT0803 0.8 µm	0.72	0.43	0.94
Marinomonas sp. MWYL1 0.8 µm	0.46	0.11	0.92
Roseobacter sp. OCh 114 3.0 µm	0.76	0.54	0.91
Flavobacterium psychrophilum strain JIP02/86 0.8 µm	0.63	0.32	0.89
Silicibacter pomeroyi DSS-3 0.8 µm	0.75	0.69	0.86
Brachyspira hyodysenteriae strain WA1 3.0 µm	0.47	0.19	0.84
"Candidatus Ruthia magnifica" str. Cm (<i>Calyptogena magnifica</i>) 3.0 µm	0.34	0.21	0.82
Pseudoalteromonas haloplanktis strain TACI25 3.0 µm	0.22	0.33	0.77
Robiginitalea biformata strain HTCC2501 0.8 µm	0.61	0.40	0.74
Nitrosopumilus maritimus SCM1 0.1 µm	0.27	0.01	0.72
Gramella forsetii strain KT0803 3.0 µm	0.59	0.59	0.71
Lysinibacillus sphaericus strain C3-41 3.0 µm	0.29	0.02	0.71
Nitrosopumilus maritimus SCM1 0.8 µm	0.25	0.01	0.70
Silicibacter sp. TM1040 0.8 µm	0.59	0.55	0.69

contributed by OTUs from the 3.0 μm fraction, 37% by the 0.8 μm fraction, and 9% by the 0.1 μm fraction. Notably, OTUs within several taxonomic groups that had high contribution to variance covaried in their relative representation in the NZ and SZ. For example, Bacteroidetes and GSO-EOSA-1 representatives were on average more abundant in the SZ; while *Prochlorococcus* and *Synechococcus* species, SAR11 and SAR116 were on average more abundant in the NZ (Fig. 3.3). Some groups, such as the Alteromonadales, had variable relative representation depending on size fraction.

Fragment recruitment to verify OTU identification

As predicted, read recruitment to reference genomes produced fairly even coverage, especially in the high-abundance OTUs. This indicates that OTU identification was accurate and unlikely to be confounded by small regions of genomic identity. Fig. 3.5 gives examples of read recruitment plots to three high-abundance and three low-abundance OTUs.

Additional samples to test alternative “polynya hypothesis”

The ANOSIM analysis strongly supported the “PF hypothesis” ($R = 0.435, p = 0.002$), i.e. that the PF was primarily responsible for the difference observed between the NZ and SZ, over the “polynya hypothesis” ($R = 0.29, p = 0.005$) that the influence of the Mertz Glacier polynya was responsible. This also provides evidence that the PF effect is robust over a longitudinal range and over time.

Functional analysis of metagenomic data

ANOSIM analysis of the samples’ KEGG ortholog group and module profiles revealed that the zones had significantly different functional potential (ortholog group: $R = 0.642, p = 0.001$; module: $R = 0.871, p = 0.001$). SIMPER was performed on the profiles in order to identify the specific functional differences between the zones. The highest-contributing modules are given in Table 3.5, and a complete list in the supplementary material (PF-modules-SIMPER.csv). The highest-contributing ortholog groups are given in Table 3.6, and a complete list in the supplementary material (PF-ortholog-groups-SIMPER.csv). No single ortholog group or module contributed more than 2.2% of the variance, indicating a complex and diverse pattern of functional differences. There was a strong trend for ortholog groups and modules with higher contributions to variance to be overrepresented in the NZ in the 3.0 μm fraction but the SZ in the smaller fractions, indicating that the functional diversity of each zone was strongly segregated by size fraction.

Discussion

Taxonomic groups differentiating the zones

GSO-EOSA-1

The Gammaproteobacterial Sulfur Oxidizer-EOSA-1 (GSO-EOSA-1) cluster, represented in RefSeq by the OTUs “*Candidatus Vesicomyosocius okutanii*” strain HA and “*Candidatus Ruthia magnifica*” strain Cm. (*Calyptogena magnifica*) (Walsh *et al.*, 2009), made a large contribution to variance between the NZ and SZ, with higher abundance in the SZ: relative abundances of GSO-EOSA-1 in the SZ were 5.2%, 3.4% and 0.25% in the 0.1, 0.8 and 3.0 μm size fractions respectively, compared to 1.1%, 0.84% and 0.30% in the NZ (Table 3.3). The contribution to variance of this group was highest in the 0.1 μm size fraction, followed by 0.8 μm and 3.0 μm (Table 3.4). This pattern most likely represents a small cell size and lack of association with particulate matter.

“*Ca. R. magnifica*” and “*Ca. V. okutanii*” are chemoautotrophic endosymbionts of deep-sea bivalves (Kuwahara *et al.*, 2007; Newton *et al.*, 2007) and are thus unlikely to be present in open ocean surface waters. However, GSO-EOSA-1 representative ARCTIC96BD-19 has recently been reported at high abundance in Antarctic coastal waters (Ghilione and Murray, 2011; Grzymski *et al.*, 2012). The majority of 16S rRNA genes from this metagenome with best BLASTN hits to “*Ca. R. magnifica*” and “*Ca. V. okutanii*” clustered with ARTIC96BD-19 in a neighbour-joining phylogenetic tree (Fig. 3.4), indicating this is the dominant GSO-EOSA-1 representative. Single-cell genomic analysis of ARCTIC96BD-19

Table 3.5: The thirty KEGG modules with the highest contributions to the difference between the NZ and SZ. Abundances are zonal averages and have been standardised and log-transformed.

KEGG module	Abundance South	Abundance North	Contribution to variance (%)
Photosystem II	0.42	0.57	2.21
Complex I (NADH dehydrogenase), NADH dehydrogenase I/diaphorase subunit of the bidirectional hydrogenase	0.01	0.24	1.80
Photosystem I	0.43	0.34	1.70
Pyrimidine deoxyribonucleotide biosynthesis, CDP/CTP → dCDP/dCTP,dTDP/dTTP	0.51	0.66	1.16
Histidine degradation, histidine → N-formiminoglutamate → glutamate	0.42	0.31	1.14
Methionine salvage pathway	0.29	0.43	1.14
sn-Glycerol 3-phosphate transport system	0.29	0.16	1.11
Complex I (NADH dehydrogenase), NADH dehydrogenase I	1.08	1.05	1.06
Branched-chain amino acid transport system	0.79	0.83	0.96
Dipeptide transport system	0.14	0.02	0.95
Adenine nucleotide biosynthesis, IMP → ADP/dADP,ATP/dATP	0.62	0.74	0.95
Glycine betaine/proline transport system	0.66	0.56	0.94
Sulfur reduction, sulfate → H ₂ S	0.54	0.44	0.91
Simple sugar transport system	0.46	0.39	0.90
Peptides/nickel transport system	0.99	0.98	0.89
Ribosome, eukaryotes	0.26	0.27	0.89
Multiple sugar transport system	0.55	0.55	0.86
Type II general secretion system	0.21	0.21	0.82
Sulfonate/nitrate/taurine transport system	0.45	0.37	0.82
Guanine nucleotide biosynthesis, IMP → GDP/dGDP,GTP/dGTP	0.72	0.82	0.81
RNA polymerase II, eukaryotes	0.11	0.20	0.76
Histidine biosynthesis, PRPP → histidine	0.94	0.86	0.76
Putrescine transport system	0.18	0.09	0.72
Leucine biosynthesis, pyruvate → 2-oxoisovalerate → leucine	1.29	1.37	0.71
C5 isoprenoid biosynthesis, non-mevalonate pathway	0.70	0.77	0.71
Leucine degradation, leucine → acetoacetate + acetyl-CoA	0.64	0.59	0.71
Thiamine transport system	0.13	0.05	0.69
Spliceosome, 35S U5-snRNP	0.18	0.20	0.68
Cytochrome b6f complex	0.14	0.12	0.67
Menaquinone biosynthesis, chorismate → menaquinone	0.25	0.27	0.66

Table 3.6: The thirty KEGG ortholog groups with the highest contribution to the difference between the NZ and SZ. Abundances are zonal averages and have been standardised and log-transformed. As each ortholog group on each size fraction was encoded as a separate variable in the SIMPER analysis, the size fraction is given after each ortholog group name.

KEGG ortholog group	Abundance South	Abundance North	Contribution to variance (%)
Hypothetical protein 3.0 μm	0.11	0.24	0.26
Hypothetical protein 0.8 μm	0.68	0.57	0.24
Ribonucleoside-diphosphate reductase alpha chain [EC:1.17.4.1] 0.8 μm	0.17	0.24	0.15
DNA polymerase III subunit alpha [EC:2.7.7.7] 0.8 μm	0.25	0.19	0.14
Hypothetical protein 0.1 μm	0.26	0.24	0.12
Proline dehydrogenase / delta 1-pyrroline-5-carboxylate 0.8 μm	0.10	0.04	0.12
Aminomethyltransferase [EC:2.1.2.10] 0.8 μm	0.25	0.19	0.12
Ribonucleoside-diphosphate reductase alpha chain [EC:1.17.4.1] 3.0 μm	0.02	0.08	0.12
Sarcosine oxidase, subunit alpha [EC:1.5.3.1] 0.8 μm	0.22	0.17	0.12
Integrator complex subunit 6 3.0 μm	0.07	0.05	0.11
Multicomponent $\text{Na}^+:\text{H}^+$ antiporter subunit D 0.8 μm	0.11	0.05	0.11
Glutamine synthetase [EC:6.3.1.2] 0.8 μm	0.24	0.19	0.11
Pyruvate dehydrogenase E1 component [EC:1.2.4.1] 0.8 μm	0.15	0.10	0.11
Cobaltochelatase CobN [EC:6.6.1.2] 0.8 μm	0.11	0.06	0.11
Formate dehydrogenase, alpha subunit [EC:1.2.1.2] 0.8 μm	0.15	0.10	0.11
DNA-directed RNA polymerase subunit beta [EC:2.7.7.6] 3.0 μm	0.03	0.08	0.11
Glutamate synthase (NADPH/NADH) large chain [EC:1.4.1.13 1.4.1.14] 0.8 μm	0.25	0.22	0.11
Dimethylglycine dehydrogenase [EC:1.5.99.2] 0.8 μm	0.17	0.14	0.11
Flagellin 0.8 μm	0.06	0.10	0.10
DNA-directed RNA polymerase subunit beta [EC:2.7.7.6] 3.0 μm^a	0.03	0.08	0.10
Photosystem II PsbA protein 0.8 μm	0.01	0.06	0.09
Aldehyde dehydrogenase (NAD+) [EC:1.2.1.3] 0.8 μm	0.17	0.13	0.09
Glutamate synthase (NADPH/NADH) large chain [EC:1.4.1.13 1.4.1.14] 3.0 μm	0.02	0.07	0.09
Thymidylate synthase (FAD) [EC:2.1.1.148] 0.8 μm	0.02	0.06	0.09
Topoisomerase IV subunit A [EC:5.99.1.-] 0.8 μm	0.11	0.07	0.09
DNA mismatch repair protein MutS 0.8 μm	0.13	0.08	0.09
Glutamate dehydrogenase [EC:1.4.1.2] 0.8 μm	0.07	0.03	0.09
DNA polymerase I [EC:2.7.7.7] 0.1 μm	0.12	0.11	0.09
GTP-binding protein 0.8 μm	0.26	0.21	0.09
GTP-binding protein 3.0 μm	0.03	0.07	0.09



Figure 3.4: Neighbour-joining tree of GSO-EOSA-1-like 16S rRNA gene sequences from the metagenomes in this study. Sequences labeled in black text are reads from the metagenomes. Red labels are 16S rRNA gene sequences from Gammaproteobacterial Sulfur Oxidizers (GSO) and other Gammaproteobacteria. The tree was constructed using ARB (Ludwig et al., 2004).






Figure 3.5: Examples of read recruitment (fragment recruitment) plots of metagenomic reads to reference genomes. The even distribution of recruitment sites across the genome suggests that OTUs identifications are relatively accurate; an OTU identification resulting from a small region of genomic identity would generate a plot with read recruitment concentrated in small islands.

from global mesopelagic waters indicates the lineage is probably mixotrophic, able to couple carbon fixation to oxidation of reduced sulphur compounds as well as assimilate organic carbon (Swan *et al.*, 2011). GSO-EOSA-1 cytochrome C oxidase (CoxII) has been identified in a winter metaproteome of Antarctic Peninsula coastal waters, suggesting the capacity for aerobic respiration (Williams *et al.*, 2012a). Taken together, this evidence suggests the GSO-EOSA-1 representative in Antarctic coastal waters is a versatile chemolithoautotroph capable of aerobic respiration.

It has been proposed that during the winter months, chemolithoautotrophy is dominant over photoautotrophy as the major carbon fixation process in AZ waters due to the lack of available light, both from seasonal darkness and ice shading (Grzymski *et al.*, 2012). The high relative abundance of GSO-EOSA-1 in SZ compared to NZ waters may therefore represent the remnants of an annual winter increase in population in the marginal ice zone which does not occur in the open ocean.

Ammonia-oxidizing Crenarchaeota

Nitrosopumilus maritimus SCM1 and *Cenarchaeum symbiosum* are chemolithoautotrophic, nitrifying members of the Marine Group I Crenarchaeota (MGI) (Preston *et al.*, 1996; Walker *et al.*, 2010) and are the only representatives in the reference database of the Ammonia-Oxidizing Archaea (AOA). The contribution of OTUs of *C. symbiosum* to the AOA signature was low. As *C. symbiosum* is a sponge symbiont (Preston *et al.*, 1996) and given the poor representation of AOA in RefSeq, it is likely this OTU has attracted sequences originating from planktonic AOA and *C. symbiosum* itself is not present. AOA were moderate contributors to variance between the NZ and SZ, and were overrepresented in the SZ in all size fractions (Fig. 3.3). As with the GSO-EOSA-1 cluster, MGI have been proposed to be abundant chemolithoautotrophs and therefore major drivers of winter carbon fixation in Antarctic coastal waters (Grzymski *et al.*, 2012; Williams *et al.*, 2012a).

Sample 353 had a particularly high relative abundance of *N. maritimus* OTUs (7.5% of the 0.1 µm fraction; 0.8 µm: 11%; 3.0 µm: 12%). This sample was taken closer to the Antarctic continent (3.7 km) than any other, in relatively shallow (180 m) waters 17.6 km from the Mertz Glacier. The high abundance of ammonia oxidizers may reflect an input of ammonia from terrestrial sources (e.g. penguin guano), or resuspension of benthic sediments in which MGI are abundant (Bowman and McCuaig, 2003) by near-shore turbulence and iceberg scouring. Breakdown of water column stratification has been previously suggested as a cause of increased AOA abundance in Antarctic coastal surface waters (Kalanetra *et al.*, 2009).

Cyanobacteria

OTUs of the cyanobacterial genera *Prochlorococcus* and *Synechococcus* were overrepresented in the NZ in all size fractions (Fig. 3.3). The mean relative abundance of cyanobacteria in samples 367 and 368, the two northernmost samples, was strikingly higher than the mean abundance across all other samples in the NZ. *Synechococcus* sp. CC9902 alone composed greater than 22% of the 0.8 µm fraction in these samples, consistent with *Synechococcus* species' average cell diameter of approximately 0.9 µm. The high abundance of both cyanobacterial genera on the 3.0 µm fraction has previously been reported (Lauro *et al.*, 2011) and may be attributable to aggregation (Lomas and Moran, 2011).

Samples 367 and 368 were separated from the other samples north of the PF by the STF. While the STF was not a significant boundary on the assemblage level, it may mark a significant biogeographic boundary for these cyanobacteria. *Synechococcus* and *Prochlorococcus* together represent a large proportion of both phytoplankton abundance and carbon fixation in temperate and tropical waters, in many regions contributing more than half of total primary production (Liu *et al.*, 1997, 1998; André *et al.*, 1999). The role of the STF in determining the latitudinal range of *Synechococcus* and *Prochlorococcus* is therefore important, as it will affect models of ocean productivity under changing climactic conditions, and warrants further investigation. Despite the high abundance of cyanobacteria north of the STF, they were also a significant feature of the SAZ; for example, *Synechococcus* sp. CC9902 composed 3–5% of the 0.8 µm fraction in SAZ samples.

These results extend the latitudinal distribution of both *Prochlorococcus* and *Synechococcus* to include presence at very low abundance as far south as the Antarctic coast. *Prochlorococcus* have been reported to be restricted to tropical and subtropical waters within 40° of latitude (Partensky *et al.*, 1999), and to be a negligible (Ghiglione and Murray, 2011) or undetectable (Grzymski *et al.*, 2012) component of marine microorganisms in Antarctic waters. However, these findings are consistent with findings of a logarithmic relationship of cyanobacterial numbers with temperature, where cyanobacteria were found at concentrations of 103–104 cells per litre even in the coldest waters, approximately four orders of magnitude less than in waters around Tasmania (Marchant *et al.*, 1987). Cyanophage proteins have also been detected in a metaproteomic analysis of Antarctic Peninsula coastal surface waters (Williams *et al.*, 2012a).

SAR11 and SAR116 clades

“*Candidatus Pelagibacter ubique*” HTCC1062 is a good representative of total SAR11 abundance in this study, as it is a member of the SAR11 phylotype which is most abundant in SO waters (Brown *et al.*, 2012). “*Ca. P. ubique*” HTCC1062 was the most abundant OTU across all samples and fractions (NZ average: 62%, 25% and 24% of the 0.1 µm, 0.8 µm and 3.0 µm fractions respectively; SZ: 59%, 22% and 18%) and one of the most significant contributors to variance between the NZ and SZ. The high abundance of SAR11 in the 0.1 µm fraction is consistent with the small size of SAR11 cells (Rappé *et al.*, 2002). The higher representation in the NZ may reflect the competitiveness of SAR11 members in regions with low Dissolved Organic Carbon (DOC) concentrations due to low primary productivity (Giovannoni *et al.*, 2005; Alonso and Perenthaler, 2006), such as the High Nutrient, Low Chlorophyll (HNLC) SAZ. Overall, these findings are consistent with reports that SAR11 is ubiquitous in the world’s oceans (Mary *et al.*, 2006; Carlson *et al.*, 2009) and more abundant north of the ACC (Giebel *et al.*, 2009).

OTUs of “*Candidatus Puniceispirillum marinum*” from the SAR116 clade were a moderate contributor to variance between the NZ and SZ with higher abundance in the NZ (Fig. 3.3). A genomic analysis reported “*Ca. P. marinum*” IMCC1322 to be a metabolic generalist with genes for aerobic CO fixation, C1 metabolism and a “*Ca. P. ubique*”-like dimethylsulfoniopropionate (DMSP) demethylase, suggesting SAR116 and SAR11 occupy similar ecological niches (Oh *et al.*, 2010). In the Scotia Sea, SAR116 abundance (determined using Fluorescence *In Situ* Hybridization (FISH)) was reported to be higher in more productive waters where SAR11 numbers were lower (Topping *et al.*, 2006). However, this analysis across an extended latitudinal transect indicates that overall SAR11 and SAR116 have similar biogeographic distributions.

Bacteroidetes

OTUs of the phylum Bacteroidetes, in particular members of the class Flavobacteria, were found to be abundant (NZ average: 1.2%, 5.0% and 6.9% of the 0.1 µm, 0.8 µm and 3.0 µm fractions respectively; SZ: 2.3%, 9.8% and 9.1%) and significant contributors to variance between the NZ and SZ (Fig. 3.3). Flavobacteria have been previously reported to compose the majority of both Bacteroidetes (Murray and Grzymski, 2007) and total planktonic biomass (Abell and Bowman, 2005b) in the SO, as well as being abundant in sea ice (Brown and Bowman, 2001). As heterotrophic degraders of High Molecular Weight (HMW) compounds in the form of both Dissolved Organic Matter (DOM) and Particulate Organic Matter (POM) (Kirchman, 2002), marine Flavobacteria are major components of marine aggregates (Rath *et al.*, 1998; Crump *et al.*, 1999; Zhang *et al.*, 2007). The higher abundance of Flavobacteria OTUs on the 0.8 µm and 3.0 µm fractions indicates their association with particulate matter. Similar size partitioning of SO Flavobacteria has previously been reported (Abell and Bowman, 2005b).

The higher abundance of OTUs of Flavobacteria in the SZ may reflect an input of cells from melting sea ice (Brown and Bowman, 2001), the higher rates of primary productivity in the south, and the role of the Flavobacteria as degraders of HMW DOM. Because deposition of marine snow is a major route for sequestration of fixed carbon in the ocean (e.g. Hessen *et al.*, 2004), the Flavobacteria that associate with this particulate matter represent a remineralising shunt, which would decrease carbon sequestration by this route.

Rhodobacterales

Members of the order Rhodobacterales were abundant (NZ average: 1.2%, 10% and 5.5% of the 0.1 µm, 0.8 µm and 3.0 µm fractions respectively; SZ: 1.6%, 13% and 7.9%) and high contributors to variance, overrepresented in the SZ on all size fractions. As several members of the Roseobacter clade have been shown to have symbiotic relationships with marine eukaryotic algae (Buchan *et al.*, 2005; Wagner-Döbler and Biebl, 2006), and their abundance in the SO has previously been linked to phytoplankton blooms (West *et al.*, 2008; Obernosterer *et al.*, 2011), it is likely that their overrepresentation in the SZ is related to the higher density of phytoplankton in the AZ.

OTUs of *Roseobacter denitrificans* Och114 and *Silicibacter pomeroyi* DSS-3 were consistently the most abundant Roseobacter clade representatives. *R. denitrificans* and *S. pomeroyi* fall within a subclade of Aerobic Anoxygenic Phototrophic (AAP) members of the Roseobacter clade (Swingley *et al.*, 2007). These species have diverse mixotrophic metabolisms, with genomic and experimental evidence of photoheterotrophic respiration of organic carbon, fixation of CO₂, oxidation of CO, oxidation of reduced sulfur compounds, and utilization of the abundant marine osmolyte DMSP (King, 2003; Moran *et al.*, 2004; Wagner-Döbler and Biebl, 2006; Swingley *et al.*, 2007; Brinkhoff *et al.*, 2008; Howard *et al.*, 2008). This metabolic diversity suggests a complex ecological role, particularly with respect to the capture and release of climatically active gases (CO₂, CO, dimethylsulfide) involved in carbon and sulfur cycling.

Alteromonadales

Members of the gammaproteobacterial order Alteromonadales were large contributors to variance. Most OTUs were overrepresented in the SZ but some were overrepresented in the NZ on the 3.0 µm fraction (Fig. 3.3). *Colwellia psychrerythraea* 34H was one of the most abundant OTUs in the Alteromonadales that exhibited this distribution (NZ average: 0.14%, 2.2% and 16% of the 0.1 µm, 0.8 µm and 3.0 µm fractions respectively; SZ: 0.52%, 5.1% and 10%). *C. psychrerythraea* 34H was isolated from Arctic sediment, grows well at low temperatures and secretes extracellular polysaccharides (Huston *et al.*, 2000; Junge *et al.*, 2003; Methé *et al.*, 2005). Similar to other *Colwellia* species grown under laboratory conditions, cells have widths of 0.4–0.8 µm and lengths of 1.5–4.5 µm (Jung *et al.*, 2006). Growth temperature can have a major impact on cell morphology, enzyme secretion and global gene expression in psychrophiles (e.g. Feller and Gerday, 2003; Junge *et al.*, 2003; Williams *et al.*, 2011; Cavicchioli, 2006; Campanaro *et al.*, 2011). Moreover, marine bacteria can alter their cell dimensions in response to nutrient flux (e.g. Kjelleberg *et al.*, 1987). It is therefore possible that the populations of Alteromonadales captured on the 3.0 µm filters (overrepresented in the NZ) had different physiological properties to those on the 0.1 and 0.8 µm filters (overrepresented in the SZ).

Verrucomicrobia

Two representatives of the phylum Verrucomicrobia, *Coraliomargarita akajimensis* and *Akkermansia* sp. Muc-30, were moderate contributors to variance and overrepresented in the SZ (Fig. 3.3). Surprisingly given the small cell size of *C. akajimensis* (Yoon *et al.*, 2007), its contribution to variance increased with size fraction. A global survey reported a similar fractionation pattern, and suggested marine Verrucomicrobia may be predominantly particle attached (Freitas *et al.*, 2012). However, little else is known about the distribution and ecological roles of marine Verrucomicrobia (Freitas *et al.*, 2012).

Functional capacities differentiating the zones

A number of modules with transport functions (sn-glycerol 3-phosphate transport system, dipeptide transport system, peptides/nickel transport system, simple sugar transport system, sulfonate/nitrate/taurine transport system) were overrepresented in the SZ (Table 3.5). As the genomes of copiotrophic bacteria have evolved to have a higher number of narrow-specificity transporters relative to oligotrophic genomes (Lauro *et al.*, 2009), these differences may reflect the higher nutrient availability and thus a dominance of copiotrophs in the SZ. The taxonomic contributors to these modules were varied, although members of the Rhodobacterales were prominent (Fig. 3.6).

The glycine betaine/proline transport module was also overrepresented in the SZ, though this probably reflects glycine betaine's role as an osmo- and cryoprotectant in the colder SZ waters. This




Figure 3.6: Decomposition of KEGG modules of interest to contributing classes, orders or genera. The left side of each stack (S) indicates the proportion of the module abundance contributed by each class, order or genus in the South Zone, while the right side (N) represents the North Zone. As the contributions are relative and represent unitless module abundances, no axis is given and proportions are not comparable between modules. Contributing classes, orders or genera are arranged in descending order of the difference in the relative contributions between the zones. Only the eight highest contributors for each module are shown, with the remainder collapsed into the “Other” group. The taxonomic ranks to which each module was decomposed are as follows: sn-glycerol 3-phosphate transport, peptide-nickel transport, simple sugar transport and sulfonate/nitrate/taurine transport were decomposed to order; glycine betaine/proline transport and branched-chain amino acid transport to genus; pyrimidine deoxyribonucleotide biosynthesis, adenine nucleotide biosynthesis and guanine nucleotide biosynthesis to order; methionine salvage to genus; sulphur reduction to class; photosystem I and photosystem II to genus; histidine degradation to glutamate and histidine biosynthesis to class.

is supported by the major taxonomic contributor to this module, genus *Psychromonas*, which includes several psychrophilic species.

One exception to this pattern was the branched-chain amino acid transport system module, overrepresented in the NZ. The genera *Pelagibacter* and *Puniceispirillum* were major contributors to this module’s overabundance in the NZ (Fig. 3.6). As both SAR11 (Giovannoni *et al.*, 2005) and SAR116 (Grote *et al.*, 2011) representatives encode branched-chain amino acid transporters, the abundance of this module is likely to represent taxonomic differences between the zones.

Biosynthesis pathways for all major nucleic acids (pyrimidine deoxyribonucleotide biosynthesis, adenine nucleotide biosynthesis, guanine nucleotide biosynthesis) were consistently high contributors to variance and overabundant in the NZ. This pattern seems inconsistent with the more oligotrophic nature of the NZ, as oligotrophic cells generally have smaller genomes (Lauro *et al.*, 2009) and slower growth rates than copiotrophs, and would therefore be expected to require a lower rate of de novo nucleotide biosynthesis. A possible explanation for this is that SZ cells have higher availability of extracellular DNA as a byproduct of decaying phytoplankton (Lomas and Moran, 2011), which can be imported and salvaged for nucleic acids (Paul *et al.*, 1988) thus reducing the requirement for de novo synthesis. No single taxonomic group contributed a large fraction of the difference in this module (Fig. 3.6), suggesting this is a widespread adaptation.

The methionine salvage pathway module had a large contribution to variance between the zones and was overrepresented north of the PF. This may reflect the higher availability of DMSP in the SZ as a byproduct of blooming eukaryotic algae. DMSP is a major carbon and sulfur source for marine microorganisms, and is commonly assimilated by bacteria through demethylation to methylmercaptopyruvate (MMPA), followed by further catabolism to the climatically important compounds dimethylsulfide or methanethiol (review in Curson *et al.*, 2011). However, when DMSP is scarce, MMPA may be derived from methionine through the alternative methionine salvage pathway (Reisch *et al.*, 2011). The genus *Synechococcus*, a noted contributor to marine DMSP uptake and assimilation (Vila-Costa *et al.*, 2006), was a very high contributor to the abundance of this module in the NZ (Fig. 3.6), suggesting *Synechococcus* species may use this route when DMSP is unavailable.

The sulfur reduction module was overrepresented in the SZ, and it is likely that this result is strongly driven by taxonomic differences. While the taxonomic breakdown indicated a large number of genera contributed to the difference, the Gammaproteobacteria were the highest-contributing class (Fig. 3.6). This module also includes the assimilatory sulfate reduction pathway, which is widespread in marine bacteria, but is absent from SAR11, with known representatives reported to lack genes for assimilatory sulfate reduction (cysDNCHI) (Tripp *et al.*, 2008). The higher relative abundance of SAR11 in the NZ may therefore contribute to the lower abundance of genes for assimilatory sulfate reduction in that zone.

The sulfur reduction module also included adenylylsulfate reductase (APS reductase, encoded by aprAB), an enzyme implicated in sulfite detoxification during heterotrophic growth on organosulfonates (Meyer and Kuever, 2007) (N.B. in recent KEGG releases, aprA is no longer included in this module). As the GSO-EOSA-1 representative SUP05 has been found to encode APS reductase, the overabundance of this module may reflect sulfur oxidation through the reverse dissimilatory sulfate reduction pathway (Walsh *et al.*, 2009). Also, Roseobacter clade bacteria are involved in the decomposition of abundant organic sulfur compounds (e.g. DMSP, organosulfonates), and hence have been

accorded an important role in marine sulfur cycling (Moran *et al.*, 2007).

The photosystem II module was overrepresented in the NZ. Given the underrepresentation of cyanobacterial OTUs in the SZ, this may reflect a dominance of primary production by eukaryotic algae south of the PF and cyanobacteria to the north. Decomposition of the taxonomic affiliations of ortholog groups contributing to this module found OTUs of *Synechococcus* and *Prochlorococcus* to be major contributors to the difference (Fig. 3.6). Variation in the photosystem I module, which was marginally overrepresented in the SZ, could largely be attributed to diatoms and other eukaryotic phytoplankton (Fig. 3.6), again supporting a dominance of eukaryotic phytoplankton in SZ primary production. Diatoms have previously been reported at higher abundance south of the PF, and their distribution is likely to be linked to the higher concentration of dissolved silica in that region (Trull *et al.*, 2001). As both eukaryotic phytoplankton and cyanobacteria would be expected to encode both complete photosystems, the differences in module abundance probably reflect the degree of similarity between the photosystem I and II genes in the KEGG database and those found in the sampled environments.

The histidine degradation to glutamate module, which comprises four ortholog groups mediating the degradation of histidine to glutamate via N-formiminoglutamate, was overrepresented in the SZ. The histidine biosynthesis module was also overrepresented in the SZ. While the concentration of dissolved histidine in the SO is generally low (Kawahata and Ishizuka, 2000), blooming eukaryotic phytoplankton (which are more prevalent in the SZ) may deplete nitrate while releasing Dissolved Free Amino Acids (DFAA). As DFAA become available, they are used by bacteria to sense the decaying bloom. Histidine may therefore act as a proxy for DFAA to regulate the expression of bacterial aminopeptidases, which are involved in lysing diatoms (Bidle and Azam, 2001). The class Bacteroidetes, while a small contributor to the histidine biosynthesis module in the SZ, was a large contributor to histidine degradation (Fig. 3.6), supporting an association between Bacteroidetes and phytoplanktonic bloom products. It is also possible that uptake and degradation of histidine to glutamate (which generates ammonia as a by-product) may function as a limited nitrogen source.

Conclusions: Biogeographic role of the Polar Front

These results show that there are major taxonomic and functional differences across the PF. The differences in functional potential between the NZ and SZ reflect both their taxonomic profiles and fundamental trophic and ecological differences. In particular, they provide genomic support that the NZ is more oligotrophic than the SZ (Pollard *et al.*, 2002; Giovannoni *et al.*, 2005; Alonso and Pernthaler, 2006; Lauro *et al.*, 2009), and are consistent with the observation that primary production is higher south of the PF (Strutton *et al.*, 2000; Williams *et al.*, 2010). Our findings extend previous work in defining the PF as a strong biogeographic boundary which shapes not only the composition, but also the functional capacity of microbial communities in the SO.

These results do not exclude the possibility that other major SO fronts, particularly the STF and SAF, are also significant biogeographic boundaries, as has been reported in some previous reports for specific taxonomic groups (e.g. Abell and Bowman, 2005b). While the sampling resolution in this study was not sufficient to resolve the effects of other fronts, there are some indications in the data of further structure within the zones. The two samples north of the STF had significantly larger cyanobacterial populations than the remaining NZ samples (see discussion of *Prochlorococcus* and *Synechococcus*, above). Future sampling across these fronts at higher resolution will provide the data necessary to investigate finer biogeographic patterns.

The nature and function of microbial communities in the SO are of global significance. Knowledge of these communities and their biogeographic drivers has relevance for understanding and predicting the long-term effects of climate change. These findings provide a basis for predicting how climate change-driven shifts in the SO may affect microbial communities; in particular, the effects of changes in the nature and location of the ACC on the ecosystem functions of SO microorganisms.

References

- Abell G. C. J. and Bowman J. P. (2005). Colonization and community dynamics of class *Flavobacteria* on diatom detritus in experimental mesocosms based on Southern Ocean seawater. *FEMS Microbiology Ecology*, 53(3):379–391.
- Abell G. G. J. and Bowman J. P. (2005). Ecological and biogeographic relationships of class Flavobacteria in the Southern Ocean. *FEMS Microbiology Ecology*, 51:265–277.
- Agogué H., Lamy D., Neal P. R., Sogin M. L., and Herndl G. J. (2011). Water mass-specificity of bacterial communities in the North Atlantic revealed by massively parallel sequencing. *Molecular Ecology*, 20(2):258–274.
- Alonso C. and Pernthaler J. (2006). Roseobacter and SAR11 dominate microbial glucose uptake in coastal North Sea waters. *Environmental Microbiology*, 8(11):2022–2030.
- Anderson L. A. and Sarmiento J. L. (1994). Redfield ratios of remineralization determined by nutrient data analysis. *Global Biogeochemical Cycles*, 8(1):65–80.
- André J. M., Navarette C., Blanchot J., and Radenac M. H. (1999). Picophytoplankton dynamics in the equatorial Pacific: Growth and grazing rates from cytometric counts. *Journal of Geophysical Research*, 104(C2):3369–3380.
- Angly F. E., Felts B., Breitbart M., Salamon P., Edwards R. A., Carlson C., Chan A. M., Haynes M., Kelley S., Liu H., Mahaffy J. M., Mueller J. E., Nulton J., Olson R., Parsons R., Rayhawk S., Suttle C. A., and Rohwer F. (2006). The marine viromes of four oceanic regions. *PLoS Biology*, 4(11):e368.
- Angly F. E., Willner D., Prieto-Davó A., Edwards R. A., Schmieder R., Vega-Thurber R., Antonopoulos D. A., Barott K., Cottrell M. T., Desnues C., Dinsdale E. A., Furlan M., Haynes M., Henn M. R., Hu Y., Kirchman D. L., McDole T., McPherson J. D., Meyer F., Miller R. M., Mundt E., Naviaux R. K., Rodriguez-Mueller B., Stevens R., Wegley L., Zhang L., Zhu B., and Rohwer F. (2009). The GAAS Metagenomic Tool and Its Estimations of Viral and Microbial Average Genome Size in Four Major Biomes. *PLoS Computational Biology*, 5(12):e1000593.
- Aoki S., Yoritaka M., and Masuyama A. (2003). Multidecadal warming of subsurface temperature in the Indian sector of the Southern Ocean. *Journal of Geophysical Research*, 108(C4):8081–8088.
- Baas Becking L. G. M. *Geobiologie Of Inleiding Tot De Milieukunde*. W.P. Van Stockum & Zoon, The Hague, 1934.
- Beja O., Aravind L., Koonin E. V., Suzuki M. T., Hadd A., Nguyen L. P., Jovanovich S. B., Gates C. M., Feldman R. A., Spudich J. L., Spudich E. N., and DeLong E. F. (2000). Bacterial rhodopsin: evidence for a new type of phototrophy in the sea. *Science*, 289(5486):1902–1906.
- Béjà O., Suzuki M. T., Heidelberg J. F., Nelson W. C., Preston C. M., Hamada T., Eisen J. A., Fraser C. M., and DeLong E. F. (2002). Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature*, 415(6872):630–633.
- Berg I. A., Kockelkorn D., Buckel W., and Fuchs G. (2007). A 3-Hydroxypropionate/4-Hydroxybutyrate Autotrophic Carbon Dioxide Assimilation Pathway in Archaea. *Science*, 318(5857):1782–1786.

- Bidle K. D. and Azam F. (2001). Bacterial control of silicon regeneration from diatom detritus: significance of bacterial ectohydrolases and species identity. *Limnology and Oceanography*, 46(7):1606–1623.
- Biebl H., Allgaier M., Tindall B. J., Kobližek M., Lünsdorf H., Pukall R., and Wagner-Döbler I. (2005). *Dinoroseobacter shibae* gen. nov., sp. nov., a new aerobic phototrophic bacterium isolated from dinoflagellates. *International Journal of Systematic and Evolutionary Microbiology*, 55(Pt 3):1089–1096.
- Bissett A., Richardson A. E., Baker G., Wakelin S., and Thrall P. H. (2010). Life history determines biogeographical patterns of soil bacterial communities over multiple spatial scales. *Molecular Ecology*, 19(19):4315–4327.
- Böning C. W., Dispert A., Visbeck M., Rintoul S. R., and Schwarzkopf F. U. (2008). The response of the Antarctic Circumpolar Current to recent climate change. *Nature Geoscience*, 1(12):864–869.
- Bowman J. P. and McCuaig R. D. (2003). Biodiversity, community structural shifts, and biogeography of prokaryotes within Antarctic continental shelf sediment. *Applied and Environmental Microbiology*, 69(5):2463–2483.
- Bowman J. P., Rea S. M., McCammon S. A., and McMeekin T. A. (2000). Diversity and community structure within anoxic sediment from marine salinity meromictic lakes and a coastal meromictic marine basin, Vestfold Hills, Eastern Antarctica. *Environmental Microbiology*, 2(2):227–237.
- Boyd P. W., Jickells T., Law C. S., Blain S., Boyle E. A., Buesseler K. O., Coale K. H., Cullen J. J., Baar H. J. W.de, Follows M., Harvey M., Lancelot C., Levasseur M., Owens N. P. J., Pollard R., Rivkin R. B., Sarmiento J., Schoemann V., Smetacek V., Takeda S., Tsuda A., Turner S., and Watson A. J. (2007). Mesoscale Iron Enrichment Experiments 1993–2005: Synthesis and Future Directions. *Science*, 315(5812):612–617.
- Brinkhoff T., Giebel H.-A., and Simon M. (2008). Diversity, ecology, and genomics of the Roseobacter clade: a short overview. *Archives of Microbiology*, 189(6):531–539.
- Brinkmeyer R., Knittel K., Jürgens J., Weyland H., Amann R., and Helmke E. (2003). Diversity and Structure of Bacterial Communities in Arctic versus Antarctic Pack Ice. *Applied and Environmental Microbiology*, 69(11):6610–6619.
- Brown M. V. and Bowman J. P. (2001). A molecular phylogenetic survey of sea-ice microbial communities (SIMCO). *FEMS Microbiology Ecology*, 35(3):267–275.
- Brown M. V., Lauro F. M., DeMaere M. Z., Muir L., Wilkins D., Thomas T., Riddle M. J., Fuhrman J. A., Andrews-Pfannkoch C., Hoffman J. M., McQuaid J. B., Allen A., Rintoul S. R., and Cavicchioli R. (2012). Global biogeography of SAR11 marine bacteria. *Molecular systems biology*, 8.
- Buchan A., González J. M., and Moran M. A. (2005). Overview of the marine Roseobacter lineage. *Applied and Environmental Microbiology*, 71(10):5665–5677.
- Callahan J. E. (1972). The structure and circulation of deep water in the Antarctic. *Deep Sea Research and Oceanographic Abstracts*, 19(8):563–575.
- Campanaro S., Williams T. J., Burg D. W., De Francisci D., Treu L., Lauro F. M., and Cavicchioli R. (2011). Temperature-dependent global gene expression in the Antarctic archaeon *Methanococcoides burtonii*. *Environmental Microbiology*, 13(8):2018–2038.
- Canfield D. E., Stewart F. J., Thamdrup B., De Brabandere L., Dalsgaard T., DeLong E. F., Revsbech N. P., and Ulloa O. (2010). A Cryptic Sulfur Cycle in Oxygen-Minimum-Zone Waters off the Chilean Coast. *Science*, 330(6009):1375–1378.
- Caporaso J. G., Kuczynski J., Stombaugh J., Bittinger K., Bushman F. D., Costello E. K., Fierer N., Pena A. G., Goodrich J. K., and Gordon J. I. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nature methods*, 7(5):335–336.
- Carlson C. A., Morris R., Parsons R., Treusch A. H., Giovannoni S. J., and Vergin K. (2009). Seasonal dynamics of SAR11 populations in the euphotic and mesopelagic zones of the northwestern Sargasso Sea. *The ISME Journal*, 3(3):283–295.

- Cavicchioli R. (2006). Cold-adapted archaea. *Nature Reviews Microbiology*, 4(5):331–343.
- Cerovečki I., Talley L. D., and Mazloff M. R. (2011). A comparison of Southern Ocean air-sea buoyancy flux from an ocean state estimate with five other products. *Journal of Climate*, 24:6283–6306.
- Chiba S., Ishimaru T., Hosie G. W., and Fukuchi M. (2001). Spatio-temporal variability of zooplankton community structure off east Antarctica (90 to 160°E). *Marine Ecology Progress Series*, 216:95–108.
- Cho J. C. and Giovannoni S. J. (2004). Cultivation and Growth Characteristics of a Diverse Group of Oligotrophic Marine Gammaproteobacteria. *Applied and Environmental Microbiology*, 70(1):432–440.
- Cho J.-C. and Tiedje J. M. (2000). Biogeography and degree of endemism of fluorescent Pseudomonas strains in soil. *Applied and Environmental Microbiology*, 66(12):5448–5456.
- Christaki U., Obernosterer I., Van Wambeke F., Veldhuis M., Garcia N., and Catala P. (2008). Microbial food web structure in a naturally iron-fertilized area in the Southern Ocean (Kerguelen Plateau). *Deep Sea Research Part II: Topical Studies in Oceanography*, 55(5-7):706–719.
- Church M. J., DeLong E. F., Ducklow H. W., Karner M. B., Preston C. M., and Karl D. M. (2003). Abundance and distribution of planktonic Archaea and Bacteria in the waters west of the Antarctic Peninsula. *Limnology and Oceanography*, 48(5):1893–1902.
- Clarke K. R. and Gorley R. N. *PRIMER v6: User Manual / Tutorial*, 1st edition edition, 2006.
- Clarke K. R. and Warwick R. M. *Change in marine communities: an approach to statistical analysis and interpretation*. PRIMER-E, Plymouth, 2nd edition, 2001.
- Coleman M. L. M. and Chisholm S. W. S. (2010). Ecosystem-specific selection pressures revealed through comparative population genomics. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 107(43):18634–18639.
- Cottrell M. T. and Kirchman D. L. (2000). Community Composition of Marine Bacterioplankton Determined by 16S rRNA Gene Clone Libraries and Fluorescence In Situ Hybridization. *Applied and Environmental Microbiology*, 66(12):5116–5122.
- Cottrell M. T., Waidner L. A., Yu L., and Kirchman D. L. (2005). Bacterial diversity of metagenomic and PCR libraries from the Delaware River. *Environmental Microbiology*, 7(12):1883–1895.
- Cox P. M., Betts R. A., Jones C. D., Spall S. A., and Totterdell I. J. (2000). Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature*, 408(6809):184–187.
- Crump B. C., Armbrust E. V., and Baross J. A. (1999). Phylogenetic analysis of particle-attached and free-living bacterial communities in the Columbia River, its estuary, and the adjacent coastal ocean. *Applied and Environmental Microbiology*, 65(7):3192–3204.
- Curson A. R. J., Todd J. D., Sullivan M. J., and Johnston A. W. B. (2011). Catabolism of dimethylsulphoniopropionate: microorganisms, enzymes and genes. *Nature Reviews Microbiology*, 9(12):849–859.
- Wit R.de and Bouvier T. (2006). ‘*Everything is everywhere, but, the environment selects*'; what did Baas Becking and Beijerinck really say? *Environmental Microbiology*, 8(4):755–758.
- Deacon G. E. R. (1982). Physical and biological zonation in the Southern Ocean. *Deep Sea Research Part A. Oceanographic Research Papers*, 29(1):1–15.
- Declerck S. A. J., Winter C., Shurin J. B., Suttle C. A., and Matthews B. (2013). Effects of patch connectivity and heterogeneity on metacommunity structure of planktonic bacteria and viruses. *The ISME Journal*, 7(3):533–542.
- DeLong E. F., Franks D. G., and Alldredge A. L. (1993). Phylogenetic Diversity of Aggregate-Attached vs. Free-Living Marine Bacterial Assemblages. *Limnology and Oceanography*, 38(5):924–934.
- DeLong E. F., Wu K. Y., Prézelin B. B., and Jovine R. V. (1994). High abundance of Archaea in Antarctic marine picoplankton. *Nature*, 371(6499):695–697.

- Dinsdale E. A., Edwards R. A., Hall D., Angly F., Breitbart M., Brulc J. M., Furlan M., Desnues C., Haynes M., Li L., McDaniel L., Moran M. A., Nelson K. E., Nilsson C., Olson R., Paul J., Brito B. R., Ruan Y., Swan B. K., Stevens R., Valentine D. L., Thurber R. V., Wegley L., White B. A., and Rohwer F. (2008). Functional metagenomic profiling of nine biomes. *Nature*, 452(7187):629–632.
- Dixon J. L., Beale R., and Nightingale P. D. (2011). Rapid biological oxidation of methanol in the tropical Atlantic: significance as a microbial carbon source. *Biogeosciences Discussions*, 8(2):3899–3921.
- Ducklow H. W., Myers K., Erickson M., Ghiglione J. F., and Murray A. E. (2011). Response of a summertime Antarctic marine bacterial community to glucose and ammonium enrichment. *Aquatic Microbial Ecology*, 64(3):205–220.
- Dupont C. L., Rusch D. B., Yooseph S., Lombardo M.-J., Richter R. A., Valas R., Novotny M., Yee-Greenbaum J., Selengut J. D., Haft D. H., Halpern A. L., Lasken R. S., Nealson K., Friedman R., and Venter J. C. (2011). Genomic insights to SAR86, an abundant and uncultivated marine bacterial lineage. pages 1–14.
- Eilers H., Pernthaler J., Glöckner F. O., and Amann R. (2000). Culturability and In Situ Abundance of Pelagic Bacteria from the North Sea. *Applied and Environmental Microbiology*, 66(7):3044–3051.
- El-Sayed S. Z. (2005). History and evolution of primary productivity studies of the Southern Ocean. *Polar Biology*, 28(6):423–438.
- Esper O. and Zonneveld K. A. F. (2002). Distribution of organic-walled dinoflagellate cysts in surface sediments of the Southern Ocean (eastern Atlantic sector) between the Subtropical Front and the Weddell Gyre. *Marine Micropaleontology*, 46(1):177–208.
- Evans C., Pearce I., and Brussaard C. P. D. (2009). Viral-mediated lysis of microbes and carbon release in the sub-Antarctic and Polar Frontal zones of the Australian Southern Ocean. *Environmental Microbiology*, 11(11):2924–2934.
- Evans C., Thomson P. G., Davidson A. T., Bowie A. R., Enden R. van den, Witte H., and Brussaard C. P. D. (2011). Potential climate change impacts on microbial distribution and carbon cycling in the Australian Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 58(21-22):2150–2161.
- Fandino L. B., Riemann L., Steward G. F., Long R. A., and Azam F. (2001). Variations in bacterial community structure during a dinoflagellate bloom analyzed by DGGE and 16S rDNA sequencing. *Aquatic Microbial Ecology*, 23:119.
- Feller G. and Gerday C. (2003). Psychrophilic enzymes: hot topics in cold adaptation. *Nature Reviews Microbiology*, 1(3):200–208.
- Firing Y. L., Chereskin T. K., and Mazloff M. R. (2011). Vertical structure and transport of the Antarctic Circumpolar Current in Drake Passage from direct velocity observations. *Journal of Geophysical Research*, 116(C8):C08015.
- Foldvik A. and Gammelsrød T. (1988). Notes on Southern Ocean hydrography, sea-ice and bottom water formation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 67(1-2):3–17.
- Freitas S., Hatosy S., Fuhrman J. A., Huse S. M., Welch D. B. M., Sogin M. L., and Martiny A. C. (2012). Global distribution and diversity of marine *Verrucomicrobia*. *The ISME Journal*, 6(8):1499–1505.
- Fuhrman J. A., Schwalbach M. S., and Stingl U. (2008). Proteorhodopsins: an array of physiological roles? *Nature Reviews Microbiology*, 6:488–494.
- Fyfe J. C. and Saenko O. A. (2005). Human-induced change in the Antarctic Circumpolar Current. *Journal of Climate*, 18(15):3068–3073.
- Galand P. E., Potvin M., Casamayor E. O., and Lovejoy C. (2009). Hydrography shapes bacterial biogeography of the deep Arctic Ocean. *Nature*, 44(4):564–576.

- García-Martínez J. and Rodríguez-Valera F. (2000). Microdiversity of uncultured marine prokaryotes: the SAR11 cluster and the marine Archaea of Group I. *Molecular Ecology*, 9(7):935–948.
- Gentile G., Giuliano L., D'Auria G., Smedile F., Azzaro M., De Domenico M., and Yakimov M. M. (2006). Study of bacterial communities in Antarctic coastal waters by a combination of 16S rRNA and 16S rDNA sequencing. *Environmental Microbiology*, 8(12):2150–2161.
- Ghiglione J. F. and Murray A. E. (2011). Pronounced summer to winter differences and higher winter time richness in coastal Antarctic marine bacterioplankton. *Environmental Microbiology*, 14(3):617–629.
- Ghiglione J.-F., Galand P. E., Pommier T., Pedrós-Alió C., Maas E. W., Bakker K., Bertilson S., Kirchman D. L., Lovejoy C., Yager P. L., and Murray A. E. (2012). Pole-to-pole biogeography of surface and deep marine bacterial communities. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 109(43):17633–17638.
- Giebel H.-A., Brinkhoff T., Zwisler W., Selje N., and Simon M. (2009). Distribution of *Roseobacter* RCA and SAR11 lineages and distinct bacterial communities from the subtropics to the Southern Ocean. *Environmental Microbiology*, 11(8):2164–2178.
- Giebel H.-A., Kalhoefer D., Lemke A., Thole S., Gahl-Janssen R., Simon M., and Brinkhoff T. (2010). Distribution of *Roseobacter* RCA and SAR11 lineages in the North Sea and characteristics of an abundant RCA isolate. *The ISME Journal*, 5:8–19.
- Gille S. T. (2002). Warming of the Southern Ocean Since the 1950s. *Science*, 295(5558):1275–1277.
- Giovannoni S. J., Tripp H. J., Givan S., Podar M., Vergin K. L., Baptista D., Bibbs L., Eads J., Richardson T. H., Noordewier M., Rappé M. S., Short J. M., Carrington J. C., and Mathur E. J. (2005). Genome streamlining in a cosmopolitan oceanic bacterium. *Science*, 309(5738):1242–1245.
- Giovannoni S. J., Hayakawa D. H., Tripp H. J., Stingl U., Givan S. A., Cho J.-C., Oh H.-M., Kitner J. B., Vergin K. L., and Rappé M. S. (2008). The small genome of an abundant coastal ocean methylotroph. *Environmental Microbiology*, 10(7):1771–1782.
- Glöckner F. O., Fuchs B. M., and Amann R. (1999). Bacterioplankton compositions of lakes and oceans: a first comparison based on fluorescence in situ hybridization. *Applied and Environmental Microbiology*, 65(8):3721–3726.
- González J. M., Fernández-Gómez B., Fernández-Guerra A., Gómez-Consarnau L., Sánchez O., Coll-Lladó M., Del Campo J., Escudero L., Rodríguez-Martínez R., Alonso-Sáez L., Latasa M., Paulsen I., Nedashkovskaya O., Lekunberri I., Pinhassi J., and Pedrós-Alió C. (2008). Genome analysis of the proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria). *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 105(25):8724–8729.
- Grossart H. P., Schlingloff A., Bernhard M., Simon M., and Brinkhoff T. (2004). Antagonistic activity of bacteria isolated from organic aggregates of the German Wadden Sea. *FEMS Microbiology Ecology*, 47(3):387–396.
- Grote J., Bayindirli C., Bergauer K., Moraes P.Carpintero de, Chen H., D'Ambrosio L., Edwards B., Fernández-Gómez B., Hamisi M., Logares R., Nguyen D., Rii Y. M., Saeck E., Schutte C., Widner B., Church M. J., Steward G. F., Karl D. M., DeLong E. F., Eppley J. M., Schuster S. C., Kyrpides N. C., and Rappé M. S. (2011). Draft genome sequence of strain HIMB100, a cultured representative of the SAR116 clade of marine *Alphaproteobacteria*. *Standards in Genomic Sciences*, 5(3):269–278.
- Grzymski J. J., Carter B. J., DeLong E. F., Feldman R. A., Ghadiri A., and Murray A. E. (2006). Comparative Genomics of DNA Fragments from Six Antarctic Marine Planktonic Bacteria. *Applied and Environmental Microbiology*, 72(2):1532–1541.
- Grzymski J. J., Riesenfeld C. S., Williams T. J., Dussaq A. M., Ducklow H., Erickson M., Cavicchioli R., and Murray A. E. (2012). A metagenomic assessment of winter and summer bacterioplankton from Antarctica Peninsula coastal surface waters. *The ISME Journal*, 6(10):1901–1915.

- Guixa-Boixereu N., Vaqué D., Gasol J. M., Sánchez-Cámara J., and Pedrós-Alio C. (2002). Viral distribution and activity in Antarctic waters. *Deep Sea Research Part II: Topical Studies in Oceanography*, 49 (4):827–845.
- Hamdan L. J., Coffin R. B., Sikaroodi M., Greinert J., Treude T., and Gillevet P. M. (2013). Ocean currents shape the microbiome of Arctic marine sediments. *The ISME Journal*, 7(4):685–696.
- Hamilton A. K., Lovejoy C., Galand P. E., and Ingram R. G. (2008). Water masses and biogeography of picoeukaryote assemblages in a cold hydrographically complex system. *Limnology and Oceanography*, pages 922–935.
- Hanson C. A., Fuhrman J. A., Horner-Devine M. C., and Martiny J. B. H. (2012). Beyond biogeographic patterns: processes shaping the microbial landscape. *Nature Reviews Microbiology*, 10(7):497–506.
- Head I. M., Hiorns W. D., Embley T. M., McCarthy A. J., and Saunders J. R. (1993). The phylogeny of autotrophic ammonia-oxidizing bacteria as determined by analysis of 16S ribosomal RNA gene sequences. *Journal of General Microbiology*, 139(6):1147–1153.
- Heikes B. G., Chang W., Pilson M. E. Q., Swift E., Singh H. B., Guenther A., Jacob D. J., Field B. D., Fall R., Riemer D., and Brand L. (2002). Atmospheric methanol budget and ocean implication. *Global Biogeochemical Cycles*, 16(4):1133.
- Hessen D. O., Ågren G. I., Anderson T. R., Elser J. J., and de Ruiter, P.C. (2004). Carbon sequestration in ecosystems: the role of stoichiometry. *Ecology*, 85(5):1179–1192.
- Hollibaugh J. T., Bano N., and Ducklow H. W. (2002). Widespread Distribution in Polar Oceans of a 16S rRNA Gene Sequence with Affinity to *Nitrosospira*-Like Ammonia-Oxidizing Bacteria. *Applied and Environmental Microbiology*, 68(3):1478–1484.
- Howard E. C., Sun S., Biers E. J., and Moran M. A. (2008). Abundant and diverse bacteria involved in DMSP degradation in marine surface waters. *Environmental Microbiology*, 10(9):2397–2410.
- Hunt B. P. V., Pakhomov E. A., and McQuaid C. D. (2001). Short-term variation and long-term changes in the oceanographic environment and zooplankton community in the vicinity of a sub-Antarctic archipelago. *Marine Biology*, 138:369–381.
- Huntley M. E., Lopez M. D., and Karl D. M. (1991). Top predators in the Southern ocean: a major leak in the biological carbon pump. *Science*, 253(5015):64–66.
- Huson D. H., Auch A. F., Qi J., and Schuster S. C. (2007). MEGAN analysis of metagenomic data. *Genome Research*, 17(3):377–386.
- Huston A. L., Krieger-Brockett B. B., and Deming J. W. (2000). Remarkably low temperature optima for extracellular enzyme activity from Arctic bacteria and sea ice. *Environmental Microbiology*, 2(4): 383–388.
- Ingalls A. E., Shah S. R., Hansman R. L., Aluwihare L. I., Santos G. M., Druffel E. R. M., and Pearson A. (2006). Quantifying archaeal community autotrophy in the mesopelagic ocean using natural radiocarbon. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 103 (17):6442–6447.
- Iverson V., Morris R. M., Frazar C. D., Berthiaume C. T., Morales R. L., and Armbrust E. V. (2012). Untangling Genomes from Metagenomes: Revealing an Uncultured Class of Marine Euryarchaeota. *Science*, 335(6068):587–590.
- Jacobs S. S. (2004). Bottom water production and its links with the thermohaline circulation. *Antarctic Science*, 16(04):427–437.
- Jamieson R. E., Rogers A. D., Billett D., Smale D. A., and Pearce D. A. (2012). Patterns of marine bacterioplankton biodiversity in the surface waters of the Scotia Arc, Southern Ocean. *FEMS Microbiology Ecology*, 80:452–468.

- Jung S.-Y., Oh T.-K., and Yoon J.-H. (2006). *Colwellia aestuarii* sp. nov., isolated from a tidal flat sediment in Korea. *International Journal of Systematic and Evolutionary Microbiology*, 56(1):33–37.
- Junge K., Eicken H., and Deming J. W. (2003). Motility of *Colwellia psychrerythraea* Strain 34H at Subzero Temperatures. *Applied and Environmental Microbiology*, 69(7):4282–4284.
- Kalanetra K. M., Bano N., and Hollibaugh J. T. (2009). Ammonia-oxidizing Archaea in the Arctic Ocean and Antarctic coastal waters. *Environmental Microbiology*, 11(9):2434–2445.
- Kawahata H. and Ishizuka T. (2000). Amino acids in interstitial waters from ODP Sites 689 and 690 on the Maud Rise, Antarctic Ocean. *Geochemical Journal*, 34(4):247–261.
- King G. M. (2003). Molecular and Culture-Based Analyses of Aerobic Carbon Monoxide Oxidizer Diversity. *Applied and Environmental Microbiology*, 69(12):7257–7265.
- Kirchman D. L. (2002). The ecology of *Cytophaga–Flavobacteria* in aquatic environments. *FEMS Microbiology Ecology*, 39(2):91–100.
- Kirchman D. L. *Microbial ecology of the oceans*. John Wiley & Sons, Inc., Hoboken, New Jersey, second edition, 2008.
- Kjelleberg S., Hermansson M., and Mårdén P. (1987). The transient phase between growth and non-growth of heterotrophic bacteria, with emphasis on the marine environment. *Annual Review of Microbiology*, 41:25–49.
- Knights D., Kuczynski J., Charlson E. S., Zaneveld J., Mozer M. C., Collman R. G., Bushman F. D., Knight R., and Kelley S. T. (2011). Bayesian community-wide culture-independent microbial source tracking. *Nature methods*, 8(9):761–763.
- Koh E. Y., Phua W., and Ryan K. G. (2011). Aerobic anoxygenic phototrophic bacteria in Antarctic sea ice and seawater. *Environmental Microbiology Reports*, 3(6):710–716.
- Kuwahara H., Yoshida T., Takaki Y., Shimamura S., Nishi S., Harada M., Matsuyama K., Takishita K., Kawato M., Uematsu K., Fujiwara Y., Sato T., Kato C., Kitagawa M., Kato I., and Maruyama T. (2007). Reduced Genome of the Thioautotrophic Intracellular Symbiont in a Deep-Sea Clam, *Calyptogena okutanii*. *Current Biology*, 17(10):881–886.
- Laubscher R. K., Perissinotto R., and McQuaid C. D. (1993). Phytoplankton production and biomass at frontal zones in the Atlantic sector of the Southern Ocean. *Polar Biology*, 13(7).
- Lauro F. M., Chastain R. A., Blankenship L. E., Yayanos A. A., and Bartlett D. H. (2007). The unique 16S rRNA genes of piezophiles reflect both phylogeny and adaptation. *Applied and Environmental Microbiology*, 73(3):838–845.
- Lauro F. M., McDougald D., Thomas T., Williams T. J., Egan S., Rice S., DeMaere M. Z., Ting L., Ertan H., Johnson J., Ferriera S., Lapidus A., Anderson I., Kyrpides N., Munk A. C., Detter C., Han C. S., Brown M. V., Robb F. T., Kjelleberg S., and Cavicchioli R. (2009). The genomic basis of trophic strategy in marine bacteria. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 106(37):15527–15533.
- Lauro F. M., DeMaere M. Z., Yau S., Brown M. V., Ng C., Wilkins D., Raftery M. J., Gibson J. A., Andrews-Pfannkoch C., Lewis M., Hoffman J. M., Thomas T., and Cavicchioli R. (2011). An integrative study of a meromictic lake ecosystem in Antarctica. *The ISME Journal*, 5(5):879–895.
- Legendre P. and Anderson M. J. (1999). Distance-based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecological Monographs*, 69(1):1–24.
- Liu H., Nolla H. A., and Campbell L. (1997). *Prochlorococcus* growth rate and contribution to primary production in the equatorial and subtropical North Pacific Ocean. *Aquatic Microbial Ecology*, 12(1):39–47.

- Liu H., Campbell L., Landry M. R., Nolla H. A., Brown S. L., and Constantinou J. (1998). *Prochlorococcus* and *Synechococcus* growth rates and contributions to production in the Arabian Sea during the 1995 Southwest and Northeast Monsoons. *Deep Sea Research Part II: Topical Studies in Oceanography*, 45 (10-11):2327–2352.
- Lo Giudice A., Caruso C., Mangano S., Bruni V., Domenico M., and Michaud L. (2011). Marine Bacterioplankton Diversity and Community Composition in an Antarctic Coastal Environment. *Microbial Ecology*, 63(1):210–223.
- Lomas M. W. and Moran S. B. (2011). Evidence for aggregation and export of cyanobacteria and nano-eukaryotes from the Sargasso Sea euphotic zone. *Biogeosciences*, 8(1):203–216.
- López-García P., López-López A., Moreira D., and Rodríguez-Valera F. (2001). Diversity of free-living prokaryotes from a deep-sea site at the Antarctic Polar Front. *FEMS Microbiology Ecology*, 36(2-3): 193–202.
- Ludwig W., Strunk O., Westram R., Richter L., Meier H., Yadhukumar , Buchner A., Lai T., Steppi S., Jobb G., Förster W., Brettske I., Gerber S., Ginhart A. W., Gross O., Grumann S., Hermann S., Jost R., König A., Liss T., Lüssmann R., May M., Nonhoff B., Reichel B., Strehlow R., Stamatakis A., Stuckmann N., Vilbig A., Lenke M., Ludwig T., Bode A., and Schleifer K.-H. (2004). ARB: a software environment for sequence data. *Nucleic Acids Research*, 32(4):1363–1371.
- Malmstrom R. R., Cottrell M. T., Elifantz H., and Kirchman D. L. (2005). Biomass production and assimilation of dissolved organic matter by SAR11 bacteria in the Northwest Atlantic Ocean. *Applied and Environmental Microbiology*, 71(6):2979–2986.
- Marchant H. J., Davidson A. T., and Wright S. W. (1987). The distribution and abundance of chroococloid cyanobacteria in the Southern Ocean. *Proc. NIPR Symp. Polar Biol*, 1:1–9.
- Martiny J. B. H., Bohannan B. J. M., Brown J. H., Colwell R. K., Fuhrman J. A., Green J. L., Horner-Devine M. C., Kane M., Krumins J. A., Kuske C. R., Morin P. J., Naeem S., Ovreas L., Reysenbach A.-L., Smith V. H., and Staley J. T. (2006). Microbial biogeography: putting microorganisms on the map. *Nature Reviews Microbiology*, 4(2):102–112.
- Mary I., Heywood J. L., Fuchs B. M., Amann R., Tarran G. A., Burkhill P. H., and Zubkov M. V. (2006). SAR11 dominance among metabolically active low nucleic acid bacterioplankton in surface waters along an Atlantic meridional transect. *Aquatic Microbial Ecology*, 45(2):107–113.
- Massana R., Taylor L. T., Murray A. E., Wu K. Y., Jeffrey W. H., and DeLong E. F. (1998). Vertical Distribution and Temporal Variation of Marine Planktonic Archaea in the Gerlache Strait, Antarctica, During Early Spring. *Limnology and ...*, 43(4):607–617.
- Massana R., DeLong E. F., and Pedrós-Alio C. (2000). A Few Cosmopolitan Phylotypes Dominate Planktonic Archaeal Assemblages in Widely Different Oceanic Provinces. *Applied and Environmental Microbiology*, 66(5):1777–1787.
- Mayali X., Franks P. J. S., and Azam F. (2008). Cultivation and Ecosystem Role of a Marine *Roseobacter* Clade-Affiliated Cluster Bacterium. *Applied and Environmental Microbiology*, 74(9):2595–2603.
- Mazloff M. R., Heimbach P., and Wunsch C. (2010). An eddy-permitting Southern Ocean state estimate. *Journal of physical oceanography*, 40:880–899.
- Merbt S. N., Stahl D. A., Casamayor E. O., Martí E., Nicol G. W., and Prosser J. I. (2012). Differential photoinhibition of bacterial and archaeal ammonia oxidation. *FEMS Microbiology Letters*, 327(1): 41–46.
- Methé B. A., Nelson K. E., Deming J. W., Momen B., Melamud E., Zhang X., Moult J., Madupu R., Nelson W. C., Dodson R. J., Methe B. A., Nelson K. E., Deming J. W., Momen B., Melamud E., Zhang X., Moult J., Madupu R., Nelson W. C., Dodson R. J., Brinkac L. M., Daugherty S. C., Durkin A. S., DeBoy R. T., Kolonay J. F., Sullivan S. A., Zhou L., Davidsen T. M., Wu M., Huston A. L., Lewis M., Weaver B., Weidman J. F., Khouri H., Utterback T. R., Feldblyum T. V., and Fraser C. M.

- (2005). The psychrophilic lifestyle as revealed by the genome sequence of *Colwellia psychrerythraea* 34H through genomic and proteomic analyses. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 102(31):10913–10918.
- Meyer B. and Kuever J. (2007). Molecular Analysis of the Diversity of Sulfate-Reducing and Sulfur-Oxidizing Prokaryotes in the Environment, Using *aprA* as Functional Marker Gene. *Applied and Environmental Microbiology*, 73(23):7664–7679.
- Mikaloff Fletcher S. E., Gruber N., Jacobson A. R., Doney S. C., Dutkiewicz S., Gerber M., Follows M., Joos F., Lindsay K., Menemenlis D., Mouchet A., Müller S. A., and Sarmiento J. L. (2006). Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean. *Global Biogeochemical Cycles*, 20(2):GB2002.
- Miller T. R. and Belas R. (2004). Dimethylsulfoniopropionate Metabolism by *Pfiesteria*-Associated *Roseobacter* spp. *Applied and Environmental Microbiology*, 70(6):3383–3391.
- Mira A., Ochman H., and Moran N. A. (2001). Deletional bias and the evolution of bacterial genomes. *Trends in genetics : TIG*, 17(10):589–596.
- Moore J. K., Abbott M. R., and Richman J. G. (1999). Location and dynamics of the Antarctic Polar Front from satellite sea surface temperature data. *Journal of Geophysical Research*, 104:3052–3073.
- Moran M. A., Belas R., Schell M. A., González J. M., Sun F., Sun S., Binder B. J., Edmonds J., Ye W., Orcutt B., Howard E. C., Meile C., Palefsky W., Goesmann A., Ren Q., Paulsen I., Ulrich L. E., Thompson L. S., Saunders E., and Buchan A. (2007). Ecological Genomics of Marine Roseobacters. *Applied and Environmental Microbiology*, 73(14):4559–4569.
- Moran M. A., González J. M., and Kiene R. P. (2003). Linking a Bacterial Taxon to Sulfur Cycling in the Sea: Studies of the Marine Roseobacter Group. *Geomicrobiology Journal*, 20(4):375–388.
- Moran M. A., Buchan A., González J. M., Heidelberg J. F., Whitman W. B., Kiene R. P., Henriksen J. R., King G. M., Belas R., Fuqua C., Brinkac L., Lewis M., Johri S., Weaver B., Pai G., Eisen J. A., Rahe E., Sheldon W. M., Ye W., Miller T. R., Carlton J., Rasko D. A., Paulsen I. T., Ren Q., Daugherty S. C., Deboy R. T., Dodson R. J., Durkin A. S., Madupu R., Nelson W. C., Sullivan S. A., Rosovitz M. J., Haft D. H., Selengut J., and Ward N. (2004). Genome sequence of *Silicibacter pomeroyi* reveals adaptations to the marine environment. *Nature*, 432(7019):910–913.
- Morris R. M., Rappé M. S., Connon S. A., Vergin K. L., Siebold W. A., Carlson C. A., and Giovannoni S. J. (2002). SAR11 clade dominates ocean surface bacterioplankton communities. *Nature*, 420(6917):806–810.
- Morris R. M., Longnecker K., and Giovannoni S. J. (2006). *Pirellula* and OM43 are among the dominant lineages identified in an Oregon coast diatom bloom. *Environmental Microbiology*, 8(8):1361–1370.
- Murray A. E. and Grzimski J. J. (2007). Diversity and genomics of Antarctic marine micro-organisms. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1488):2259–2271.
- Murray A. E., Wu K. Y., Moyer C. L., Karl D. M., and DeLong E. F. (1999). Evidence for circumpolar distribution of planktonic Archaea in the Southern Ocean. *Aquatic Microbial Ecology*, 18(3):263–273.
- Murray A. E. A., Preston C. M. C., Massana R. R., Taylor L. T. L., Blakis A. A., Wu K. K., and DeLong E. F. (1998). Seasonal and spatial variability of bacterial and archaeal assemblages in the coastal waters near Anvers Island, Antarctica. *Applied and Environmental Microbiology*, 64(7):2585–2595.
- Murray A. E., Peng V., Tyler C., and Wagh P. (2011). Marine bacterioplankton biomass, activity and community structure in the vicinity of Antarctic icebergs. *Deep Sea Research Part II: Topical Studies in Oceanography*, 58(11-12):1407–1421.
- Newton I. L. G., Woyke T., Auchtung T. A., Dilly G. F., Dutton R. J., Fisher M. C., Fontanez K. M., Lau E., Stewart F. J., Richardson P. M., Barry K. W., Saunders E., Dettter J. C., Wu D., Eisen J. A., and Cavanaugh C. M. (2007). The *Calyptogena magnifica* Chemoautotrophic Symbiont Genome. *Science*, 315(5814):998–1000.

- Ng C., DeMaere M. Z., Williams T. J., Lauro F. M., Raftery M., Gibson J. A., Andrews-Pfannkoch C., Lewis M., Hoffman J. M., Thomas T., and Cavicchioli R. (2010). Metaproteogenomic analysis of a dominant green sulfur bacterium from Ace Lake, Antarctica. *The ISME Journal*, 4(8):1002–1019.
- Nikrad M. P., Cottrell M. T., and Kirchman D. L. (2012). Abundance and Single-Cell Activity of Heterotrophic Bacterial Groups in the Western Arctic Ocean in Summer and Winter. *Applied and Environmental Microbiology*, 78(7):2402–2409.
- Obernosterer I., Catala P., Lebaron P., and West N. J. (2011). Distinct bacterial groups contribute to carbon cycling during a naturally iron fertilized phytoplankton bloom in the Southern Ocean. *Limnology and Oceanography*, 56(6):2391–2401.
- Oh H. M., Kwon K. K., Kang I., Kang S. G., Lee J. H., Kim S. J., and Cho J. C. (2010). Complete Genome Sequence of "Candidatus Puniceispirillum marinum" IMCC1322, a Representative of the SAR116 Clade in the Alphaproteobacteria. *Journal of Bacteriology*, 192(12):3240–3241.
- Oliver J. L., Barber R. T., Smith Jr W. O., and Ducklow H. W. (2004). The heterotrophic bacterial response during the Southern Ocean iron experiment (SOFeX). *Limnology and Oceanography*, 49(6): 2129–2140.
- Orsi A. H., Whitworth T., and Nowlin W. D. (1995). On the meridional extent and fronts of the Antarctic Circumpolar Current. *Deep Sea Research Part I: Oceanographic Research Papers*, 42(5):641–673.
- Orsi A. H., Johnson G. C., and Bullister J. L. (1999). Circulation, mixing, and production of Antarctic Bottom Water. *Progress in Oceanography*, 43(1):55–109.
- O'Sullivan L. A., Fuller K. E., Thomas E. M., Turley C. M., Fry J. C., and Weightman A. J. (2004). Distribution and culturability of the uncultivated 'AGG58 cluster' of the *Bacteroidetes* phylum in aquatic environments. *FEMS Microbiology Ecology*, 47(3):359–370.
- Paris C. B., Helgers J., Sebille E.van, and Srinivasan A. (2013). Connectivity Modeling System: A probabilistic modeling tool for the multi-scale tracking of biotic and abiotic variability in the ocean. *Environmental Modelling and Software*, 42(C):47–54.
- Partensky F., Hess W. R., and Vaulot D. (1999). *Prochlorococcus*, a marine photosynthetic prokaryote of global significance. *Microbiology and Molecular Biology Reviews*, 63(1):106–127.
- Paul J. H., DeFlaun M. F., and Jeffrey W. H. (1988). Mechanisms of DNA utilization by estuarine microbial populations. *Applied and Environmental Microbiology*, 54(7):1682–1688.
- Pham V. D., Konstantinidis K. T., Palden T., and DeLong E. F. (2008). Phylogenetic analyses of ribosomal DNA-containing bacterioplankton genome fragments from a 4000 m vertical profile in the North Pacific Subtropical Gyre. *Environmental Microbiology*, 10(9):2313–2330.
- Pinhassi J., Sala M. M., Havskum H., Peters F., Guadayol Ò., Malits A., and Marrasé C. (2004). Changes in bacterioplankton composition under different phytoplankton regimens. *Applied and Environmental Microbiology*, 70(11):6753–6766.
- Piquet A. M. T., Bolhuis H., Meredith M. P., and Buma A. G. J. (2011). Shifts in coastal Antarctic marine microbial communities during and after melt water-related surface stratification. *FEMS Microbiology Ecology*, 76(3):413–427.
- Pollard R. T., Lucas M. I., and Read J. F. (2002). Physical controls on biogeochemical zonation in the Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 49(16):3289–3305.
- Pommier T., Canbäck B., Riemann L., Boström K. H., Simu K., Lundberg P., Tunlid A., and Hagström Å. (2007). Global patterns of diversity and community structure in marine bacterioplankton. *Molecular Ecology*, 16(4):867–880.
- Poorvin L., Rinta-Kanto J. M., Hutchins D. A., and Wilhelm S. W. (2004). Viral release of iron and its bioavailability to marine plankton. *Limnology and Oceanography*, 49(5):1734–1741.

- Powell L. M., Bowman J. P., Skerratt J. H., Franzmann P. D., and Burton H. R. (2005). Ecology of a novel *Synechococcus* clade occurring in dense populations in saline Antarctic lakes. *Marine Ecology Progress Series*, 291(28 April):65–80.
- Preston C. M., Wu K. Y., Molinski T. F., and DeLong E. F. (1996). A psychrophilic crenarchaeon inhabits a marine sponge: *Cenarchaeum symbiosum* gen. nov., sp. nov. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 93(13):6241–6246.
- Qin J., Li R., Raes J., Arumugam M., Burgdorf K. S., Manichanh C., Nielsen T., Pons N., Levenez F., Yamada T., Mende D. R., Li J., Xu J., Li S., Li D., Cao J., Wang B., Liang H., Zheng H., Xie Y., Tap J., Lepage P., Bertalan M., Batto J.-M., Hansen T., Le Paslier D., Linneberg A., Nielsen H. B., Pelletier E., Renault P., Sicheritz-Ponten T., Turner K., Zhu H., Yu C., Li S., Jian M., Zhou Y., Li Y., Zhang X., Li S., Qin N., Yang H., Wang J., Brunak S., Doré J., Guarner F., Kristiansen K., Pedersen O., Parkhill J., Weissenbach J., MetaHIT Consortium , Bork P., Ehrlich S. D., and Wang J. (2010). A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*, 464(7285):59–65.
- Quast C., Pruesse E., Yilmaz P., Gerken J., Schneemann T., Yarza P., Peplies J., and Glöckner F. O. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, 41(Database issue):D590–6.
- Ramette A. and Tiedje J. M. (2007). Multiscale responses of microbial life to spatial distance and environmental heterogeneity in a patchy ecosystem. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 104(8):2761–2766.
- Rappé M. S., Connon S. A., Vergin K. L., and Giovannoni S. J. (2002). Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature*, 418(6898):630–633.
- Rath J., Wu K. Y., Herndl G. J., and DeLong E. F. (1998). High phylogenetic diversity in a marine-snow-associated bacterial assemblage. *Aquatic Microbial Ecology*, 14(3):261–269.
- Reisch C. R., Stoudemayer M. J., Varaljay V. A., Amster I. J., Moran M. A., and Whitman W. B. (2011). Novel pathway for assimilation of dimethylsulphoniopropionate widespread in marine bacteria. *Nature*, 473(7346):208–211.
- Rosenberg M. and Rintoul S. R. Aurora Australis Marine Science Cruise AU1203 – Oceanographic Field Measurements and Analysis. Technical report, 2012.
- Rosenberg M. S. and Anderson C. D. (2011). PASSaGE: pattern analysis, spatial statistics and geographic exegesis. Version 2. *Methods in Ecology and Evolution*, 2(3):229–232.
- Rusch D. B., Halpern A. L., Sutton G., Heidelberg K. B., Williamson S., Yooseph S., Wu D., Eisen J. A., Hoffman J. M., Remington K., Beeson K., Tran B., Smith H., Baden-Tillson H., Stewart C., Thorpe J., Freeman J., Andrews-Pfannkoch C., Venter J. E., Li K., Kravitz S., Heidelberg J. F., Utterback T., Rogers Y.-H., Falcón L. I., Souza V., Bonilla-Rosso G., Eguiarte L. E., Karl D. M., Sathyendranath S., Platt T., Bermingham E., Gallardo V., Tamayo-Castillo G., Ferrari M. R., Strausberg R. L., Nealson K., Friedman R., Frazier M., and Venter J. C. (2007). The Sorcerer II Global Ocean Sampling expedition: northwest Atlantic through eastern tropical Pacific. *PLoS Biology*, 5(3):e77–e77.
- Sabine C. L., Feely R. A., Gruber N., Key R. M., Lee K., Bullister J. L., Wanninkhof R., Wong C. S., Wallace D. W. R., Tilbrook B., Millero F. J., Peng T.-H., Kozyr A., Ono T., and Rios A. F. (2004). The Oceanic Sink for Anthropogenic CO₂. *Science*, 305(5682):367–371.
- Scanlan D. J., Ostrowski M., Mazard S., Dufresne A., Garczarek L., Hess W. R., Post A. F., Hagemann M., Paulsen I., and Partensky F. (2009). Ecological Genomics of Marine Picocyanobacteria. *Microbiology and Molecular Biology Reviews*, 73(2):249–299.
- Selje N. N., Simon M. M., and Brinkhoff T. T. (2004). A newly discovered *Roseobacter* cluster in temperate and polar oceans. *Nature*, 427(6973):445–448.
- Short C. M. and Suttle C. A. (2005). Nearly Identical Bacteriophage Structural Gene Sequences Are Widely Distributed in both Marine and Freshwater Environments. *Applied and Environmental Microbiology*, 71(1):480–486.

- Short S. M. and Suttle C. A. (2002). Sequence Analysis of Marine Virus Communities Reveals that Groups of Related Algal Viruses Are Widely Distributed in Nature. *Applied and Environmental Microbiology*, 68(3):1290–1296.
- Simon M., Glöckner F. O., and Amann R. (1999). Different community structure and temperature optima of heterotrophic picoplankton in various regions of the Southern Ocean. *Aquatic Microbial Ecology*, 18(3):275–284.
- Sinha V., Williams J., Meyerhöfer M., Riebesell U., Paulino A. I., and Larsen A. (2007). Air-sea fluxes of methanol, acetone, acetaldehyde, isoprene and DMS from a Norwegian fjord following a phytoplankton bloom in a mesocosm experiment. *Atmospheric Chemistry and Physics*, 7(3):739–755.
- Sokolov S. and Rintoul S. R. (2002). Structure of Southern Ocean fronts at 140°E. *Journal of Marine Systems*, 37(1):151–184.
- Sokolov S. and Rintoul S. R. (2009). Circumpolar structure and distribution of the Antarctic Circumpolar Current fronts: 1. Mean circumpolar paths. *Journal of Geophysical Research*, 114(C11):C11018.
- Sowell S. M., Wilhelm L. J., Norbeck A. D., Lipton M. S., Nicora C. D., Barofsky D. F., Carlson C. A., Smith R. D., and Giovannoni S. J. (2009). Transport functions dominate the SAR11 metaproteome at low-nutrient extremes in the Sargasso Sea. *The ISME Journal*, 3(1):93–105.
- Speer K., Rintoul S. R., and Sloyan B. (2000). The Diabatic Deacon Cell. *Journal of physical oceanography*, 30(12):3212–3222.
- Steindler L., Schwalbach M. S., Smith D. P., Chan F., and Giovannoni S. J. (2011). Energy Starved *Candidatus Pelagibacter Ubique* Substitutes Light-Mediated ATP Production for Endogenous Carbon Respiration. *PLoS ONE*, 6(5):e19725.
- Stingl U., Tripp H. J., and Giovannoni S. J. (2007). Improvements of high-throughput culturing yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast and the Bermuda Atlantic Time Series study site. *The ISME Journal*, 1:361–371.
-). The concept of taxon invariance in ecology: do diversity patterns vary with changes in taxonomic resolution? *Folia Geobotanica*, 43:329–344.
- Straza T. R. A., Ducklow H. W., Murray A. E., and Kirchman D. L. (2010). Abundance and single-cell activity of bacterial groups in Antarctic coastal waters. *Limnology and Oceanography*, 55(6):2526–2536.
- Strous M., Fuerst J. A., Kramer E. H. M., Logemann S., Muyzer G., Van De Pas-Schoonen K. T., Webb R., Kuenen J. G., and Jetten M. S. M. (1999). Missing lithotroph identified as new planctomycete. *Nature*, 400(6743):446–449.
- Strutton P. G., Griffiths F. B., Waters R. L., Wright S. W., and Bindoff N. L. (2000). Primary productivity off the coast of East Antarctica (80- 150°E): January to March 1996. *Deep Sea Research Part II: Topical Studies in Oceanography*, 47:2327–2362.
- Sul W. J., Oliver T. A., Ducklow H. W., Amaral-Zettler L. A., and Sogin M. L. (2013). Marine bacteria exhibit a bipolar distribution. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 110(6):2342–2347.
- Swan B. K., Martinez-Garcia M., Preston C. M., Sczyrba A., Woyke T., Lamy D., Reinthal T., Poulton N. J., Masland E. D. P., Gomez M. L., Sieracki M. E., DeLong E. F., Herndl G. J., and Stepanauskas R. (2011). Potential for Chemolithoautotrophy Among Ubiquitous Bacteria Lineages in the Dark Ocean. *Science*, 333(6047):1296–1300.
- Swingley W. D., Sadekar S., Mastrian S. D., Matthies H. J., Hao J., Ramos H., Acharya C. R., Conrad A. L., Taylor H. L., Dejesa L. C., Shah M. K., O'Huallachain M. E., Lince M. T., Blankenship R. E., Beatty J. T., and Touchman J. W. (2007). The Complete Genome Sequence of *Roseobacter denitrificans* Reveals a Mixotrophic Rather than Photosynthetic Metabolism. *Journal of Bacteriology*, 189(3):683–690.

- Teske A., Alm E., Regan J. M., Toze S., Rittmann B. E., and Stahl D. A. (1994). Evolutionary relationships among ammonia- and nitrite-oxidizing bacteria. *Journal of Bacteriology*, 176(21):6623–6630.
- Thomalla S. J., Waldron H. N., Lucas M. I., Read J. F., Ansorge I. J., and Pakhomov E. (2011). Phytoplankton distribution and nitrogen dynamics in the southwest Indian subtropical gyre and Southern Ocean waters. *Ocean Science*, 7(1):113–127.
- Thompson D. W. J. and Solomon S. (2002). Interpretation of Recent Southern Hemisphere Climate Change. *Science*, 296(5569):895–899.
- Topping J. N., Heywood J. L., Ward P., and Zubkov M. V. (2006). Bacterioplankton composition in the Scotia Sea, Antarctica, during the austral summer of 2003. *Aquatic Microbial Ecology*, 45(3):229–235.
- Tréguer P., Nelson D. M., Van Bennekom A. J., DeMaster D. J., Leynaert A., and Quéquier B. (1995). The silica balance in the world ocean: a reestimate. *Science*, 268(5209):375–379.
- Tripp H. J., Kitner J. B., Schwalbach M. S., Dacey J. W. H., Wilhelm L. J., and Giovannoni S. J. (2008). SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature*, 452(7188):741–744.
- Trull T., Rintoul S. R., Hadfield M., and Abraham E. R. (2001). Circulation and seasonal evolution of polar waters south of Australia: implications for iron fertilization of the Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, 48(11):2439–2466.
- Sebille E. van, Johns W. E., and Beal L. M. (2012). Does the vorticity flux from Agulhas rings control the zonal pathway of NADW across the South Atlantic? *Journal of Geophysical Research*, 117(C5):C05037.
- Venter J. C., Remington K., Heidelberg J. F., Halpern A. L., Rusch D., Eisen J. A., Wu D., Paulsen I., Nelson K. E., Nelson W., Fouts D. E., Levy S., Knap A. H., Lomas M. W., Nealson K., White O., Peterson J., Hoffman J., Parsons R., Baden-Tillson H., Pfannkoch C., Rogers Y.-H., and Smith H. O. (2004). Environmental Genome Shotgun Sequencing of the Sargasso Sea. *Science*, 304(5667):66–74.
- Vila-Costa M., Simó R., Harada H., Gasol J. M., Slezak D., and Kiene R. P. (2006). Dimethylsulfonio-propionate Uptake by Marine Phytoplankton. *Science*, 314(5799):652–654.
- Wagner-Döbler I. and Biebl H. (2006). Environmental Biology of the Marine *Roseobacter* Lineage. *Annual Review of Microbiology*, 60(1):255–280.
- Walker C. B., Torre J. R. de la, Klotz M. G., Urakawa H., Pinel N., Arp D. J., Brochier-Armanet C., Chain P., Chan P. P., Gollabgir A., Hemp J., Hügler M., Karr E. A., Könekke M., Shin M., Lawton T. J., Lowe T., Martens-Habbena W., Sayavedra-Soto L. A., Langf D., Sievert S. M., Rosenzweig A. C., Manning G., and Stahl D. A. (2010). *Nitrosopumilus maritimus* genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, 107(19):8818–8823.
- Walsh D. A., Zaikova E., Howes C. G., Song Y. C., Wright J. J., Tringe S. G., Tortell P. D., and Hallam S. J. (2009). Metagenome of a Versatile Chemolithoautotroph from Expanding Oceanic Dead Zones. *Science*, 326(5952):578–582.
- Ward P., Whitehouse M., Brandon M., Shreeve R., and Woold-Walker R. (2003). Mesozooplankton community structure across the Antarctic Circumpolar Current to the north of South Georgia: Southern Ocean. *Marine Biology*, 143(1):121–130.
- Weber T. S. and Deutsch C. (2010). Ocean nutrient ratios governed by plankton biogeography. *Nature*, 467(7315):550–554.
- Weinbauer M. G., Arrieta J. M., Griebler C., and Herndl G. J. (2009). Enhanced viral production and infection of bacterioplankton during an iron-induced phytoplankton bloom in the Southern Ocean. *Limnol. Oceanogr.*, 54(3):774–784.
- West N. J., Obernosterer I., Zemb O., and Lebaron P. (2008). Major differences of bacterial diversity and activity inside and outside of a natural iron-fertilized phytoplankton bloom in the Southern Ocean. *Environmental Microbiology*, 10(3):738–756.

- Whitworth T. (1980). Zonation and geostrophic flow of the Antarctic Circumpolar Current at Drake Passage. *Deep Sea Research Part I: Oceanographic Research Papers*, 27(7):497–507.
- Whitworth III T. and Nowlin Jr. W. D. (1987). Water masses and currents of the Southern Ocean at the Greenwich Meridian. *Journal of Geophysical Research*, 92(C6):6462–6476.
- Wilhelm S. W. and Suttle C. A. (1999). Viruses and nutrient cycles in the sea. *BioScience*, 49(10):781–788.
- Wilkins D., Lauro F. M., Williams T. J., DeMaere M. Z., Brown M. V., Hoffman J. M., Andrews-Pfannkoch C., McQuaid J. B., Riddle M. J., Rintoul S. R., and Cavicchioli R. (2012). Biogeographic partitioning of Southern Ocean microorganisms revealed by metagenomics. *Environmental Microbiology*.
- Wilkins D., Yau S., Williams T. J., Allen M. A., Brown M. V., DeMaere M. Z., Lauro F. M., and Cavicchioli R. (2012). Key microbial drivers in Antarctic aquatic environments. *FEMS Microbiology Reviews*.
- Williams G. D., Nicol S., Aoki S., Meijers A. J. S., Bindoff N. L., Iijima Y., Marsland S. J., and Klocker A. (2010). Surface oceanography of BROKE-West, along the Antarctic margin of the south-west Indian Ocean (30–80°E). *Deep Sea Research Part II: Topical Studies in Oceanography*, 57(9-10):738–757.
- Williams T. J., Lauro F. M., Ertan H., Burg D. W., Poljak A., Raftery M. J., and Cavicchioli R. (2011). Defining the response of a microorganism to temperatures that span its complete growth temperature range (-2 °C to 28 °C) using multiplex quantitative proteomics. *Environmental Microbiology*, 13 (8):2186–2203.
- Williams T. J., Long E., Evans F., DeMaere M. Z., Lauro F. M., Raftery M. J., Ducklow H., Grzymski J. J., Murray A. E., and Cavicchioli R. (2012). A metaproteomic assessment of winter and summer bacterioplankton from Antarctic Peninsula coastal surface waters. *The ISME Journal*, 6(10):1883–1900.
- Williams T. J., Wilkins D., Long E., Evans F., DeMaere M. Z., Raftery M. J., and Cavicchioli R. (2012). The role of planktonic Flavobacteria in processing algal organic matter in coastal East Antarctica revealed using metagenomics and metaproteomics. *Environmental Microbiology*.
- Wright T. D., Vergin K. L., Boyd P. W., and Giovannoni S. J. (1997). A novel āLČ-subdivision proteobacterial lineage from the lower ocean surface layer. *Applied and Environmental Microbiology*, 63 (4):1441–1448.
- Wunsch C. and Heimbach P. (2007). Practical global oceanic state estimation. *Physica D: Nonlinear Phenomena*, 230(1):197–208.
- Ye Y. and Doak T. G. (2009). A parsimony approach to biological pathway reconstruction/inference for genomes and metagenomes. *PLoS Computational Biology*, 5(8):e1000465.
- Yoon J., Yasumoto-Hirose M., Katsuta A., Sekiguchi H., Matsuda S., Kasai H., and Yokota A. (2007). *Coraliomargarita akajimensis* gen. nov., sp. nov., a novel member of the phylum 'Verrucomicrobia' isolated from seawater in Japan. *International Journal of Systematic and Evolutionary Microbiology*, 57(5): 959–963.
- Youssef N., Sheik C. S., Krumholz L. R., Najar F. Z., Roe B. A., and Elshahed M. S. (2009). Comparison of species richness estimates obtained using nearly complete fragments and simulated pyrosequencing-generated fragments in 16S rRNA gene-based environmental surveys. *Applied and Environmental Microbiology*.
- Zaballos M., López-López A., Ovreas L., Bartual S. G., D'Auria G., Alba J. C., Legault B., Pushker R., Daae F. L., and Rodriguez-Valera F. (2006). Comparison of prokaryotic diversity at offshore oceanic locations reveals a different microbiota in the Mediterranean Sea. *FEMS Microbiology Ecology*, 56(3): 389–405.
- Zhang R., Liu B., Lau S. C. K., Ki J.-S., and Qian P.-Y. (2007). Particle-attached and free-living bacterial communities in a contrasting marine environment: Victoria Harbor, Hong Kong. *FEMS Microbiology Ecology*, 61(3):496–508.

Zubkov M. V., Sleigh M. A., Tarran G. A., Burkill P. H., and Leakey R. J. G. (1998). Picoplanktonic community structure on an Atlantic transect from 50°N to 50°S. *Deep Sea Research Part I: Oceanographic Research Papers*, 45(8):1339–1355.