## My PhD thesis

David Wilkins

September 20, 2012

## **Contents**

1	Intr	roduction	3
	1.1	Some test text to preview layout etc	3
	1.2	Microbial ecology of the Southern Ocean	3
	1.3	Oceanography of the Southern Ocean	3
		1.3.1 Water masses and fronts	3
		1.3.2 Effect of climate change	3
	1.4	Role of the Polar Front in biogeography	3
	1.5	Project questions and hypotheses	3
2	The	Polar Front as a major biogeographic boundary in the Southern Ocean	5
	2.1	Summary	5
	2.2	Introduction	5
	2.3	Methods	6
		2.3.1 Sampling	6
		2.3.2 Metagenomic sequencing	6
		2.3.3 Phylogenetic analysis of metagenomic data	6
		2.3.4 Functional analysis of metagenomic data	6
	2.4	Results	6
		2.4.1 Metagenomic sequencing	6
		2.4.2 Phylogenetic analysis of metagenomic data	6
		2.4.3 Functional analysis of metagenomic data	6
	2.5	Discussion	6
	2.6	Conclusions	6
3	Mes	so-scale biogeographic drivers of planktonic diversity	7
4	Con	nclusions	(

# **List of Figures**

## **List of Tables**

# Acknowledgements

## **Abstract**

### Introduction

### 1.1 Some test text to preview layout *etc*.

The Roseobacter clade is an abundant and ecologically significant group of marine bacteria, found at high (> 15%) abundance in most marine surface environments (Ano (2005) and references therein). Unlike some other major proteobacterial groups which are strongly associated with a particular ecological niche (e.g. the SAR11 clade), roseobacters have diverse metabolic abilities, with members capable (for example) of aerobic anoxygenic phototrophy (Biebl, 2005; Béjà *et al.*, 2002), degradation of dimethylsulfoniopropionate (DMSP) by at least two pathways (Moran *et al.*, 2007; ?), carbon monoxide oxidation (King, 2003) and heterotrophic utilisation of a broad range of substrates (reviewed in (Brinkhoff *et al.*, 2008)). Roseobacters are found in the planktonic fraction as well as in commensal association with phytoplankton and metazoans (reviewed in Ano (2005)).

- 1.2 Microbial ecology of the Southern Ocean
- 1.3 Oceanography of the Southern Ocean
- 1.3.1 Water masses and fronts
- 1.3.2 Effect of climate change
- 1.4 Role of the Polar Front in biogeography
- 1.5 Project questions and hypotheses

# The Polar Front as a major biogeographic boundary in the Southern Ocean

### 2.1 Summary

#### 2.2 Introduction

### 2.3 Methods

- 2.3.1 Sampling
- 2.3.2 Metagenomic sequencing
- 2.3.3 Phylogenetic analysis of metagenomic data
- 2.3.4 Functional analysis of metagenomic data

### 2.4 Results

- 2.4.1 Metagenomic sequencing
- 2.4.2 Phylogenetic analysis of metagenomic data
- 2.4.3 Functional analysis of metagenomic data
- 2.5 Discussion
- 2.6 Conclusions

Meso-scale biogeographic drivers of planktonic diversity

# **Conclusions**

## **Bibliography**

- Buchan, Moran 2005 Overview of the Marine Roseobacter Lineage. pages 1-13, September 2005.
- Béjà O., Suzuki M. T., Heidelberg J. F., Nelson W. C., Preston C. M., Hamada T., Eisen J. A., Fraser C. M., and DeLong E. F. Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature*, 415(6872):630–633, February 2002.
- Biebl H. Dinoroseobacter shibae gen. nov., sp. nov., a new aerobic phototrophic bacterium isolated from dinoflagellates. *INTERNATIONAL JOURNAL OF SYSTEMATIC AND EVOLUTIONARY MI-CROBIOLOGY*, 55(3):1089–1096, May 2005.
- Brinkhoff T., Giebel H.-A., and Simon M. Diversity, ecology, and genomics of the Roseobacter clade: a short overview. *Archives of Microbiology*, 189(6):531–539, February 2008.
- King G. M. Molecular and culture-based analyses of aerobic carbon monoxide oxidizer diversity. *Applied and Environmental Microbiology*, 69(12):7257–7265, November 2003.
- Lauro F. M., Demaere M. Z., Yau S., Brown M. V., Ng C., Wilkins D., Raftery M. J., Gibson J. A., Andrews-Pfannkoch C., Lewis M., Hoffman J. M., Thomas T., and Cavicchioli R. An integrative study of a meromictic lake ecosystem in Antarctica. *The ISME journal*, 5(5):879–895, December 2010.
- Moran M. A., Belas R., Schell M. A., Gonzalez J. M., Sun F., Sun S., Binder B. J., Edmonds J., Ye W., Orcutt B., Howard E. C., Meile C., Palefsky W., Goesmann A., Ren Q., Paulsen I., Ulrich L. E., Thompson L. S., Saunders E., and Buchan A. Ecological Genomics of Marine Roseobacters. *Applied and Environmental Microbiology*, 73(14):4559–4569, July 2007.
- Weber T. S. and Deutsch C. Ocean nutrient ratios governed by plankton biogeography. *Nature*, 467 (7315):550–554, September 2010.
- Ye Y. and Doak T. G. A Parsimony Approach to Biological Pathway Reconstruction/Inference for Genomes and Metagenomes. *PLoS Computational Biology*, 5(8):e1000465, August 2009.